

**This Page Is Inserted by IFW Operations
and is not a part of the Official Record**

BEST AVAILABLE IMAGES

**Defective images within this document are accurate representations of
the original documents submitted by the applicant.**

Defects in the images may include (but are not limited to):

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
18.10.2000 Bulletin 2000/42

(51) Int. Cl.⁷: **C07D 239/46**, C07D 401/04,
C07D 401/12, C07D 409/04,
A61K 31/505

(21) Application number: **92307389.4**

(22) Date of filing: **12.08.1992**

(54) **Pyrimidinyl acetamides as elastase inhibitors**

Pyrimidinyl-Essigsäureamide als Elastase Inhibitoren

Pyrimidinyl-acétamides comme inhibiteurs de l'élastase

(84) Designated Contracting States:
AT BE CH DE DK FR GB IT LI LU MC NL PT SE

(30) Priority: **15.08.1991 GB 9117641**
16.04.1992 GB 9208378
08.07.1992 GB 9214447

(43) Date of publication of application:
24.02.1993 Bulletin 1993/08

(73) Proprietor: **ZENECA LIMITED**
London W1Y 6LN (GB)

(72) Inventors:
• **Bernstein, Peter Robert**
Wilmington DE 19897 (US)
• **Edwards, Philip Duke**
Wilmington DE 19897 (US)
• **Shaw, Andrew**
Wilmington DE 19897 (US)

- **Thomas, Royston Martin**
Wilmington DE 19897 (US)
- **Veale, Chris Allan**
Wilmington DE 19897 (US)
- **Warner, Peter**
Macclesfield, Cheshire, GB SK10 4TG (GB)
- **Wolanin, Donald John**
Wilmington DE 19897 (US)

(74) Representative:
Giles, David Eric et al
AstraZeneca PLC
Global Intellectual Property
Mereside
Alderley Park
Macclesfield, Cheshire SK10 4TG (GB)

(56) References cited:
US-A- 4 910 190

EP 0 528 633 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention relates to certain substituted heterocycles, in particular, certain 1-pyrimidinylacetamide compounds, which are inhibitors of human leukocyte elastase (HLE), also known as human neutrophil elastase (HNE), making them useful whenever such inhibition is desired, such as for research tools in pharmacological, diagnostic and related studies and in the treatment of diseases in mammals in which HLE is implicated. For example, HLE has been implicated in the pathogenesis of acute respiratory distress syndrome (ARDS), rheumatoid arthritis, atherosclerosis, pulmonary emphysema, and other inflammatory disorders, including airway inflammatory diseases characterized by increased and abnormal airway secretion such as chronic bronchitis and cystic fibrosis. Also, HLE has been implicated in certain vascular diseases and related conditions (and their therapy) in which neutrophil participation is involved or implicated, for example, in hemorrhage associated with acute non-lymphocytic leukemia, as well as in reperfusion injury associated with, for example, myocardial ischaemia and related conditions associated with coronary artery disease such as angina and infarction, cerebrovascular ischaemia such as transient ischaemic attack and stroke, peripheral occlusive vascular disease such as intermittent claudication and critical limb ischaemia, venous insufficiency such as venous hypertension, varicose veins and venous ulceration, as well as impaired reperfusion states such as those associated with reconstructive vascular surgery, thrombolysis and angioplasty. The invention also includes intermediates useful in the synthesis of these heterocyclic amides, processes for preparing the heterocyclic amides, pharmaceutical compositions containing such heterocyclic amides and methods for their use.

[0002] In U.S. Patent 4,910,190, of 20 March 1990, assigned to ICI Americas Inc., there is disclosed a series of peptidoyl trifluoromethane derivatives which are HLE inhibitors. Disclosed herein is a series of substituted 2-(6-oxo-1,6-dihydro-1-pyrimidinyl)-N-[3,3,3-trifluoro-1-(lower alkyl)-2-oxopropyl]acetamide derivatives, which unexpectedly possess inhibitory properties against HLE, which provides the basis for the present invention.

[0003] According to the invention there is provided a Compound of the invention which is a compound of formula I (formula set out, together with other formulae referred to by Roman numerals, following the Examples) wherein:

R⁰ is (1-5C)alkyl;

R is hydrogen; or

R is an acyl group of formula A.X.CO- in which A.X-, taken together, is hydrogen, trifluoromethyl, 2,2,2-trifluoroethoxy, amino, methoxyamino, 2,2,2-trifluoroethylamino, RbRcN.O-, RaOCONH-, R¹SO₂NH-, RaOCO-, RbRcNCO- or RaCO-; or

R is an acyl group of formula A.X.CJ- in which

J is oxygen or sulfur;

X is a direct bond, imino, oxy or thio; and

A is as defined below or

A is tetrahydropyran-4-yl, 1-methylpiperid-4-yl, or 5-methyl-1,3-dioxacyclohex-5-ylmethyl; or

R is a sulfonyl group of formula D.W.SO₂- in which D.W-, taken together, is hydroxy, amino, di(1-4C)alkylamino, 2,2,2-trifluoroethylamino, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl or trifluoromethyl; or

W is a direct bond, imino, carbonylimino, oxycarbonylimino or iminocarbonylimino; and

D is as defined below; or

R is an alkyl, aryl or heteroaryl group G as defined below;

The group A, D or G is (1-6C)alkyl, (3-6C)cycloalkyl, (3-6C)cycloalkyl-(1-3C)alkyl, aryl, aryl(1-3C)alkyl, heteroaryl or heteroaryl(1-3C)alkyl wherein an aryl moiety is selected from phenyl, indenyl and naphthyl, and a heteroaryl moiety is selected from furyl, imidazolyl, tetrazolyl, pyridyl, pyridyl N-oxide, thienyl, pyrimidinyl, pyrimidinyl N-oxide, indolyl, quinolyl and quinolyl N-oxide, and wherein an aryl or heteroaryl moiety may bear one or more halogeno, nitro, methyl or trifluoromethyl groups and further wherein the group A, D or G may bear one or more substituents selected from a group consisting of hydroxy, (1-4C)alkoxy, acyloxy containing 1 to 5 carbon atoms, COORa, CONRbRc, COO(CH₂)₂NRdRf, cyano, SO₂R¹, CONRdSO₂R¹, NRdRf, NRgCHO, NRgCOR², NRgCOOR², NRhCQNRiRj, NRkSO₂R³, SO₂NRlRm, SO₂NRnCOR⁴ and P(O)(ORa)₂ in which

Q is oxygen or sulfur;

Ra-Rn are independently hydrogen, benzyl or (1-4C)alkyl; or, independently, a group NRbRc, NRdRf, NRiRj or NRlRm is a cyclic radical selected from a group consisting of 1-pyrrolidinyl, piperidino, morpholino or 1-piperazinyl which may bear a (1-4C)alkyl substituent at the 4-position; or, independently, a group NRdRf is a cyclic radical selected from a group consisting of 2-pyrrolidinon-1-yl, succinimido, oxazolidin-2-on-3-yl, 2-benzoxazolinon-3-yl, phthalimido and cis-hexahydrophthalimido; and

R¹-R⁴ are independently trifluoromethyl, (1-6C)alkyl, (3-6C)cycloalkyl, aryl or heteroaryl wherein an aryl moiety is selected from phenyl, indenyl and naphthyl, and a heteroaryl moiety is selected from furyl, imidazolyl, tetrazolyl, pyridyl, pyridyl N-oxide, thienyl, pyrimidinyl, pyrimidinyl N-oxide, indolyl, quinolyl and quinolyl N-oxide, and wherein an aryl or heteroaryl moiety may bear one or more substituents selected from a group consisting of (1-4C)alkyl,

hydroxy, (1-4C)alkoxy, halogeno or trifluoromethyl;

R^6 is (1-5C)alkyl which has no tertiary carbon, (3-7C)cycloalkyl, aryl or heteroaryl, wherein an aryl moiety is selected from phenyl, indenyl and naphthyl, and a heteroaryl moiety is selected from furyl, imidazolyl, tetrazolyl, pyridyl, pyridyl N-oxide, thienyl, pyrimidinyl, pyrimidinyl N-oxide, indolyl, quinolyl and quinolyl N-oxide, and wherein an aryl or heteroaryl moiety independently may bear one or more of the substituents defined for the group A or an aryl or heteroaryl moiety thereof; and

provided that no aliphatic carbon is bonded to more than one nitrogen or oxygen, except as part of a cyclic ketal or where the nitrogen bears a carbonyl group; or, for a compound of formula I which is acidic or basic, a pharmaceutically acceptable salt thereof.

[0004] In this specification, the following definitions are used, unless otherwise described: Halogeno is fluoro, chloro, bromo or iodo. Alkyl, alkoxy, etc. denote both straight and branched groups; but reference to an individual radical such "propyl" embraces only the straight chain ("normal") radical, a branched chain isomer such as "isopropyl" being specifically referred to. It will be appreciated that, owing to the asymmetrically substituted carbon atom at the chiral center indicated by *** in formula I, a compound of formula I may exist in, and be isolated in, optically active and racemic forms. If a compound of formula I contains an additional chiral element, such compound of formula I may exist in, and be isolated in, the form of a diastereomeric mixture or as a single diastereomer. It is to be understood that the present invention encompasses a compound of formula I as a mixture of diastereomers, as well as in the form of an individual diastereomer, and that the present invention encompasses a compound of formula I as a mixture of enantiomers, as well as in the form of an individual enantiomer. When R^0 is isopropyl, a compound of formula I may be viewed as an alanyl trifluoromethane derivative. In general, a compound of formula I having the (S)-configuration at the chiral center indicated by ***, which corresponds to the L-alanyl configuration, is preferred. Accordingly, it may be preferred to use the compound of formula I in a form which is characterized as containing, for example, at least 95%, 98% or 99% enantiomeric excess (ee) of the (S)-form. However, owing to the interconvertibility of the (S)-isomer and the (R)-isomer by the facile epimerization of the chiral center indicated by *** in formula I, it may be preferred to utilize a compound of formula I as a mixture of the (S)- and (R)-isomers at the center indicated by *** in formula I.

[0005] As will be appreciated by those skilled in the art, a trifluoromethyl ketone of formula I can exist as a solvate, particularly a hydrate; and such a solvate of a compound of formula I is encompassed by the present invention.

[0006] A compound of formula I may exhibit polymorphism. The compound may form solvates in addition to a ketone solvate mentioned above. A compound may exist in more than one tautomeric form. It is to be understood, therefore, that the present invention encompasses any racemic or optically-active form, any polymorphic form, any tautomer or any solvate, or any mixture thereof, which form possesses inhibitory properties against HLE, it being well known in the art how to prepare optically active forms (for example, by resolution of the racemic form or by synthesis from optically-active starting materials) and how to determine the inhibitory properties against HLE by the standard tests described hereinafter.

[0007] It is preferred that the radicals R^0 , R and R^6 not contain nor introduce an additional element of chirality into the molecule beyond the chiral center indicated by *** in formula I.

[0008] Particular values are listed below for radicals, substituents and ranges for illustration only and they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

[0009] A particular value for R^0 is methyl, ethyl, propyl, isopropyl or isobutyl.

[0010] A particular value for W is a direct bond or imino.

[0011] A particular value for G is (1-3C)alkyl, aryl(1-3C)alkyl or heteroaryl(1-2C)alkyl which may bear one or more substituents as defined above for G or a part thereof.

[0012] A particular value of (1-6C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, 3-methylbutyl, 1-ethylpropyl, hexyl or 4-methylpentyl. A particular value of (3-6C)cycloalkyl is cyclopropyl, cyclopentyl or cyclohexyl. A particular value for the (1-3C)alkyl portion of (3-6C)cycloalkyl-(1-3C)alkyl, aryl(1-3C)alkyl or heteroaryl(1-3C)alkyl is methylene, ethylene or trimethylene. A particular value for (1-4C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl or t-butyl. A particular value for acyloxy containing 1 to 5 carbon atoms is acetoxy. A particular value for (1-4C)alkoxy is methoxy, ethoxy, propoxy, isopropoxy or t-butoxy. A particular value for halogeno is bromo, chloro or fluoro.

[0013] A particular value for A.X-, taken together, is 2,2,2-trifluoroethoxy. A particular value for COORa is carboxy or methoxycarbonyl. A particular value for CONRbRc is carbamoyl or N,N-dimethylcarbamoyl. A particular value for NR_gCOR² is trifluoroacetyl amino. A particular value of CONRdSO₂R¹ is N-phenylsulfonylcarbamoyl or N-(4-chlorophenylsulfonyl)carbamoyl. A particular value for A.X- is tris(hydroxymethyl)methylamino, tris(acetoxyethyl)methylamino or 2,2-bis(hydroxymethyl)propoxy.

[0014] A particular value for D.W-, taken together, is 2,2,2-trifluoroethylamino or 3,3,3-trifluoropropyl.

[0015] A particular value for R^6 is, for example, isopropyl, cyclopentyl, cyclohexyl, phenyl, furyl, thienyl or pyridyl in which a phenyl or heteroaryl may bear one or two substituents as defined above.

[0016] A more particular value for R^0 is isopropyl. A more particular value for A.X-, taken together, is 2,2,2-trifluor-

oethoxy. A more particular value for J is oxygen. A more particular value for X is a direct bond, imino or oxy. A more particular value for A is methyl, ethyl, phenyl, benzyl, phenethyl, pyridyl, thienyl, 5-tetrazolyl, thiazolyl, pyridylmethyl, thenyl, 5-tetrazolylmethyl, 2-(pyridyl)ethyl, 2-(thienyl)ethyl or 2-(thiazolyl)ethyl wherein the phenyl or heteroaryl group may bear one or two halogeno or methyl groups and further wherein the group A may bear a substituent selected from hydroxy, methoxy, *t*-butoxy, acetoxy, pivaloyloxy, carboxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl, dimethylcarbamoyl, 2-(dimethylamino)ethoxycarbonyl, cyano, methylsulfonyl, phenylsulfonyl, N-methylsulfonylcarbamoyl, N-phenylsulfonylcarbamoyl, amino, dimethylamino, oxazolidin-2-on-3-yl, acetylamino, trifluoroacetylamino, ureldo, methylsulfonyl, sulfamoyl, dimethylphosphoryl or diethylphosphoryl. A more particular value for D.W., taken together, is 2,2,2-trifluoroethylamino or 3,3,3-trifluoropropyl. A more particular value for D is methyl, ethyl, isopropyl, *tert*-butyl, cyclohexyl, phenyl, benzyl, phenethyl, pyridyl, thienyl, 5-tetrazolyl, thiazolyl, quinolonyl, pyridylmethyl, thenyl, 5-tetrazolylmethyl, 2-(pyridyl)ethyl, 2-(thienyl)ethyl or 2-(thiazolyl)ethyl wherein the phenyl or heteroaryl group may bear one or two halogeno or methyl groups and further wherein the group D may bear a substituent selected from hydroxy, methoxy, *t*-butoxy, acetoxy, pivaloyloxy, carboxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl, dimethylcarbamoyl, 2-(dimethylamino)ethoxycarbonyl, cyano, methylsulfonyl, phenylsulfonyl, N-methylsulfonylcarbamoyl, N-phenylsulfonylcarbamoyl, N-(4-chlorophenylsulfonyl)carbamoyl, methylsulfonylamino, amino, dimethylamino, oxazolidin-2-on-3-yl, acetylamino, trifluoroacetylamino, ureldo, methylsulfonyl, sulfamoyl, dimethylphosphoryl or diethylphosphoryl. A more particular value for G is methyl, ethyl, benzyl, phenethyl, pyridyl, pyridylmethyl, thenyl, 5-tetrazolylmethyl, or 2-(pyridyl)ethyl, wherein an alkyl carbon may bear an oxo group and wherein the phenyl or heteroaryl group may bear one or two halogeno or methyl groups and further wherein the group G may bear a substituent selected from hydroxy, methoxy, acetoxy, carboxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl, dimethylcarbamoyl, phenylcarbamoyl, pyridylcarbamoyl, methylsulfonylamino, amino, dimethylamino, acetylamino, nicotinoylamino, or trifluoroacetylamino.

[0017] A particular value for R is, for example, hydrogen, formyl, trifluoroacetyl, 2,2,2-trifluoroethoxycarbonyl, hydroxyoxalyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, 2-methoxyethoxycarbonyl, 4-fluorophenoxycarbonyl, 4-bromophenoxycarbonyl, 4-methoxyphenoxycarbonyl, benzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-pyridylmethoxycarbonyl, 3-methylpyrid-4-ylmethoxycarbonyl, 2,6-dimethylpyrid-4-ylmethoxy-carbonyl, 2-pyridylmethoxycarbonyl, 6-methylpyrid-2-ylmethoxycarbonyl, 2-dimethylaminoethoxycarbonyl, acetyl, carbamoylmethylaminocarbonyl, 4-(N-phenylsulfonylcarbamoyl)-phenylacetyl, methylthiocarbonyl, sulfo, aminosulfonyl, dimethylaminosulfonyl, 2,2,2-trifluoroethylaminosulfonyl, 3,3,3-trifluoroethylsulfonyl, trifluoromethylsulfonyl, methylsulfonyl (which may bear a methoxycarbonyl, carboxy or ethylsulfonyl substituent), methylaminosulfonyl, isopropylaminosulfonyl, butylsulfonyl, butylaminosulfonyl, *tert*-butylaminosulfonyl, cyclohexylaminosulfonyl, phenylsulfonyl (in which the phenyl may bear a chloro, nitro, amino, formylamino, acetylamino, trifluoroacetylamino, methoxy, carboxy, N-(4-chlorophenylsulfonyl)carbamoyl, or methylsulfonylamino substituent at the 3- or 4-position), anilino, pyridylsulfonyl, quinolonylsulfonyl, benzylsulfonyl (in which the phenyl ring may bear a nitro or amino substituent at the 3- or 4-position), pyridylmethylsulfonyl, 2-(pyridyl)ethylsulfonyl, benzylaminosulfonyl, methyl, ethyl, benzyl, phenethyl or pyridylmethyl.

[0018] A more particular value for R is, for example, hydrogen, formyl, trifluoroacetyl, 2,2,2-trifluoroethoxycarbonyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, 2-methoxyethoxycarbonyl, 4-bromophenoxycarbonyl, benzyloxycarbonyl, 2,6-dimethylpyrid-4-ylmethoxy-carbonyl, methylthiocarbonyl, *tert*-butylaminosulfonyl, 4-acetylaminophenylsulfonyl, 4-[N-(4-chlorophenylsulfonyl)carbamoyl]phenylsulfonyl, benzylsulfonyl, benzylaminosulfonyl or ethyl.

[0019] A particular group of compounds of formula I is one in which R⁰ and R have any of the values defined above and R⁶ is 2-furyl, 2-thienyl, 3-pyridyl or phenyl in which the phenyl may bear one or two halogeno, trifluoromethyl, methyl, hydroxy, methoxy, *tert*-butoxy, methoxycarbonyl or carboxy substituents; and, more particularly, R⁶ is phenyl, 4-fluorophenyl or 2-thienyl.

[0020] A more particular group of compounds of formula I is one in which R⁰ is isopropyl, R is hydrogen, formyl, 2,2,2-trifluoroethoxycarbonyl, isopropoxycarbonyl, methylthiocarbonyl or ethyl, and R⁶ is 2-furyl, 2-thienyl, 3-pyridyl or phenyl in which the phenyl may bear one or two halogeno, trifluoromethyl, methyl, hydroxy, methoxy, *tert*-butoxy, methoxycarbonyl or carboxy substituents; and, more particularly, R⁶ is phenyl, 4-fluorophenyl or 2-thienyl.

[0021] Specific compounds of formula I are described in the accompanying Examples. Of these, compounds of particular interest, along with their pharmaceutically acceptable salts, include those described in Examples 12, 15, 51, 82, 99, 100, 102, 106, 157 and 159, based upon their activity in *in vivo* tests.

[0022] A pharmaceutically acceptable salt of an acidic compound of formula I is one made with a base which affords a pharmaceutically acceptable cation, which includes alkali metal salts (especially lithium, sodium and potassium), alkaline earth metal salts (especially calcium and magnesium), aluminum salts and ammonium salts, as well as salts made from appropriate organic bases such as triethylamine, morpholine, piperidine and triethanol amine. A pharmaceutically acceptable salt of a basic compound of formula I includes an acid-addition salt made with an acid which provides a pharmaceutically acceptable anion, including for example, a strong acid such as hydrochloric, sulfuric or phosphoric acid.

[0023] A compound of formula I may be made by processes which include processes known in the chemical art for the production of structurally analogous heterocyclic and peptidic compounds. Such processes and intermediates for

the manufacture of a compound of formula I as defined above are provided as further features of the invention and are illustrated by the following procedures in which the meanings of generic radicals are as defined above:

- (A) Oxidizing a corresponding alcohol of formula II. If R is hydrogen or a group G, it will be recognized that protection of the pyridone 3-amino substituent prior to oxidation and removal of the protecting group after oxidation may be preferred or required if the amino group is not stable to the oxidation conditions employed. A convenient method is the use of excess dimethyl sulfoxide and a water soluble carbodiimide, with dichloroacetic acid as a catalyst, in an inert solvent such as toluene at about room temperature, for example as described in Example 1. Other methods which may be useful include the use of alkaline aqueous potassium permanganate solution; the use of oxalyl chloride, dimethyl sulfoxide and a tertiary amine; the use of acetic anhydride and dimethyl sulfoxide; the use of chromium trioxide pyridine complex in methylene chloride; and the use of a hypervalent iodine reagent, such as a periodinane, for example 1,1,1-triacetoxy-2,1-benzoxido-3(3H)-one with trifluoroacetic acid in dichloromethane.
- (B) For a compound of formula I which contains an N-H residue, removal by using a conventional method of the nitrogen protecting group of a corresponding compound bearing a conventional nitrogen protecting group to afford the compound of formula I which contains an amino N-H residue, particularly for a compound of formula I in which R is hydrogen, removal of a group from a corresponding compound of formula I, or for a compound of formula I in which R has a value of G, the removal of an activating/protecting group Rx from a corresponding compound of formula Vb. Rx is a group which protects and activates a primary amino group for substitution, such as for example benzyloxycarbonyl or trifluoroacetyl. Conventional methods include, for example, removal of a benzyloxycarbonyl group by hydrogenolysis, as described in Example 6; removal of a benzyloxycarbonyl by treatment with a strong acid, as described in Example 12, for example with trifluoromethanesulfonic acid in an inert solvent such as dichloromethane; and basic hydrolysis of a trifluoroacetyl group.
- (C) For a compound of formula I wherein R is an acyl group, acylation of a corresponding amine of formula I wherein R is hydrogen. Convenient methods include those described below for acylation of an amine of formula XIII, for example, when J is oxygen, the use of an activated carboxylic acid derivative, such as an acid halide, the use of a carboxylic acid and a coupling reagent, the use of an isocyanate for a compound wherein X is imino, and the use of a deactivated carbonic acid derivative, for example, carbonyldiimidazole, phosgene, diphosgene (trichloromethyl chloroformate) or triphosgene (bis(trichloromethyl) carbonate) with an alcohol of formula A.OH, a thiol of formula A.SH or an amine of formula A.NH₂ and a base, such as triethylamine or, when J is sulfur, the use of an activated thiocarboxylic acid derivative, such as a thioyl chloride or a lower alkyl ester of a dithioic acid, the use of a thioic acid and a coupling reagent, the use of an isothiocyanate for a compound wherein X is imino, and the use of a deactivated thiocarbonic acid derivative, for example, dimethyl trithiocarbonate, with an alcohol of formula A.OH, a thiol of formula A.SH or an amine of formula A.NH₂. In addition, for a compound of formula I in which R is an acyl group of formula A.X.CO- and X is oxy or imino, the acylation may be carried out by converting the corresponding amine of formula I in which R is hydrogen into its corresponding isocyanate, followed by reaction of the isocyanate with an alcohol of formula A.OH or an amine of formula A.NH₂, respectively, using a method similar to that described for Example 7. For an acylation using an isocyanate, for example for a compound of formula I wherein X is imino and J is oxygen, using a catalyst, for example, cuprous chloride, as described in Example 113, may be preferred.
- (D) For a compound of formula I wherein R is a sulfonyl group, sulfonylation of a corresponding amine of formula I wherein R is hydrogen with a corresponding sulfonic acid of formula D.W.SO₂.OH, or an activated derivative thereof, such as an acid halide, particularly a sulfonyl (or sulfamoyl) chloride of formula D.W.SO₂.Cl. The sulfonylation is conveniently carried out in an inert solvent or diluent, such as dichloromethane, tetrahydrofuran or toluene, at about ambient temperature, using an organic base such as, for example, triethylamine or pyridine, or an inorganic base, such as sodium or potassium carbonate, as an acid acceptor. If a sulfonyl chloride is not commercially available, it may be obtained by a conventional method.
- (E) For a compound of formula I in which R is a group G, substitution of the group L of a corresponding compound of formula G-L, wherein L is a conventional leaving group, such as for example halogeno, methylsulfonyloxy, trifluoromethylsulfonyloxy or diazonium, with a corresponding amine of formula I wherein R is hydrogen, optionally using a conventional catalyst.
- (F) For a compound of formula I which bears a hydroxy substituent on an aryl or heteroaryl group, cleaving the alkyl ether or acyloxy ester of a corresponding compound of formula I which bears a (1-4C)-alkoxy or containing 1 to 5 carbon atoms acyloxy substituent on an aryl or heteroaryl group. Convenient methods include, for example, the cleavage of a methoxy group using boron tribromide or pyridinium chloride and the cleavage of a t-butoxy group using trifluoroacetic acid for an alkyl ether, and the acidic or alkaline hydrolysis of an acyloxy group.
- (G) For a compound of formula I which bears a group of formula COORa in which Ra is hydrogen (a carboxy group), decomposing the ester group of a corresponding ester made with a conveniently removed acid protecting group, for example a corresponding compound of formula I in which Ra is not hydrogen. The decomposition may

be carried out using any one of the variety of procedures well known in organic chemistry, for example basic hydrolysis using lithium or sodium hydroxide, or by hydrogenolysis of a benzyl ester.

(H) For a compound of formula I bearing a moiety of formula COORa , CONRbRc , $\text{COO}(\text{CH}_2)_2\text{NRfRf}$ or $\text{CONRdSO}_2\text{R}^1$, acylation of a corresponding compound of formula HORA , HNRbRc , $\text{HO}(\text{CH}_2)_2\text{NRfRf}$ or $\text{HNRdSO}_2\text{R}^1$ with a corresponding acid of formula I bearing a moiety of formula COORa in which Ra is hydrogen, or an activated derivative thereof.

(I) For a compound of formula I bearing a containing 1 to 5 carbon atoms acyloxy group or a group of formula NRgCHO , NRgCOR^2 , NRgCOOR^2 , NRhCQNRiRj or NRkSO_2R^3 , acylation or sulfonylation of a corresponding compound of formula I bearing a hydroxy group or an amino group of formula NHRg , NHRh or NHRk (i.e. an amino group of formula NRfRf in which Rf is hydrogen and Rf is Rg, Rh or Rk) with an activated derivative of a corresponding acid of formula HOCHO , HOCOR^2 , HOCOOR^2 , HOCQNRiRj (including an isocyanate or isothiocyanate) or HOSO_2R^3 , respectively, using a conventional method.

(J) For a compound of formula I which bears a heteroaryl N-oxide group, oxidation of a corresponding compound of formula I which bears a heteroaryl group using a conventional oxidant, such as for example with peracetic acid or with dioxirane in acetone.

(K) For a compound of formula I which bears a primary amino group, reduction of a corresponding compound bearing a nitro group using a conventional reducing method, such as for example, hydrogenation over a palladium catalyst, or reduction with tin(II) chloride or with iron in acetic acid.

[0024] Whereafter, for any of the above procedures, when a pharmaceutically acceptable salt of an acidic or basic compound of formula I is required, it may be obtained by reacting the acidic or basic form of such a compound of formula I with a base or acid affording a physiologically acceptable counterion or by any other conventional procedure.

[0025] If not commercially available, the necessary starting materials for the above procedures may be made by procedures which are selected from standard techniques of heterocyclic chemistry and peptide chemistry, techniques which are analogous to the synthesis of known, structurally similar compounds, and techniques which are analogous to the above described procedures or the procedures described in the Examples. For uniformity and clarity, compounds herein are represented as the 6-pyrimidone, rather than the 6-hydroxypyrimidine, tautomers.

[0026] As will be clear to one skilled in the art, a variety of sequences is available for preparation of the starting materials. According to one of the available routes, a key intermediate pyrimidin-6-one-1-acetic acid of formula III may be prepared as shown in Scheme I (set out, together with other Schemes, following Examples) and as described in the Examples. In the Schemes, CBZ represents a benzyloxycarbonyl group.

[0027] In general, a nitrile of formula R^6CN is converted into a corresponding imidic ester of formula IV wherein R^7 is methyl or ethyl, conveniently isolated as its hydrochloride, if the imidic ester is not commercially available. Reaction of the imidic ester with an amine of formula $\text{H}_2\text{NCH}_2\text{R}^8$ in which R^8 is a latent or protected carboxaldehyde group, such as vinyl, dimethoxymethyl or diethoxyethyl, affords a corresponding amidine of formula V, conveniently isolated as its hydrochloride salt. Cyclization of an amidine of formula V with diethyl ethoxyethylenemalonate affords a corresponding ethyl 1,2-disubstituted-6-pyrimidone-5-carboxylate of formula VI which is hydrolyzed to the 1,2-disubstituted-6-pyrimidone-5-carboxylic acid of formula VII.

[0028] An acid of formula VII may be converted into a corresponding isocyanate of formula VIII by a conventional method, for example by using diphenylphosphoryl azide in an inert solvent, as described in the examples. Conveniently, the isocyanate is not isolated, but is converted into a benzyl urethane of formula IX as also is shown in Scheme I. It will be clear to one skilled in the art that, in general, treatment of an isocyanate of formula VIII with a selected alcohol or amine of formula A.X.H in which X is oxy or imino will provide a corresponding product of formula IXa in which X is oxy or imino, and that the product of formula IXa may be carried forward to a corresponding product of formula I using one of the routes outlined below.

[0029] Transformation of R^8 into a carboxaldehyde to afford a corresponding compound of formula X from a compound of formula IX is the next step. If R^8 is a vinyl group, the transformation may be carried out using N-methylmorpholine-N-oxide and osmium tetroxide, as described in Example 1, part e. If R^8 is a dimethoxymethyl or diethoxyethyl group, the acetal may be hydrolyzed with dilute hydrochloric acid, as described in Example 8, part f. Oxidation of an acetaldehyde derivative of formula X to provide a corresponding substituted acetic acid of formula III is conveniently carried out as described in Example 1f using sodium chlorite as the oxidant.

[0030] Elaboration of an acetic acid derivative of formula III into a corresponding intermediate alcohol of formula II or amine of formula Vb may be carried out as outlined in Scheme II. Thus, coupling a substituted acetic acid of formula III with an amino alcohol of formula XI, using, for example the method described in Example 1, part g, affords a corresponding alcohol of formula II in which R is benzyloxycarbonyl. Oxidation using a procedure as described above in process (A) provides a corresponding ketone of formula I in which R is benzyloxycarbonyl. As described above in process (B), by removing the benzyloxycarbonyl group, for example by hydrogenolysis as described in Example 6, an aminoketone of formula I in which R is hydrogen is obtained. Alternatively, an alcohol of formula II in which R is

benzyloxycarbonyl may be converted into a corresponding compound of formula XII in which R_p represents an alcohol protecting group, conveniently *tert*-butyldimethylsilyl, for example as described in Example 2, part a. Removal of the benzyloxycarbonyl group of a compound of formula XII by a conventional method, for example as noted in process (B) above, affords a corresponding 5-amino pyrimidone derivative of formula XIII.

- 5 [0031] A 3-amino pyridone of formula XIII may then be acylated, sulfonylated or b substituted with a group G by using a conventional method to afford a corresponding pyridone of formula XIV. Conventional acylation and sulfonylation methods and methods for introducing a group R include those described above in processes (C), (D) and (E) for substituting an amine of formula I wherein R is hydrogen. (Should a portion of bis-sulfonylated product be obtained, treatment with aqueous base at an elevated temperature may be used to remove the more labile second sulfonyl group at a convenient stage in the synthesis.) Removal of a *tert*-butyldimethylsilyl group to provide a corresponding alcohol of formula II may be carried out using tetrabutylammonium fluoride in an inert solvent, for example as described in Example 2, part d; it may be preferred to use acetic acid to buffer the reaction conditions.

- 10 [0032] Alternatively, for preparation of an intermediate of formula Vb, oxidation of an alcohol of formula II wherein R is benzyloxycarbonyl, using a method similar to one described in process (A), affords a corresponding ketone of formula I wherein R is benzyloxycarbonyl. Removal of the nitrogen protecting group of a ketone of formula I wherein R is benzyloxycarbonyl by hydrogenolysis or by treatment with a strong acid affords a corresponding amine of formula I wherein R is hydrogen. A preferred method for introducing the substituent R when it is a group G, particularly when it is an alkyl or substituted alkyl group, is by the use of a corresponding compound in which the pyrimidone 5-amino substituent bears an activating/protecting group of formula R_x, for example, benzyloxycarbonyl or trifluoroacetyl. Thus, acylation of a compound of formula I wherein R is hydrogen with trifluoroacetic anhydride affords a corresponding compound of formula Va in which R_x is trifluoroacetyl, which compound also may be prepared by an alternative order of steps via the corresponding compound of formula XIII. It will be noted that each of a compound of formula Va in which R_x is benzyloxycarbonyl or trifluoroacetyl is also a compound of formula I in which R is an acyl group. Alkylation, using a corresponding reagent of formula G.L in which G is alkyl or substituted alkyl, then provides a corresponding intermediate of formula Vb.

- 15 [0033] The trifluoromethyl amino alcohols of Formula XI required for the synthesis routes described above may be prepared by known routes. For example, 3-amino-1,1,1-trifluoro-4-methyl-2-pentanol (as its hydrochloride salt) conveniently may be obtained as described in U.S. Patent 4,910,190 in Example 4 (as a single diastereomer) or Example 6 (as a single enantiomer of a single diastereomer). If it is desired to carry out a chiral synthesis of a compound of formula I, using the single enantiomer in a substantially enantiomerically pure form and using methods and conditions which avoid epimerization at the center indicated by *** in formula I provide such a synthesis.

- 20 [0034] It may be desired optionally to use a protecting group during all or portions of the above described processes; the protecting group then may be removed when the final compound or a required starting material is to be formed. As will be clear to one skilled in the art, the order of steps in the sequences leading to the starting materials and products of the invention may be altered if appropriate considerations relative to coupling methods, racemization, deprotection methods, etc. are followed.

- 25 [0035] The utility of a compound of the invention or a pharmaceutically acceptable salt thereof (hereinafter, collectively referred to as a "Compound") may be demonstrated by standard tests and clinical studies, including those described below.

40

Inhibition Measurements:

- 45 [0036] The potency of a Compound to act as an inhibitor of human leukocyte elastase (HLE) on the low molecular weight peptide substrate methoxy-succinyl-alanyl-alanyl-prolyl-valine-p-nitroanilide is determined as described in U.S. Patent 4,910,190. The potency of an inhibitor is evaluated by obtaining a kinetic determination of the dissociation constant, K_i, of the complex formed from the interaction of the inhibitor with HLE. If a Compound is found to be a "slow-binding" inhibitor of HLE, special methods of analysis to accurately determine K_i values for the inhibition of HLE are carried out as described in U.S. Patent 4,910,190. In general, the K_i values for Compounds of the invention which were tested are generally on the order of 10⁻⁷ M or much less. For example, a K_i of 27 nM was determined for the Compound of the invention described as Example 51.

50

Acute Lung Injury Model:

- 55 [0037] Animal models of emphysema include intratracheal (i.t.) administration of an elastolytic protease to cause a slowly progressive, destructive lesion of the lung. These lesions are normally evaluated a few weeks to a few months after the initial insult. However, these proteases also induce a lesion that is evident in the first few hours. The early lesion is first hemorrhagic, progresses to an inflammatory lesion by the end of the first 24 hours and resolves in the first week post insult. To take advantage of this early lesion, the following model was used.

[0038] Hamsters are first lightly anesthetized with Brevital. Phosphate buffered saline (PBS) pH 7.4, either alone or containing human leukocyte elastase (HLE), is then administered directly into the trachea. Twenty-four hours later the animals are killed and the lungs removed and carefully trimmed of extraneous tissue. Following determination of wet lung weight, the lungs are lavaged with PBS and total lavagable red and white cells recovered are determined. The values for wet lung weights, total lavagable red cells and total lavagable white cells are elevated in a dose-dependent manner following administration of HLE. Compounds that are effective elastase inhibitors can prevent or diminish the severity of the enzyme-induced lesion resulting in lower wet lung weight and reduced values for total lavagable cells, both red and white, relative to administration of HLE alone. Compounds can be evaluated by administering them intratracheally as solutions or suspensions in PBS, either with or at various times prior to the HLE challenge (400 µg), or by dosing them intravenously or orally as solutions at various times prior to the HLE challenge (100 µg) to determine their utility in preventing an HLE lesion. A solution of a Compound is conveniently prepared using 10% polyethylene glycol 400/PBS or 10% polyethylene glycol 400/water. For a Compound which is acidic or basic, base (e.g. sodium hydroxide solution) or acid (e.g. hydrochloric acid) may be added as indicated to achieve solution. Compounds of this invention produced statistically significant reductions in wet lung weight and total lavagable cells relative to HLE alone.

Acute Hemorrhagic Assay:

[0039] This assay relies on monitoring only the amount of hemorrhage in the lung following intratracheal administration of human neutrophil elastase (HNE). Hemorrhage is quantified by disrupting erythrocytes recovered in lung lavage fluid and comparing that to dilutions of whole hamster blood. The screening protocol, similar to that described in Fletcher et al., *American Review of Respiratory Disease* (1990), 141, 672-677, is as follows. Compounds demonstrated to be HNE inhibitors *in vitro* are conveniently prepared for dosing as described above for the Acute Lung Injury Model. The compounds are then dosed by mouth to male Syrian hamsters at a fixed time, such as 30 or 90 min, prior to intratracheal administration of 50 µg/animal of HNE in 300 µL phosphate buffered saline (PBS) pH 7.4. Four hours enzyme administration, the animals are killed with an overdose of pentobarbital sodium, the thorax opened and the lungs and trachea removed. The excised lungs are lavaged with three changes of 2 mL normal saline via a tracheal cannula. The recovered lavages are pooled, the volumes (about 5 mL) are recorded and the lavages stored at 4 °C until assayed. For calculation of the amount of blood in each sample, the thawed lavages and a sample of whole hamster blood are sonicated to disrupt erythrocytes and appropriately diluted into individual wells of a 96-well microtiter plate. The optical densities (OD) of the disrupted lavages and blood samples are determined at 405 nm. The (µL blood equivalents) / (mL lavage) are determined by comparing the OD of the test samples with the OD of the standard curve prepared from whole hamster blood. The total µL equivalents of blood recovered is determined by multiplying recovered lavage volume by the (µL blood equivalents) / (mL lavage) for each sample. Results are reported as % inhibition of hemorrhage with respect to PBS treated controls when the test compound is given at a specified dose and time prior to administration of HNE. For example, the Compound of the invention described as Example 51 provided statistically significant inhibition of hemorrhage when administered at a dose of 2.5 mg/kg 30 or 90 min prior to administration of HNE.

[0040] No overt toxicity was observed when Compounds of the invention were administered in the above *in vivo* tests.

[0041] It will be appreciated that the implications of a Compound's activity in the Acute Lung Injury Model or Acute Hemorrhagic Assay are not limited to emphysema, but, rather, that the test provides evidence of general *in vivo* inhibition of HLE.

[0042] Compounds of the present invention which were tested exhibited activity in at least one of the tests described above under Inhibition Measurement, Acute Lung Injury Model and Acute Hemorrhagic Assay. It should be noted that there was not always a direct correlation between the activities of the compounds measured as K_i values in the Inhibition Measurement test and the reduced values for total lavagable cells and wet lung weights relative to the administration of HLE alone obtained in the Acute Lung Injury Model test or inhibition of hemorrhage in the Acute Hemorrhagic Assay.

[0043] According to a further feature of the invention, there is provided a pharmaceutical composition comprising a pharmaceutically effective amount of a Compound and a pharmaceutically acceptable diluent or carrier. As noted above, another feature of the invention is a method of using a Compound of the invention in the treatment of a disease or condition in a mammal, especially a human, in which HLE is implicated.

[0044] A Compound of the present invention may be administered to a warm-blooded animal, particularly a human, in need thereof for treatment of a disease in which HLE is implicated, in the form of a conventional pharmaceutical composition, for example as generally disclosed in U.S. Patent 4,910,190. The preferred mode of administration may be via a powdered or liquid aerosol. In a powdered aerosol, a Compound of the invention may be administered in the same manner as cromolyn sodium via a 'Spinhaler' (a trademark) turbo-inhaler device obtained from Fisons Corp. of Bedford, Massachusetts at a rate of about 0.1 to 50 mg per capsule, 1 to 8 capsules being administered daily for an average human. Each capsule to be used in the turbo-inhaler contains the required amount of a Compound of the invention with

the remainder of the 20 mg capsule being a pharmaceutically acceptable carrier such as lactose. In alliquid aerosol, a Compound of the invention may be administered using a nebulizer such as, for example, a 'Retec' (trademark) nebulizer, in which the solution is nebulized with compressed air. The aerosol may be administered, for example, at the rate of one to about eight times per day as follows: A nebulizer is filled with a solution of a Compound, for example 3.5 mL of solution containing 10 mg/mL; the solution in the nebulizer is nebulized with compressed air; and the patient breathes normally (tidal volume) for eight minutes with the nebulizer in his mouth.

[0045] Alternatively, the mode of administration may be oral or parenteral, including subcutaneous deposit by means of an osmotic pump. A compound of the invention may be conventionally formulated in an oral or parenteral dosage form by compounding about 10 to 250 mg per unit of dosage with conventional vehicle, excipient, binder, preservative, stabilizer, flavor or the like as called for by accepted pharmaceutical practice, e.g. as described in U.S. Patent 3,755,340. For parenteral administration, a 1 to 10 mL intravenous, intramuscular or subcutaneous injection would be given containing about 0.02 mg to 10 mg/kg of body weight of a compound of the invention 3 or 4 times daily. The injection would contain a compound of the invention in an aqueous isotonic sterile solution or suspension optionally with a preservative such as phenol or a solubilizing agent such as ethylenediaminetetraacetic acid (EDTA).

[0046] For parenteral administration or use in an aerosol, a 10 mg/mL aqueous formulation of an acidic Compound may be prepared, for example by dissolving the Compound (10 mg), dibasic sodium phosphate heptahydrate, USP (11.97 mg), monobasic sodium phosphate, USP (0.74 mg), sodium chloride, USP (4.50 mg) and sufficient 1 N sodium hydroxide solution or 0.05 M monobasic sodium phosphate solution to achieve pH 7.0-7.5 in sufficient water for injection, USP to afford 1.0 mL (1.01 g), followed by aseptic filtration, and sterile storage using standard procedures.

[0047] In general, a Compound of the invention will be administered to humans at a daily dose in the range of, for example, 5 to 100 mg of the Compound by aerosol or 50 to 1000 mg intravenously, or a combination of the two. However, it readily will be understood that it may be necessary to vary the dose of the Compound administered in accordance with well known medical practice to take account of the nature and severity of the disease under treatment, concurrent therapy, and the age, weight and sex of the patient receiving treatment. It similarly will be understood that generally equivalent amounts of a pharmaceutically acceptable salt of the Compound also may be used. Protocols for the administration of the HLE inhibitor and evaluation of the patients are described in the European Patent Applications with Publication Numbers 458535, 458536, 458537, and 463811 for the treatment or prevention of cystic fibrosis, ARDS, bronchitis, and hemorrhage associated with acute non-lymphocytic leukemia or its therapy, respectively; and a Compound of the invention may be used similarly for the treatment of those diseases and conditions either alone or in combination with another therapeutic agent customarily indicated for the treatment of the particular condition. For therapeutic or prophylactic treatment of a vascular disease or related condition in a mammal in which neutrophils are involved or implicated, a Compound of the invention may conveniently be administered by a parenteral route, either alone or simultaneously or sequentially with other therapeutically active agents customarily administered for the condition.

[0048] The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

(i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25 °C;

(ii) organic solutions were dried over anhydrous sodium sulfate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 pascals; 4.5-30 mm Hg) with a bath temperature of up to 60 °C;

(iii) chromatography means 'flash chromatography' (method of Still) carried out on Merck Kieselgel (Art 9385 from E. Merck, Darmstadt, Germany); if 'acidic silica gel' is indicated, material custom prepared by J. T. Baker Chemical Co., Phillipsburg, NJ, USA, and having a pH of about 6 when slurried in water was used; reversed phase chromatography means flash chromatography over octadecylsilane (ODS) coated support having a particle diameter of 32-74 µ, known as 'PREP-40-ODS' (Art 731740-100 from Bodman Chemicals, Aston, PA, USA); thin layer chromatography (TLC) was carried out on 0.25 mm silica gel GHLF plates (Art 21521 from Analtech, Newark, DE, USA); reversed phase-TLC (RP-TLC) was carried out Whatman MKC₁₈F plates (Art 4803-110 from Bodman Chemicals);

(iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;

(v) melting points are uncorrected and (dec) indicates decomposition; the melting points given are those obtained for the materials prepared as described; polymorphism may result in isolation of materials with different melting points in some preparations;

(vi) final products had satisfactory nuclear magnetic resonance (NMR) spectra;

(vii) yields are given for illustration only and are not necessarily those which may be obtained by diligent process development; preparations were repeated if more material was required;

(viii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 250 MHz using DMSO-d₆ as solvent; conventional abbreviations for signal shape are used; for AB spectra the directly observed shifts are reported;

(bx) chemical symbols have their usual meanings; SI units and symbols are used;

(x) reduced pressures are given as absolute pressures in pascals (Pa); elevated pressures are given as gauge pressures in bars;

(xd) solvent ratios are given in volume:volume (v/v) terms; and

(xii) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical ionization mode using a direct exposure probe; where indicated ionization was effected by electron impact (EI) or fast atom bombardment (FAB); generally, only peaks which indicate the parent mass are reported.

EXAMPLE 1

2-(5-Benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0049] To a solution of 2-(5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide (0.537 g), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.9 g) in toluene (4 mL) and dimethyl sulfoxide (4 mL) was added dichloroacetic acid (0.33 mL) and the resulting solution was allowed to stir for 0.5 h. The reaction mixture was diluted with ethyl acetate, washed (saturated ammonium chloride, water), dried and evaporated to provide a white solid which was collected and washed with ether:hexane (1:1) to give 2-(5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (0.330 g). Chromatography of the mother liquor with ethyl acetate:dichloromethane (5:95, 10:90) as the eluent gave additional 2-(5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid (0.190 g); NMR (DMSO- D_2O): 8.46 (s,1), 7.44 (m,10), 5.20 (s,2), 4.54 (m,2), 4.05 (d,1), 2.23 (m,1), 0.85 (d,3), 0.75 (d,3); MS: $m/z=531(M+1)$.

Analysis for $C_{26}H_{25}F_3N_4O_5$:			
Calculated:	C, 58.8;	H, 4.74;	N, 10.5
Found:	C, 58.4;	H, 4.75;	N, 10.6

[0050] The intermediate 2-(5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. N-Allyl benzamidinium hydrochloride.

[0051] To a solution of ethyl benzimidate hydrochloride (20 g) in methanol at 0 °C was added allyl amine. The resulting solution was allowed to stand for 2 days at 5 °C before it was evaporated to yield a solid which was collected and washed with ether to give N-allyl benzamidinium hydrochloride (21.5 g) as a white solid; 300 MHz NMR: 10.1 (s,1), 9.68 (s,1), 9.29 (s,1), 7.72 (s,5), 5.92 (m,1), 5.35 (d,2), 5.26 (d,2), 4.14 (s,2).

b. Ethyl 1-allyl-2-phenylpyrimidin-6(1H)-one-5-carboxylate.

[0052] The free base of N-allyl benzamidinium hydrochloride was generated by dissolving N-allyl benzamidinium hydrochloride (79.7 g) in 1 N sodium hydroxide. The free base was then extracted into dichloromethane, which was dried and evaporated to provide N-allyl benzamidinium (65.2 g). This was added to diethyl ethoxymethylene malonate (78 mL) in ethanol (50 mL). The resulting solution was heated at 120 °C for 2 h. The solution was cooled, diluted with ethyl acetate, washed (saturated ammonium chloride, water), dried, and evaporated to give a solid, which was collected and washed two times with ether:hexane (1:1), to provide ethyl 1-allyl-2-phenylpyrimidin-6(1H)-one-5-carboxylate as a white solid (62.5 g); 300 MHz NMR: 8.56 (s,1), 7.54 (m,5), 5.80 (m,1), 5.09 (d,1), 4.82 (d,1), 4.47 (d,2), 4.28 (q,2), 1.28 (t,3).

c. 1-Allyl-2-phenylpyrimidin-6(1H)-one-5-carboxylic acid.

[0053] To a solution of ethyl 1-allyl-2-phenylpyrimidin-6(1H)-one-5-carboxylate (25.6 g) in tetrahydrofuran (300 mL) at 0 °C was added a solution of 0.5 N sodium hydroxide (198 mL). The resulting solution was allowed to stir for 1 h, was poured into dichloromethane and the organic layer removed. The remaining basic aqueous fraction was extracted with dichloromethane, made acidic with 1 N hydrochloric acid (to pH 2), and extracted with dichloromethane. The organic

layers from the acidic extractions were dried and evaporated to give an oil which crystallized upon addition of ether. The resulting white solid was collected and washed with ether:hexane (1:1) to give 1-allyl-2-phenylpyrimidin-6(1H)-one-5-carboxylic acid (11.1 g); 300 MHz NMR: 13.0 (s broad, 1), 8.69 (s, 1), 7.58 (m, 5), 5.82 (m, 1), 5.16 (d, 1), 4.87 (d, 1), 4.51 (d, 2).

5

d. 1-Allyl-5-benzoyloxycarbonylamino-2-phenylpyrimidin-6(1H)-one.

[0054] To a solution of 1-allyl-2-phenylpyrimidin-6(1H)-one-5-carboxylic acid (30.2 g) and triethylamine (32.8 mL) in dioxane (390 mL) was added diphenylphosphoryl azide (25.6 mL), and the resulting solution was heated at 100 °C for 2 h. Benzyl alcohol (24.5 mL) was added and the resulting solution was heated at 100 °C for 12 h. The solution was cooled and the solvent evaporated. The resulting residue was dissolved in ethyl acetate, washed (saturated ammonium chloride, 1 N sodium hydroxide, water), dried, and evaporated to give an oil which crystallized upon addition of ether to give a white solid. The solid was collected and washed with ether to provide 1-allyl-5-benzoyloxycarbonylamino-2-phenylpyrimidin-6(1H)-one (25.1 g); 300 MHz NMR: 8.93 (s, 1), 8.45 (s, 1), 7.43 (m, 10), 5.75 (m, 1), 5.18 (s, 2), 5.08 (d, 1), 4.82 (d, 1), 4.46 (d, 2).

15

e. 5-Benzoyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinylacetaldehyde.

[0055] To a solution of 1-allyl-5-benzoyloxycarbonylamino-2-phenylpyrimidin-6(1H)-one in tetrahydrofuran (200 mL) and water (30 mL) was added N-methylmorpholine-N-oxide (9.82 g) and osmium tetroxide (4.4 mL, 4% in water). The resulting solution was allowed to stir overnight. N-Methylmorpholine-N-oxide (1.65 g) was added and the solution was allowed to stir for 4 h. Sodium thiosulfate (saturated aqueous solution, 10 mL) and diatomaceous earth (30 g) were added and the mixture was stirred for 0.5 h. The mixture was filtered and evaporated to give an oil. This oil was dissolved in ethyl acetate, washed (saturated aqueous sodium thiosulfate solution, 1 N hydrochloric acid, brine), and evaporated to give an oil. This oil was dissolved in ethanol (230 mL) and a solution of sodium periodate (27 g) in water (40 mL) was added. The mixture was stirred for 2 h, filtered through diatomaceous earth and evaporated. The residue was dissolved in ethyl acetate and the solution was washed with water, dried, and evaporated to provide 5-benzoyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinylacetaldehyde (25 g) as a white solid; TLC: ethyl acetate:diethyl ether (1:1), $R_f=0.8$.

30

f. 5-Benzoyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinylacetic acid.

[0056] To a solution of 5-benzoyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinylacetaldehyde (25 g) in tert-butyl alcohol (175 mL), and 2-methyl-2-butene (148 mL) at 0 °C was added a solution of sodium chlorite (57 g) and sodium dihydrogen phosphate monohydrate (67 g) in water (190 mL). The mixture was allowed to stir for 3 h and was evaporated. The resulting material was diluted with ethyl acetate and extracted with 1 N aqueous sodium hydroxide. The aqueous solution was acidified to pH 3 with hydrochloric acid and was extracted with dichloromethane. The organic extracts were dried and evaporated to give a white solid, which was washed with ether:hexane (1:1) to yield the acid (17.2 g); 300 MHz NMR: 13.3 (s, 1), 9.04 (s, 1), 8.48 (s, 1), 7.43 (m, 10), 5.19 (s, 2), 4.51 (s, 2).

40

g. 2-(5-Benzoyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide.

[0057] To a solution of 5-benzoyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinylacetic acid (12.9 g), 3-amino-1,1,1-trifluoro-4-methyl-2-pentanol hydrochloride (10.6 g), 1-hydroxybenzotriazole hydrate (9.2 g), and triethylamine (9.5 mL) in N,N-dimethylformamide (115 mL) at 25 °C was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (7.8 g). The resulting solution was allowed to stir for two days, was diluted with ethyl acetate, washed (saturated ammonium chloride, 1 N sodium hydroxide, water), dried, and evaporated to give a white solid, which was collected and washed with ether:hexane (1:1) to provide pure 2-(5-benzoyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide (15.5 g); NMR: 8.95 (s, 1), 8.45 (s, 1), 8.00 (d, 1), 7.45 (m, 9), 8.51 (d, 1), 5.18 (s, 2), 4.46 (m, 2), 4.10 (m, 1), 3.82 (t, 1), 1.72 (m, 1), 0.86 (d, 3), 0.73 (d, 3); MS: $m/z=533(M+1)$.

50

EXAMPLE 2

55 2-(5-Acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0058] To a solution of 2-(5-acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)-acetamide (0.6 g) in toluene (4 mL) and dimethyl sulfoxide (4 mL) was added 1-(3-dimethylaminopro-

pyl)-3-ethylcarbodiimide (2.4 g) and dichloroacetic acid (0.41 mL). The resulting solution was allowed to stir for 1 h, was poured into 1 N hydrochloric acid, and was extracted with ethyl acetate. The organic solution was washed (1 N hydrochloric acid, water), dried, and evaporated. The resulting oil crystallized upon addition of ether to give a white solid, which was collected and washed with ether:hexane (1:1) to provide 2-(5-acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (0.35 g); NMR (DMSO/D₂O): 8.77 (s,1), 7.50 (m,5), 4.80 (d,1), 4.53 (d,1), 4.03 (m,1), 2.24 (m,1), 2.18 (s,3), 0.84 (d,3), 0.77 (d,3).

Analysis for C ₂₀ H ₂₁ F ₃ N ₄ O ₄ :			
Calculated:	C, 54.8;	H, 4.82;	N, 12.8
Found:	C, 53.9;	H, 4.77;	N, 12.6

[0059] The intermediate 2-(5-acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. 2-(5-Benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

[0060] To a suspension of 2-(5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide (1.13 g) and 2,6-lutidine (0.5 mL) in dichloromethane (5 mL) at 0 °C was added tert-butyldimethylsilyl trifluoromethanesulfonate (0.73 mL). The mixture was allowed to stir for 1.5 h. over which time the suspension became a solution. The mixture was poured into 1 N hydrochloric acid and extracted into ether. The organic layer was washed (water), dried and evaporated to give an oil, which crystallized upon addition of ether:hexane to give a white solid. The solid was collected and washed with hexane to yield 2-(5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a white solid (1.1 g); MS: m/z=647(M+1).

b. 2-(5-Amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

[0061] To a solution of 2-(5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide (12.4 g) in tetrahydrofuran (100 mL) and ethanol (60 mL) was added 10% (w/w) palladium on carbon (1 g) and the suspension was placed under a hydrogen atmosphere (3.4 bar) and shaken for 12 h. The solution was filtered through diatomaceous earth to give an oil which crystallized from ether to provide 2-(5-amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a white solid (8.7 g); MS: m/z=513(M+1).

c. 2-(5-Acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

[0062] To a solution of 2-(5-amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide (0.83 g) and triethylamine (0.45 mL) in tetrahydrofuran (10 mL) at 0 °C was added acetyl chloride (0.17 mL) and the resulting solution was allowed to stir for 0.5 h. The mixture was poured into ethyl acetate and the solution was washed (1 N hydrochloric acid, water), dried, and evaporated to give 2-(5-acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a foamy solid (0.86 g); TLC: R_f=0.4, methanol:dichloromethane (5:95).

d. 2-(5-Acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide.

[0063] To a solution of 2-(5-acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide (0.86 g) in tetrahydrofuran (15 mL) at 0 °C was added tetrabutylammonium fluoride (1.95 mL, 1M in tetrahydrofuran) and the resulting solution was allowed to stir for 5 min. The mixture was poured into saturated aqueous ammonium chloride and the product extracted into ethyl acetate. The organic solution was washed (water), dried, and evaporated. The resulting white solid was collected and washed with ether:hexane to provide 2-(5-acetamido-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-hydroxypropyl)aceta-

mide (0.6 g); NMR: 8.87 (s,1), 8.07 (d,1), 7.57 (m,5), 4.62 (d,1), 4.42 (d,1), 4.10 (m,1), 3.68 (t,1), 2.18 (s,3), 1.80 (m,1), 0.89 (d,3), 0.82 (d,3).

EXAMPLE 3

2-[6-Oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxo-propyl)acetamide.

[0064] The title compound was prepared from 2-[6-oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide using a similar method to that described in Example 1 to obtain 2-[6-oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid (0.6 g); NMR (DMSO/D₂O): 8.89 (d,2), 8.49 (s,1), 8.12 (d,2), 7.49 (m,7), 5.49 (s,2), 4.67 (d,1), 4.49 (d,1), 4.04 (d,1), 2.23 (m,1), 0.82 (d,3), 0.74 (d,3).

Analysis for C ₂₅ H ₂₄ F ₃ N ₅ O ₅ :			
Calculated:	C, 56.5;	H, 4.55;	N, 13.2
Found:	C, 55.6;	H, 4.68;	N, 13.2

[0065] The intermediate 2-[6-oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. 2-[6-Oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

[0066] To a solution of 2-(5-amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide (1.1 g) and triphosgene (0.93 g) in dichloromethane at 0 °C was added triethylamine (2.0 mL) and the resulting solution was allowed to stir for 0.5 h. To this solution was added 4-pyridylcarbinol and the resulting solution allowed to stir overnight. The solution was poured into saturated aqueous sodium bicarbonate solution and the product was extracted into dichloromethane. The organic solution was washed (saturated aqueous sodium bicarbonate, water), dried, and evaporated. The resulting oil was purified by chromatography, with methanol:dichloromethane (5:95) as the eluent, to provide 2-[6-oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a white solid (0.9 g); TLC: R_f=0.4, methanol:dichloromethane (5:95).

b. 2-[6-Oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide.

[0067] This compound was prepared from 2-[6-oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide by a method similar to that used in Example 2.d. to obtain pure 2-[6-oxo-2-phenyl-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide as a white solid; 300 MHz NMR: 9.23 (s,1), 8.58 (d,2), 8.46 (s,1), 8.02 (d,1), 7.45 (m,7), 6.54 (d,1), 5.23 (s,2), 4.62 (d,1), 4.43 (d,1), 4.08 (m,1), 3.80 (t,1), 4.17 (m,1), 0.88 (d,3), 0.79 (d,3).

EXAMPLE 4

2-[6-Oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxo-propyl)acetamide.

[0068] 2-[6-Oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was subjected to conditions similar to those described in Example 1. Chromatography, with methanol:dichloromethane (5:95) as the eluent, gave 2-[6-oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; NMR (DMSO/D₂O): 8.60 (s,1), 8.42 (m,3), 7.71 (m,1), 7.47 (m,5), 4.58 (d,1), 4.39 (d,1), 4.31 (s,2), 2.20 (m,1), 0.83 (d,3), 0.77 (d,3).

Analysis for C ₂₅ H ₂₅ F ₃ N ₆ O ₄ :			
Calculated:	C, 56.6;	H, 4.75;	N, 15.8
Found:	C, 56.1;	H, 4.89;	N, 15.5

10 [0069] The intermediate 2-[6-oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-hydroxypropyl)acetamide was prepared as follows:

a. 2-[6-Oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

15 [0070] 2-(5-Amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)-acetamide and 3-aminomethylpyridine were subjected to a method similar to that described in Example 3.a. Chromatography, with methanol:dichloromethane (5:95) as the eluent, gave 2-[6-oxo-2-phenyl-5-[3-(3-pyridyl)methylureido]-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a white solid; MS: m/z=647(M+1).

b. 2-[6-Oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)-acetamide.

25 [0071] 2-[6-Oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide was subjected to a method similar to that described in Example 2.d. Chromatography, with methanol:dichloromethane (5:95) as the eluent, gave 2-[6-oxo-2-phenyl-5-[3-(3-pyridyl)methylureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide as a white solid; MS: m/z=533(M+1).

30 EXAMPLE 5

2-[6-Oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

35 [0072] 2-[6-Oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 1. Chromatography, with methanol:dichloromethane (10:90) as the eluent, gave 2-[6-oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; NMR (DMSO/D₂O): 8.58 (s,1), 8.46 (bs, 2), 7.41 (m,5), 7.27 (d,2), 4.56 (d,1), 4.37 (d,1), 4.32 (s,2), 4.00 (d,1), 2.20 (m,1) 0.79 (d,3), 0.73 (d,3); MS: m/z=531(M+1).

Analysis for C ₂₅ H ₂₅ F ₃ N ₆ O ₄ :			
Calculated:	C, 56.6;	H, 4.75;	N, 15.8
Found:	C, 55.0;	H, 4.88;	N, 14.8

50 [0073] The intermediate 2-[6-oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. 2-[6-Oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

55 [0074] 2-(5-Amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide and 4-aminomethylpyridine were subjected to a procedure similar to that described in Example 3.a. Chromatog-

raphy, with methanol:dichloromethane (5:95) as the eluent, gave 2-[6-oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(2-*tert*-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a white solid; MS: $m/z=647(M+1)$.

- b. 2-[6-Oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)-acetamide.

[0075] 2-[6-Oxo-2-phenyl-5-[3-(4-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(2-*tert*-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 2.d. to provide 2-[6-oxo-2-phenyl-5-[3-(3-pyridylmethyl)ureido]-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide as a white solid; MS: $m/z=533(M+1)$.

EXAMPLE 6

- 2-[5-Amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0076] To a solution of 2-[5-benzyloxycarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (1.9 g) in tetrahydrofuran (50 mL) and ethanol (50 mL) was added 10% (w/w) palladium on carbon (0.29 g) and the resulting solution was placed under a hydrogen atmosphere (0.75 bar) for 12 h. The catalyst was removed by filtration through diatomaceous earth and the solvent was evaporated. The resulting oil was crystallized from ether. The product was collected and washed with ether:hexane (1:1) to provide 2-[5-amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (1.14 g) as a white solid; NMR (DMSO- D_6): 7.38 (m, 6), 4.55 (d, 1), 4.37 (d, 1), 3.98 (m, 1), 2.17 (m, 1), 0.79 (d, 3), 0.72 (d, 3).

Analysis for $C_{18}H_{19}F_3N_4O_3$:			
Calculated:	C, 54.5;	H, 4.83;	N, 14.1
Found:	C, 52.2;	H, 5.17;	N, 13.5

EXAMPLE 7

- 2-[6-Oxo-2-phenyl-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0077] 2-[5-Amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide and 2-pyridylcarbinol were subjected to a procedure similar to that described in Example 3.a. Chromatography, with methanol:dichloromethane (gradient, 3:97, 5:95) as the eluent, gave 2-[6-oxo-2-phenyl-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a tan solid; 300 MHz NMR (DMSO- D_6): 8.53 (d, 1), 8.43 (d, 1), 7.80 (t, 1), 7.4 (m, 7), 5.21 (s, 2), 4.57 (d, 1), 4.47 (d, 1), 4.02 (s, 1), 2.20 (m, 1), 0.81 (d, 3), 0.73 (d, 3).

Analysis for $C_{25}H_{24}F_3N_5O_5$:			
Calculated:	C, 56.5;	H, 4.55;	N, 13.1
Found:	C, 55.9;	H, 4.62;	N, 13.0

EXAMPLE 8

- 2-[5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0078] 2-[5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

propyl-2-hydroxypropyl)acetamide was oxidized by a procedure similar to that described in Example 1 to afford 2-[5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; NMR: 8.42 (s,1), 7.38 (9), 5.17 (s,2), 4.69 (d,1), 4.41 (d,1), 4.00 (s,1), 2.19 (m,1), 0.81 (d,3), 0.70 (d,3).

Analysis for C ₂₆ H ₂₄ F ₄ N ₄ O ₅ :			
Calculated:	C, 56.9;	H, 4.41;	N, 10.2
Found:	C, 56.0;	H, 4.43;	N, 10.0

[0079] The intermediate 2-[5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. Ethyl 4-fluorobenzimidate hydrochloride.

[0080] A solution of 4-fluorobenzonitrile (50 g) in tetrahydrofuran (300 mL) and ethanol (60.5 mL) at 0 °C was saturated with anhydrous hydrogen chloride gas and the resulting solution was allowed to stand overnight. The solvent was evaporated and the resulting solid was collected and washed with ether to provide ethyl 4-fluorobenzimidate hydrochloride as a white solid (20 g); NMR: 8.27 (m,2), 7.51 (m,2), 4.63 (q, 2), 1.47 (t,3).

b. N-(2,2-Diethoxyethyl)-4-fluorobenzamidine.

[0081] To a solution of ethyl 4-fluorobenzimidate hydrochloride (18.5 g) in ethanol (90 mL) at 0 °C was added aminoacetaldehyde diethyl acetal (14.5 mL) and the resulting solution was kept at 5 °C overnight. The solvent was evaporated, the resulting oil was dissolved in 1 N sodium hydroxide (200 mL), and the solution was extracted with dichloromethane. The organic extracts were dried and evaporated to yield N-(2,2-diethoxyethyl)-4-fluorobenzamidine as an oil (21 g); MS: m/z=255(M+1).

c. Ethyl 1-(2,2-diethoxyethyl)-2-(4-fluorophenyl)pyrimidin-6(1H)-one-5-carboxylate.

[0082] This compound was prepared from diethyl ethoxymethylenemalonate and N-(2,2-diethoxyethyl)-4-fluorobenzamidine by a procedure similar to that described in Example 1.b. to obtain ethyl 1-(2,2-diethoxyethyl)-2-(4-fluorophenyl)pyrimidin-6(1H)-one-5-carboxylate as an oil; MS: m/z=379(M+1).

d. 1-(2,2-Diethoxyethyl)-2-(4-fluorophenyl)pyrimidin-6(1H)-one-5-carboxylic acid.

[0083] This compound was prepared from ethyl 1-(2,2-diethoxyethyl)-2-(4-fluorophenyl)pyrimidin-6(1H)-one-5-carboxylate by a procedure similar to that described in Example 1.c. to obtain the title compound as a white solid; 300 MHz NMR: 8.66 (s,1), 7.69 (m,2), 7.40 (m,2), 4.69 (t,1), 4.05 (d,2), 3.39 (m,4), 0.99 (t,6).

e. 5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetaldehyde diethyl acetal.

[0084] This compound was prepared from 1-(2,2-diethoxyethyl)-2-(4-fluorophenyl)pyrimidin-6(1H)-one-5-carboxylic acid by a procedure similar to that used in Example 1.d. to obtain 5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetaldehyde diethyl acetal as a white solid; TLC: R_f=0.6, ether:hexane (75:25).

f. 5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetaldehyde.

[0085] A solution of 5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetaldehyde diethylacetal in tetrahydrofuran (7 mL) and 1 N hydrochloric acid (5 mL) was heated at 60 °C for 18 h. The solution was cooled and neutralized with saturated aqueous sodium bicarbonate solution (pH 6). The solution was extracted with ethyl acetate and the organic extracts were dried and evaporated to give 5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetaldehyde as a white solid; NMR: 9.51 (s,1), 9.03 (s,1), 8.47 (s,1), 7.43 (m,9), 5.19 (s,2), 4.76 (s,2).

g. 5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetic acid.

[0086] This compound was prepared from 5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetaldehyde by a procedure similar to that described in Example 1.f. to provide 5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetic acid as a white solid; 300 MHz NMR: 9.06 (s,1), 8.46 (s,1), 7.42 (m,9), 5.19 (s,2), 4.52 (s,2).

h. 2-[5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)-acetamide.

[0087] The title compound was prepared from 5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinylacetic acid and 3-amino-1,1,1-trifluoro-4-methyl-2-pentanol hydrochloride by a procedure similar to that described in Example 1.g. to provide 2-[5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)-acetamide as a white solid; NMR: 8.94 (s,1), 8.44 (s,1), 7.97 (d,1), 7.4 (m,9), 6.50 (d,1), 5.18 (s,2), 4.65 (d,1), 4.38 (d,1), 4.08 (m,1), 3.80 (t,1), 1.72 (m,1), 0.85 (d,3), 0.78 (d,3).

EXAMPLE 9

2-[2-(4-Fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)-acetamide.

[0088] 2-[2-(4-Fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 1. Chromatography, with methanol:dichloromethane (gradient, 5:95, 7:93) as the eluent, gave 2-[2-(4-fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; NMR (DMSO/D₂O): 8.54 (d,2), 8.44 (s,1), 7.53 (m,2), 7.44 (d,2), 7.22 (t,2), 5.22 (s,2), 4.70 (d,1), 4.49 (d,1), 4.01, (d,1), 2.20 (m,1), 0.85 (d,3), 0.72 (d,3).

Analysis for C ₂₅ H ₂₃ F ₄ N ₃ O ₅ :			
Calculated:	C, 54.6;	H, 4.22;	N, 12.7
Found:	C, 53.6;	H, 4.29;	N, 12.5

[0089] The intermediate 2-[2-(4-fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. 2-[5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

[0090] 2-[5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 2.a. to provide 2-[5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a white solid; MS: m/z=665(M+1).

b. 2-[5-Amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl) acetamide.

[0091] 2-[5-Benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 2.b. to provide 2-[5-amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a gray-white solid; MS: m/z=531(M+1).

c. 2-[2-(4-Fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

5 **[0092]** 2-[5-Amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide and 4-pyridylcarbinol were subjected to a procedure similar to that described in Example 3.a. Chromatography, with methanol:dichloromethane (5:95) as the eluent, gave 2-[2-(4-fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a yellow solid; MS: $m/z=666(M+1)$.

10 d. 2-[2-(4-Fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide.

15 **[0093]** 2-[2-(4-Fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 2.d. Chromatography with methanol:dichloromethane (gradient, 5:95, 7:93) as the eluent, gave 2-[2-(4-fluorophenyl)-6-oxo-5-(4-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide as a white solid; MS: $m/z=552(M+1)$.

EXAMPLE 10

20 2-[2-(4-Fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl) acetamide.

25 **[0094]** 2-[2-(4-Fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 1. Chromatography, with methanol:dichloromethane (gradient, 0:100, 5:95, 7:93) as the eluent, gave 2-[2-(4-fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; NMR (DMSO- H_2O): 8.54 (d,2), 8.43 (s,1), 7.82 (m,1), 7.53 (m,3), 7.30 (m,3), 5.22 (s,2), 4.62 (d,1), 4.42 (d,1), 4.00, (d,1), 2.19 (m,1), 0.81(d,3), 0.70 (d,3).

30

Analysis for $C_{25}H_{23}F_4N_3O_5$:				
Calculated:	C, 54.6;	H, 4.22;	N, 12.7	
Found:	C, 54.0;	H, 4.56;	N, 12.2	

35

40 **[0095]** The intermediate 2-[2-(4-fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. 2-[2-(4-Fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide.

45 **[0096]** 2-[5-Amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide and 2-pyridylcarbinol were subjected to a procedure similar to that described in Example 3.a. Chromatography, with methanol:dichloromethane (gradient, 2:98, 5:95) as the eluent, gave 2-[2-(4-fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide as a gum; MS: $m/z=666(M+1)$.

50

b. 2-[2-(4-Fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide.

55 **[0097]** 2-[2-(4-Fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(2-tert-butyldimethylsilyloxy-3,3,3-trifluoro-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in Example 2.d. to provide 2-[2-(4-fluorophenyl)-6-oxo-5-(2-pyridylmethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide as a white solid; MS: $m/z=552(M+1)$.

EXAMPLE 11

2-[5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

5

[0098] 2-[5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was oxidized using a procedure similar to that described in Example 1. Chromatography, with ethyl acetate:dichloromethane (gradient, 15:85, 30:70) as the eluent, gave 2-[5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; 300 MHz
10 NMR (DMSO/D₂O): 8.41 (s,1), 7.79 (d,1), 7.35 (m,6), 7.10 (m,1), 5.17 (s,2), 4.90 (d,1), 4.76 (d,1), 4.08 (d,1), 2.25 (m,1), 0.90 (d,3), 0.75 (d,3).

15

Analysis for C ₂₄ H ₂₃ F ₃ N ₄ O ₅ S:			
Calculated:	C, 53.7;	H, 4.32;	N, 10.4
Found:	C, 52.4;	H, 4.49;	N, 10.2

20

[0099] The intermediate 2-[5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. Ethyl 2-thiophenecarboximidate hydrochloride.

25

[0100] 2-Thiophenecarbonitrile was subjected to a procedure similar to that described in Example 8.a. to provide ethyl 2-thiophenecarboximidate hydrochloride as a white solid; MS: m/z=156(M+1).

b. N-(2,2-Dimethoxyethyl)-2-thiophenecarboximidine.

30

[0101] Ethyl 2-thiophenecarboximidate hydrochloride and aminoacetaldehyde dimethyl acetal were subjected to a procedure similar to that described in Example 8.b. to provide N-(2,2-dimethoxyethyl)-2-thiophenecarboximidine as a white solid; 300 MHz NMR (DMSO/D₂O): 8.07 (m,1), 7.91 (m,1), 7.33 (m,1), 4.67 (t,1), 3.84 (s,6), 3.60 (d,2).

35

c. Ethyl 1-(2,2-dimethoxyethyl)-2-(2-thienyl)pyrimidin-6(1H)-one-5-carboxylate.

[0102] N-(2,2-Dimethoxyethyl)-2-thiophenecarboximidate and diethyl ethoxymethylenemalonate were subjected to a procedure similar to that described in Example 1.b. to provide ethyl 1-(2,2-dimethoxyethyl)-2-(2-thienyl)pyrimidin-6(1H)-one-5-carboxylate as a yellow solid; MS: m/z=339(M+1).

40

d. 1-(2,2-Dimethoxyethyl)-2-(2-thienyl)pyrimidin-6(1H)-one-5-carboxylic acid.

45

[0103] To a solution of ethyl 1-(2,2-dimethoxyethyl)-2-(2-thienyl)pyrimidin-6(1H)-one-5-carboxylate in pyridine (33 mL) was added lithium iodide (3.32 g) and the resulting mixture was heated at 110 °C overnight. The pyridine was evaporated and residual traces of pyridine were removed by evaporation with toluene. The residue was dissolved in 1 N hydrochloric acid and the product was extracted into ethyl acetate. The solution was dried and evaporated to give an oil which crystallized from ether to provide 1-(2,2-dimethoxyethyl)-2-(2-thienyl)pyrimidin-6(1H)-one-5-carboxylic acid as a brown solid; MS: m/z=311(M+1).

50

e. 5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetaldehyde dimethyl acetal.

55

[0104] 1-(2,2-Dimethoxyethyl)-2-(2-thienyl)pyrimidin-6(1H)-one-5-carboxylic acid was subjected to a procedure similar to that described in Example 1.d. Chromatography, with ether:hexane (gradient, 25:75, 80:20) as the eluent, gave 5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetaldehyde dimethyl acetal as a yellow solid; MS: m/z=416(M+1).

f. 5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetaldehyde.

[0105] 5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetaldehyde dimethyl acetal was subjected to a procedure similar to that described in Example 8.f. to provide 5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetaldehyde as a yellow foam; MS: $m/z=370(M+1)$.

g. 5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetic acid.

[0106] 5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetaldehyde was subjected to a procedure similar to that described in Example 1.f. to provide 5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetic acid as a yellow solid; MS: $m/z=386(M+1)$.

h. 2-[5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropyl)acetamide.

[0107] 5-Benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinylacetic acid and 3-amino-1,1,1-trifluoro-4-methyl-2-pentanol hydrochloride were subjected to a procedure similar to that described in Example 1.g. to provide 2-[5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-hydroxypropyl)acetamide as a yellow solid; MS: $m/z=539(M+1)$.

EXAMPLE 12

2-[5-Amino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0108] To a solution of 2-[5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide in dichloromethane (35 mL) and anisole (1.2 mL) at 0 °C was added trifluoromethanesulfonic acid, and the resulting suspension was stirred for 15 minutes. The reaction was quenched by addition of sodium bicarbonate and the product extracted into ethyl acetate. The organic layer was washed (brine) and dried. The solvent was removed and the residue was purified by chromatography, with methanol:dichloromethane (gradient, 5:95, 7:93) as the eluent, to provide 2-[5-amino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a yellow solid; 300 MHz NMR (DMSO- D_6): 7.94 (d,1), 7.65 (d,1), 7.30 (s,1), 7.22 (d,1), 7.02 (m,1), 4.84 (d,1), 4.69 (d,1), 4.07 (d,1), 2.24 (m,1), 0.89(d,3), 0.76 (d,3).

Analysis for $C_{16}H_{17}F_3N_4O_3S$:			
Calculated:	C, 47.7;	H, 4.26;	N, 13.9
Found:	C, 56.8;	H, 4.58;	N, 13.3

EXAMPLE 13

2-[5-Benzyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0109] 2-[5-Benzyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was subjected to a procedure similar to that described in example 1 to obtain pure 2-[5-benzyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; NMR (DMSO- D_6): 8.84 (m,2), 8.44 (s,1), 7.88 (m,1); 7.39 (m,5), 5.18 (s,2), 4.63 (d,1), 4.45 (d,1), 3.97, (d,1), 2.15 (m,1), 0.77 (d,3), 0.65 (d,3).

Analysis for $C_{25}H_{24}F_3N_5O_5$:			
Calculated:	C, 56.5;	H, 4.55;	N, 13.2

(continued)

Analysis for C ₂₅ H ₂₄ F ₃ N ₅ O ₅ :			
Found:	C, 55.0;	H, 4.60;	N, 12.85

[0110] The intermediate 2-[5-benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. Ethyl 3-pyridinecarboximidate hydrochloride.

[0111] To a solution of ethanol (200 mL) in chloroform (100 mL) at 0 °C was added acetyl chloride (190 mL) and a solution of 3-cyanopyridine (25 g) in (300 mL) of chloroform. The solution was allowed to warm to 25°C and to stand for 1 day. A white solid separated from solution. The solvent was removed and the white solid collected and washed with ether to provide pure ethyl 3-pyridinecarboximidate hydrochloride as a white solid; MS: m/z=152(M+1).

b. N-(2,2-Dimethoxyethyl)-3-pyridinecarboxamidine.

[0112] Ethyl 3-pyridinecarboximidate hydrochloride and aminoacetaldehyde dimethyl acetal were subjected to a procedure similar to that described in Example 8.b. to provide pure N-(2,2-dimethoxyethyl)-3-pyridinecarboxamidine as an oil; MS: m/z=210(M+1).

c. Methyl 1-(2,2-dimethoxyethyl)-2-(3-pyridyl)pyrimidin-6(1H)-one-5-carboxylate.

[0113] Dimethyl methoxymethylenemalonate and N-(2,2-dimethoxyethyl)-3-pyridinecarboxamidine were subjected to a procedure similar to that described in Example 1.b. Chromatography, with methanol:diethyl ether (gradient, 0:100, 10:90) as the eluent, gave methyl 1-(2,2-dimethoxy-ethyl)-2-(3-pyridyl)pyrimidin-6(1H)-one-5-carboxylate as an oil; MS: m/z=320(M+1); TLC: R_f=0.4, methanol:diethyl ether (5:95).

d. 5-Benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetaldehyde dimethyl acetal.

[0114] To a solution of methyl 1-(2,2-dimethoxyethyl)-2-(3-pyridyl)pyrimidin-6(1H)-one-5-carboxylate in pyridine (200 mL) was added lithium iodide (48 g) and the mixture heated at 100 °C for 3 h. The solvent was evaporated and residual pyridine removed by addition of toluene (200 mL) followed by evaporation. The resulting material was diluted with water and the pH adjusted to pH 5. The product was extracted into ethyl acetate, dried and evaporated. The resulting crude acid was dissolved in dioxane (260 mL), and to this solution were added triethylamine (22 mL) and diphenylphosphoryl azide (18 mL). The solution was heated at 100 °C for 2 h and benzyl alcohol (17 mL) was added. The resulting solution was heated overnight at 100 °C, the solvent was removed, and the resulting material was redissolved in ethyl acetate. The solution was washed (1 N sodium hydroxide, water, brine), dried, and evaporated. Chromatography, with ethyl acetate:methanol (gradient, 100:0, 70:30) as the eluent, gave 5-benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetaldehyde dimethyl acetal as an oil; MS: m/z=365(M+1).

e. 5-Benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetaldehyde.

[0115] 5-Benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetaldehyde dimethyl acetal was subjected to a procedure similar to that described in Example 8.f. to provide 5-benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetaldehyde as a tan solid; TLC: R_f = 0.45, methanol:dichloromethane (5:95).

f. 5-Benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetic acid.

[0116] 5-Benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetaldehyde was subjected to a procedure similar to that described in Example 1.f. to provide a white solid which was washed with ether to give 5-benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetic acid as a white solid; NMR: 13.5 (s broad, 1), 9.11 (s, 1), 8.73 (m, 2), 8.51 (s, 1), 7.96 (d, 1), 7.47 (m, 5), 5.20 (s, 2), 4.57 (s, 2).

g. 2-[5-Benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide.

[0117] 5-Benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinylacetic acid and 3-amino-1,1,1-trifluoro-2-hydroxy-1-isopropylpropylamine were subjected to a procedure similar to that described in Example 1.g. to provide 2-[5-benzoyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide as a white solid; MS: m/z=528(M+1).

luoro-4-methyl-2-pentanol hydrochloride was subjected to a procedure similar to that described in Example 1.g. Chromatography, with methanol:dichloromethane (5:95) as the eluent, gave 2-[5-benzyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropyl-propyl)acetamide as a white solid; MS: $m/z=534(M+1)$.

EXAMPLE 14

2-[5-Amino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

- [0118] 2-[5-Benzyloxycarbonylamino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide was subjected to a procedure similar to that described in Example 15 to provide 2-[5-amino-6-oxo-2-(3-pyridyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; 300 MHz NMR (DMSO/D₂O): 8.56 (m,2), 7.78 (m,1), 7.41 (m,1), 7.34 (s,1), 4.56 (d,1), 4.48 (d,1), 3.94 (m,1), 2.13 (m,1), 0.75 (d,3), 0.64 (d,3).

Analysis for C ₁₇ H ₁₈ F ₃ N ₅ O ₃ :			
Calculated:	C, 51.38;	H, 4.56;	N, 17.62
Found:	C, 50.79;	H, 4.68;	N, 17.35

EXAMPLE 15

2-[5-Amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

- [0119] To a solution of 2-[5-benzyloxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (1.03 g) in ethanol (45 mL) and tetrahydrofuran (22 mL) was added 10% (w/w) palladium on carbon (15% by weight). This mixture was shaken under a hydrogen atmosphere (3 bar), filtered, and additional 10% (w/w) palladium on carbon was added. The mixture was again shaken under hydrogen (3 bar). The solution was filtered and the solvent evaporated. The resulting material was purified by chromatography, with methanol:dichloromethane (gradient, 5:95, 12:88) as the eluent, to give a white solid which was collected and washed with ether to provide 2-[5-amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid; mp 200-203 °C; 300 MHz NMR (DMSO/D₂O): 7.76 (d,1), 7.47 (m,2), 7.31 (s,1), 7.20 (t,2), 4.56 (d,1), 4.37 (d,1), 4.01 (m,1), 2.20 (m,1), 0.82 (d,3), 0.72 (d,3).

Analysis for C ₁₈ H ₁₈ F ₄ N ₄ O ₃ • 0.75 H ₂ O:			
Calculated:	C, 50.53;	H, 4.59;	N, 13.09
Found:	C, 50.41;	H, 4.69;	N, 13.09

EXAMPLE 16

2-[5-(N,N-Dimethylaminosulfonylamino)-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

- [0120] To a solution of 2-(5-amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (0.298 g) in tetrahydrofuran (3 mL) containing sodium carbonate (0.161 g) was added dimethylsulfonyl chloride (0.1 mL). The mixture was stirred at 60 °C overnight and more sodium carbonate (0.170 g) and dimethylsulfonyl chloride (0.1 mL) were added. The solution was allowed to stir for 1 day. The reaction was diluted with ethyl acetate and the solution was washed (1 N hydrochloric acid, water, 10% aqueous sodium bicarbonate, brine), dried, and evaporated. The resulting oil was purified by chromatography, with tetrahydrofuran:dichloromethane (10:90) as the eluent, to provide 2-[5-(N,N-dimethylsulfonylamino)-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid (0.117 g); NMR (DMSO/D₂O): 9.29 (s,1), 8.84 (d,1), 7.99

(s,1), 7.48 (m,5), 2.72 (s,3), 2.16 (m,1), 0.89 (d,3), 0.83 (d,3).

5

Analysis for $C_{20}H_{24}F_3N_5O_5S$:			
Calculated:	C, 47.7;	H, 4.80;	N, 13.9
Found:	C, 47.6;	H, 4.93;	N, 13.5

10

EXAMPLE 17

2-(5-Methylsulfonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

15

[0121] To a solution of 2-(5-amino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (0.345 g) in tetrahydrofuran (7mL) and triethylamine (0.36mL) at 0 °C was added methanesulfonyl chloride (0.19 mL) and the resulting solution was allowed to stir overnight. The mixture was diluted with ethyl acetate and the solution washed (1 N hydrochloric acid, water, 5% sodium bicarbonate). The organic layer was dried and evaporated. The residue was dissolved in tetrahydrofuran (1 mL), and to it was added 3 N aqueous potassium hydroxide (0.07 mL). The solution was allowed to stir for 12 h. The reaction was quenched by addition of 1 N hydrochloric acid and the product was extracted into ethyl acetate. The organic layer was washed (water, brine), and the solution was dried and evaporated. The resulting material was purified by chromatography, with tetrahydrofuran:dichloromethane (10:90) as the eluent, to provide 2-(5-methylsulfonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide as a white solid (0.146 g); NMR (DMSO/ D_2O): 9.35 (s,1), 8.84 (d,1), 8.01 (s,1), 7.50 (m,5), 4.68 (m,2), 4.55 (d,1), 3.10 (s,3), 2.16 (m,1), 0.89 (d,3), 0.83 (d,3).

25

30

Analysis for $C_{19}H_{21}F_3N_4O_5S$:			
Calculated:	C, 47.7;	H, 4.52;	N, 11.7
Found:	C, 47.7;	H, 4.62;	N, 11.5

35

EXAMPLES 18-25

[0122] Using a procedure similar to that described in Example 1, the following compounds of formula I wherein R^0 is isopropyl, R is benzyloxycarbonyl, and R^6 is the indicated group were prepared by oxidation of the corresponding alcohols of formula II.

40

[0123] Example 18: R^6 =4-nitrophenyl: Chromatography solvent: ethyl acetate:dichloromethane (gradient, 30:70 to 50:50); TLC: R_f =0.6, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.44 (s,1), 8.26 (d,2), 7.70 (d,2), 7.36 (m,5), 5.15 (s,2), 4.51 (m,2), 3.97 (d,1), 2.14 (m,1), 0.74 (d,3), 0.63 (d,3).

45

Analysis for $C_{26}H_{24}F_3N_5O_7$:			
Calculated:	C, 54.2;	H, 4.20;	N, 12.1
Found:	C, 54.1;	H, 4.23;	N, 12.2

50

[0124] Example 19: R^6 =4-trifluoromethylphenyl: Chromatography solvent: ethyl acetate:hexane (50:50); TLC: R_f =0.41, diethyl ether; NMR (DMSO/ D_2O): 9.07 (s,1), 8.79 (d,1), 8.48 (s,1), 7.84 (d,2), 7.69 (d,2), 7.39 (m,5), 5.17 (s,2), 4.80 (m,3), 2.12 (m,1), 0.83 (d,3), 0.77 (d,3).

55

Analysis for $C_{27}H_{24}F_6N_4O_5$:			
Calculated:	C, 54.2;	H, 4.04;	N, 9.36
Found:	C, 54.1;	H, 4.07;	N, 9.41.

5
10 [0125] Example 20: $R^6=3,5$ -difluorophenyl: Chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 5:95 to 10:90); TLC: $R_f=0.31$, tetrahydrofuran:dichloromethane (8:92); NMR (DMSO/ D_2O): 8.45 (s,1), 7.20-7.48 (m,8), 5.18 (s,2), 4.67 (d,1), 4.43 (d,1), 4.01 (d,1), 2.22 (m,1), 0.84 (d,3), 0.73 (d,3).

Analysis for $C_{26}H_{23}F_5N_4O_5$:			
Calculated:	C, 55.1;	H, 4.09;	N, 9.89
Found:	C, 55.2;	H, 4.13;	N, 9.89

15
20 [0126] Example 21: $R^6=4$ -methoxyphenyl: Chromatography solvent: ethyl acetate:dichloromethane (20:80); TLC: $R_f=0.28$, methanol:dichloromethane (20:80); NMR: 8.95 (s,1), 8.85 (d,1), 8.43 (s,1), 7.40 (m,7), 6.99 (d,2), 5.18 (s,2), 4.67 (t,1), 4.58 (d,1), 3.81 (s,3), 2.16 (m,1), 0.90 (d,3), 0.84 (d,3).

Analysis for $C_{27}H_{27}F_3N_4O_6$:			
Calculated:	C, 57.9;	H, 4.85;	N, 9.99
Found:	C, 57.6;	H, 4.91;	N, 9.94.

25
30
35 [0127] Example 22: $R^6=4$ -chlorophenyl: Chromatography solvent: methanol:dichloromethane (5:95); TLC: $R_f=0.4$, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.39 (s,1), 7.36 (m,9), 5.14 (s,2), 4.49 (m,2), 3.97 (d,2), 2.16 (m,1), 0.77 (d,3), 0.66 (d,3).

Analysis for $C_{26}H_{24}ClF_3N_4O_5$:			
Calculated:	C, 55.2;	H, 4.28;	N, 9.91
Found:	C, 55.1;	H, 4.23;	N, 9.91.

40
45 [0128] Example 23: $R^6=3,5$ -bis(trifluoromethyl)phenyl: Chromatography solvent: ethyl acetate:dichloromethane (gradient, 30:70, 50:50); TLC: $R_f=0.6$, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.44 (s,1), 8.26 (d,2), 7.70 (d,2), 7.36 (m,5), 5.15 (s,2), 4.51 (m,2), 3.97 (d,1), 2.14 (m,1), 0.74 (d,3), 0.63 (d,3).

Analysis for $C_{26}H_{24}F_3N_5O_7$:			
Calculated:	C, 54.2;	H, 4.20;	N, 12.1
Found:	C, 54.1;	H, 4.23;	N, 12.2

50
55 [0129] Example 24: $R^6=cyclohexyl$: Chromatography solvent: ethyl acetate:dichloromethane (15:85); TLC:

$R_f=0.55$, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.22 (s,1), 7.32 (m,5), 5.10 (s,2), 4.79 (m,2), 4.01 (d,1), 2.50 (m,1), 2.25 (m,1), 1.7-1.14 (m,10), 0.92 (d,3), 0.75 (d,3).

Analysis for $C_{26}H_{31}F_3N_4O_5$:			
Calculated:	C, 58.2;	H, 5.82;	N, 10.4
Found:	C, 57.9;	H, 4.84;	N, 10.3

[0130] Example 25: R^6 =isopropyl: Chromatography solvent: ethyl acetate:dichloromethane (gradient, 10:90 to 25:75); TLC: $R_f=0.4$, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.23 (s,1), 7.32 (m,5), 5.10 (s,2), 4.78 (m,2), 4.00 (d,1), 2.75 (m,1), 2.20 (m,1), 1.11 (d,6), 0.88 (d,3), 0.73 (d,3).

Analysis for $C_{23}H_{27}F_3N_4O_5$:			
Calculated:	C, 55.6;	H, 5.48;	N, 11.3
Found:	C, 55.6;	H, 5.53;	N, 11.3

[0131] The intermediate alcohols used in Examples 18-25 were prepared as follows.

EXAMPLES 18.a.-25.a.

[0132] Using a procedure similar to that described in Example 8.a., the following imidates of formula IV wherein R^7 is ethyl and R^6 is the indicated group were prepared from the indicated nitrile, and isolated as their hydrochloride salts.

[0133] Example 18.a: R^6 =4-nitrophenyl, from 4-nitrobenzonitrile. Purification method: trituration from diethyl ether; MS: $m/z=195(M+1-HCl)$.

[0134] Example 19.a: R^6 =4-trifluoromethylphenyl, from 4-trifluoromethylbenzonitrile. Purification method: trituration from diethyl ether; MS: $m/z=218(M+1-HCl)$.

[0135] Example 20.a: R^6 =3,5-difluorophenyl, from 3,5-difluorobenzonitrile. Purification method: trituration from diethyl ether; MS: $m/z=186(M+1-HCl)$.

[0136] Example 21.a: R^6 =4-methoxyphenyl, from 4-methoxybenzonitrile. Purification method: trituration from diethyl ether; MS: $m/z=180(M+1-HCl)$.

[0137] Example 22.a: R^6 =4-chlorophenyl, from 4-chlorobenzonitrile. Purification method: trituration from diethyl ether; MS: $m/z=169(M+1-HCl)$.

[0138] Example 23.a: R^6 =3,5-bis(trifluoromethyl)phenyl, from 3,5-bis(trifluoromethyl)benzonitrile. Purification method: trituration from diethyl ether; MS: $m/z=286(M+1-HCl)$.

[0139] Example 24.a: R^6 =cyclohexyl, from cyclohexanecarbonitrile. Purification method: trituration from diethyl ether; MS: $m/z=156(M+1-HCl)$.

[0140] Example 25.a: R^6 =isopropyl, from isopropylnitrile. Purification method: trituration from diethyl ether; MS: $m/z=115(M+1-HCl)$.

EXAMPLES 18.b.-25.b.

[0141] Using a procedure similar to that described in Example 8.b. the following compounds of formula V wherein R^8 is dimethoxymethyl and R^6 is the indicated group were prepared from the corresponding imidate hydrochlorides of formula IV and aminoacetaldehyde dimethyl acetal.

[0142] Example 18.b: R^6 =4-nitrophenyl; MS: $m/z=254(M+1)$.

[0143] Example 19.b: R^6 =4-trifluoromethylphenyl; MS: $m/z=277(M+1)$.

[0144] Example 20.b: R^6 =3,5-difluorophenyl; MS: $m/z=245(M+1)$.

[0145] Example 21.b: R^6 =4-methoxyphenyl; MS: $m/z=239(M+1)$.

[0146] Example 22.b: R^6 =4-chlorophenyl; MS: $m/z=243(M+1)$.

[0147] Example 23.b: R^6 =3,5-bis(trifluoromethyl)phenyl; MS: $m/z=345(M+1)$.

[0148] Example 24.b: R^6 =cyclohexyl; MS: $m/z=215(M+1)$.

[0149] Example 25.b.: R⁶=isopropyl; MS: m/z=175(M+1).

EXAMPLES 18.c.-25.c.

5 [0150] Using a procedure similar to that described in Example 1.b., the following pyrimidone esters of formula VI wherein R⁸ is dimethoxymethyl and R⁶ is the indicated group were prepared from the corresponding amidines of formula V and dimethylmethoxy methylenemalonate.

[0151] Example 18.c.: R⁶=4-nitrophenyl; yellow solid; TLC: R_f=0.3, diethyl ether; MS: m/z=364(M+1).

[0152] Example 19.c.: R⁶=4-trifluoromethylphenyl; TLC: R_f=0.37, diethyl ether; MS: m/z=387(M+1).

10 [0153] Example 20.c.: R⁶=3,5-difluorophenyl; TLC: R_f=0.28, diethyl ether:hexane (80:20); MS: m/z=355(M+1).

[0154] Example 21.c.: R⁶=4-methoxyphenyl; TLC: R_f=0.37, ethyl acetate; MS: m/z=349(M+1).

[0155] Example 22.c.: R⁶=4-chlorophenyl; TLC: R_f=0.5, ethyl acetate; MS: m/z=353(M+1).

[0156] Example 23.c.: R⁶=3,5-bis(trifluoromethyl)phenyl; TLC: R_f=0.6, ethyl acetate; MS: m/z=455(M+1).

[0157] Example 24.c.: R⁶=cyclohexyl; TLC: R_f=0.6, diethyl ether; MS: m/z=325(M+1).

15 [0158] Example 25.c.: R⁶=2-propyl; TLC: R_f=0.35, diethyl ether; MS: m/z=285(M+1).

EXAMPLES 18.d.-25.d.

20 [0159] Using a procedure similar to that described in Example 11.d., the pyrimidone acids of formula VII wherein R⁸ is dimethoxymethyl and R⁶ is the indicated group were prepared from the corresponding pyrimidone esters of formula VI.

[0160] Example 18.d.: R⁶=4-nitrophenyl; yellow solid; TLC: R_f=0.1, ethyl acetate; MS: m/z=350(M+1).

[0161] Example 19.d.: R⁶=4-trifluoromethylphenyl; TLC: R_f=0.24, methanol:dichloromethane (10:90); MS: m/z=373(M+1).

25 [0162] Example 20.d.: R⁶=3,5-difluorophenyl; TLC: R_f=0.40, methanol:dichloromethane (15:85); MS: m/z=341(M+1).

[0163] Example 21.d.: R⁶=4-methoxyphenyl; TLC: R_f=0.45, methanol:dichloromethane (5:95); MS: m/z=335(M+1).

[0164] Example 22.d.: R⁶=4-chlorophenyl; TLC: R_f=0.2, methanol:dichloromethane (10:90); MS: m/z=339(M+1).

30 [0165] Example 23.d.: R⁶=3,5-bis(trifluoromethyl)phenyl; TLC: R_f=0.3, methanol:dichloromethane (20:80); MS: m/z=441(M+1).

[0166] Example 24.d.: R⁶=cyclohexyl; TLC: R_f=0.2, diethyl ether; MS: m/z=311(M+1).

[0167] Example 25.d.: R⁶=2-propyl; TLC: R_f=0.8, diethyl ether; MS: m/z=376(M+1).

EXAMPLES 18.e.-25.e

35 [0168] Using a procedure similar to that described in Example 1.d, the following compounds of formula IX wherein R⁸ is dimethoxymethyl and R⁶ is the indicated group were prepared from the corresponding acids of formula VII.

[0169] Example 18.e.: R⁶=4-nitrophenyl; white solid; TLC: R_f=0.8, ethyl acetate; NMR: 9.08 (s,1), 8.49 (s,1), 8.34 (d,2), 7.85 (d,2), 7.38 (m,5), 5.21 (s,2), 4.51 (t,1), 4.02 (d,2).

40 [0170] Example 19.e.: R⁶=4-trifluoromethylphenyl; TLC: R_f=0.36, ethyl acetate:dichloromethane (5:95); MS: m/z=478(M+1).

[0171] Example 20.e.: R⁶=3,5-difluorophenyl; TLC: R_f=0.74, diethyl ether:hexane (80:20); MS: m/z=446(M+1).

[0172] Example 21.e.: R⁶=4-methoxyphenyl; TLC: R_f=0.71, ethyl acetate; MS: m/z=440(M+1).

[0173] Example 22.e.: R⁶=4-chlorophenyl; TLC: R_f=0.8, diethyl ether; MS: m/z=444(M+1).

45 [0174] Example 23.e.: R⁶=3,5-bis(trifluoromethyl)phenyl; TLC: R_f=0.61, ethyl acetate:dichloromethane (5:95); MS: m/z=546(M+1).

[0175] Example 24.e.: R⁶=cyclohexyl; oil; TLC: R_f=0.8, diethyl ether; MS: m/z=416(M+1).

[0176] Example 25.e.: R⁶=isopropyl; TLC: R_f=0.8, diethyl ether; MS: m/z=376(M+1).

50 EXAMPLES 18.f.-25.f.

[0177] Using a procedure similar to that described in Example 8.f., the following aldehydes of formula X wherein R⁶ is the indicated group were prepared from the corresponding compounds of formula IX.

[0178] Example 18.f.: R⁶=4-nitrophenyl; yellow solid; TLC: R_f=0.5, diethyl ether; MS: m/z=409(M+1).

55 [0179] Example 19.f.: R⁶=4-trifluoromethylphenyl; TLC: R_f=0.27, ethyl acetate:dichloromethane (5:95); MS: m/z=432(M+1).

[0180] Example 20.f.: R⁶=3,5-difluorophenyl; TLC: R_f=0.38, diethyl ether:hexane (70:30); MS: m/z=400(M+1).

[0181] Example 21.f.: R⁶=4-methoxyphenyl; TLC: R_f=0.65, ethyl acetate; MS: m/z=394(M+1).

[0182] Example 22.f.: R⁶=4-chlorophenyl; TLC: R_F=0.5, diethyl ether; MS: m/z=398(M+1).

[0183] Example 23.f.: R⁶=3,5-bis(trifluoromethyl)phenyl; TLC: R_F=0.72, methanol:dichloromethane (5:95); MS: m/z=500(M+1).

[0184] Example 24.f.: R⁶=cyclohexyl; TLC: R_F=0.3, ethyl acetate; MS m/z=369(M+1).

5 [0185] Example 25.f.: R⁶=isopropyl; TLC: R_F=0.2, diethyl ether; MS: m/z=330(M+1).

EXAMPLES 18.g.-25.g.

[0186] Using a procedure similar to that described in Example 1.f., the following acids of formula III wherein R⁶ is the indicated group were prepared from the corresponding aldehydes of structure X.

[0187] Example 18.g.: R⁶=4-nitrophenyl; yellow solid; TLC: R_F=0.1, ethyl acetate; MS: m/z=425(M+1).

[0188] Example 19.g.: R⁶=4-trifluoromethylphenyl; TLC: R_F=0.3, methanol:dichloromethane (20:80); MS: m/z=448(M+1).

15 [0189] Example 20.g.: R⁶=3,5-difluorophenyl; TLC: R_F=0.35, methanol:dichloromethane (20:80); MS: m/z=416(M+1).

[0190] Example 21.g.: R⁶=4-methoxyphenyl; TLC: R_F=0.27, methanol:dichloromethane (20:80); MS: m/z=410(M+1).

[0191] Example 22.g.: R⁶=4-chlorophenyl; TLC: R_F=0.2, methanol:dichloromethane (10:90); MS: m/z=414(M+1).

20 [0192] Example 23.g.: R⁶=3,5-bis(trifluoromethyl)phenyl; TLC: R_F=0.30, methanol:dichloromethane (10:90); MS: m/z=516(M+1).

[0193] Example 24.g.: R⁶=cyclohexyl; TLC: R_F=0.1, ethyl acetate; MS: m/z=385(M+1).

[0194] Example 25.g.: R⁶=isopropyl; TLC: R_F=0.2, ethyl acetate; MS: m/z=345(M+1).

EXAMPLES 18.h.-25.h.

25 [0195] Using a procedure similar to that described in Example 1.g., the following alcohols of formula II wherein R is benzyloxycarbonyl and R⁶ is the indicated group were prepared from the corresponding acids of formula III with exceptions as noted.

30 [0196] Example 18.h.: R⁶=4-nitrophenyl; Purified by chromatography, with methanol:dichloromethane (gradient, 3:97, 10:90) as the eluent, to give a yellow solid; TLC: R_F=0.2, methanol:dichloromethane (3:97); MS: m/z=578(M+1).

[0197] Example 19.h.: R⁶=4-trifluoromethylphenyl; TLC: R_F=0.65, diethyl ether; MS: m/z=601(M+1).

[0198] Example 20.h.: R⁶=3,5-difluorophenyl; TLC: R_F=0.67, tetrahydrofuran:dichloromethane (15:85); MS: m/z=569(M+1).

[0199] Example 21.h.: R⁶=4-methoxyphenyl; TLC: R_F=0.63, ethyl acetate; MS: m/z=563(M+1).

35 [0200] Example 22.h.: R⁶=4-chlorophenyl; TLC: R_F=0.5, ethyl acetate; MS: m/z=567(M+1).

[0201] Example 23.h.: R⁶=3,5-bis(trifluoromethyl)phenyl; TLC: R_F=0.61, ethyl acetate; MS: m/z=669(M+1).

[0202] Example 24.h.: R⁶=cyclohexyl; TLC: R_F=0.6, methanol:dichloromethane (5:95); MS: m/z=539(M+1).

[0203] Example 25.h.: R⁶=isopropyl; TLC: R_F=0.45, diethyl ether; MS: m/z=499(M+1).

40 EXAMPLES 26-31

[0204] Using a procedure similar to that described in Example 6, the following compounds of formula I wherein R⁰ is isopropyl, R is hydrogen, and R⁶ is the indicated group were prepared from the corresponding compounds of formula I wherein R is benzyloxycarbonyl.

45 [0205] Example 26: R⁶=4-trifluoromethylphenyl; Chromatography solvent: ethyl acetate; TLC: R_F=0.31, ethyl acetate; NMR (DMSO/D₂O): 7.71 (d,2), 7.60 (d,2), 7.35 (s,1), 4.55 (m,2), 4.04 (d,2), 2.14 (m,1), 0.76 (d,3), 0.63 (d,3).

50

Analysis for C ₁₉ H ₁₈ F ₆ N ₄ O ₃ · 0.5 H ₂ O:			
Calculated:	C, 48.2;	H, 4.05;	N, 11.8
Found:	C, 48.1;	H, 4.38;	N, 11.6

55

[0206] Example 27: R⁶=3,5-difluorophenyl; Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 12:88); TLC: R_F=0.18, tetrahydrofuran:dichloromethane (10:90); 300 MHz NMR (DMSO/D₂O): 7.35 (m,1), 7.31 (s,1), 7.14 (m,2), 4.65 (d,1), 4.40 (d,1), 4.02 (m,1), 2.22 (m,1), 0.85 (d,3), 0.74 (d,3).

Analysis for $C_{19}H_{18}F_6N_4O_3 \cdot 0.5 H_2O$:			
Calculated:	C, 48.2;	H, 4.05;	N, 11.8
Found:	C, 48.1;	H, 4.38;	N, 11.6

[0207] **Example 28:** R^6 =4-methoxyphenyl: Chromatography solvent: ethyl acetate; TLC: R_f =0.24, ethyl acetate; NMR (DMSO/ D_2O): 7.54 (d,2), 7.53 (s,1), 6.91 (d,2), 4.50 (m,2), 4.01 (d,2), 3.76 (s,3), 2.20 (m,1), 0.82 (d,3), 0.75 (d,3).

Analysis for $C_{19}H_{21}F_3N_4O_4$:			
Calculated:	C, 53.5;	H, 4.96;	N, 13.1
Found:	C, 53.5;	H, 5.04;	N, 13.1

[0208] **Example 29:** R^6 =isopropyl: NMR (DMSO/ D_2O): 7.23 (s,1), 4.75 (m,2), 4.03 (d,2), 2.72 (m,1), 2.22 (m,1), 1.06 (d,6), 0.89 (d,3), 0.75 (d,3).

Analysis for $C_{19}H_{21}F_3N_4O_3$:			
Calculated:	C, 49.7;	H, 5.84;	N, 15.5
Found:	C, 49.5;	H, 5.84;	N, 15.2

[0209] **Example 30:** R^6 =3,5-bis(trifluoromethyl)phenyl: NMR: 8.91 (d,1), 8.24 (s,1), 8.16 (s,2), 7.36 (s,1), 5.44 (s,2), 4.65 (m,3), 2.13 (m,1), 0.83 (d,3), 0.80 (d,3).

Analysis for $C_{20}H_{17}F_9N_4O_3$:			
Calculated:	C, 45.1;	H, 3.22;	N, 10.5
Found:	C, 45.6;	H, 3.37;	N, 10.7

[0210] **Example 31:** R^6 =cyclohexyl: Chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f =0.17, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 7.22 (s,1), 4.81 (m,2), 4.02 (d,1), 2.27 (m,2), 1.66 (m,5), 1.38 (m,2), 1.15 (m,3), 0.92 (d,3), 0.78 (d,3).

Analysis for $C_{18}H_{25}F_3N_4O_3 \cdot 0.5 H_2O$:			
Calculated:	C, 52.6;	H, 6.37;	N, 13.6
Found:	C, 52.6;	H, 6.26;	N, 13.5

EXAMPLES 32-33

[0211] Using a procedure similar to that described in Example 12, the following compounds of formula I wherein R^0 is isopropyl, R is hydrogen, and R^6 is the indicated group were prepared from the corresponding compounds of formula

I wherein R is benzyloxycarbonyl.

[0212] **Example 32:** R⁶=4-chlorophenyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 20:80); TLC: R_F=0.2, methanol:dichloromethane (5:95); NMR (DMSO/D₂O): 7.34 (m,5), 4.49 (m,2), 3.99 (d,2), 2.15 (m,1), 0.78 (d,3), 0.67 (d,3).

Analysis for C ₁₈ H ₁₈ ClF ₃ N ₄ O ₃ :			
Calculated:	C, 50.2;	H, 4.21;	N, 13.0
Found:	C, 50.0;	H, 4.44;	N, 12.8

[0213] **Example 33:** R⁶=4-nitrophenyl: Chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 30:70, 50:50); NMR (DMSO/D₂O): 8.21 (d,2), 7.63 (d,2), 7.35 (s,1), 4.49 (m,2), 3.99 (d,2), 2.14 (m,1), 0.76 (d,3), 0.63 (d,3).

Analysis for C ₁₈ H ₁₈ F ₃ N ₅ O ₅ • 1.1 H ₂ O:			
Calculated:	C, 46.9;	H, 4.41;	N, 15.2
Found:	C, 46.9;	H, 4.25;	N, 15.2

EXAMPLE 34

2-[5-Amino-2-(4-aminophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0214] 2-[5-Benzyloxycarbonylamino-2-(4-nitrophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide was subjected to a procedure similar to that described in Example 6 to give a brown solid; chromatography solvent: methanol: dichloromethane (gradient, 5:95, 10:90); NMR (DMSO/D₂O): 7.29 (s,1), 7.01 (d,2), 6.49 (d,2), 4.52 (m,2), 4.10 (d,2), 2.25 (m,1), 0.88 (d,3), 0.80 (d,3).

Analysis for C ₁₈ H ₂₀ F ₃ N ₅ O ₃ :			
Calculated:	C, 52.6;	H, 4.90;	N, 17.0
Found:	C, 52.4;	H, 4.96;	N, 16.7

EXAMPLE 35

2-[5-Acetylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0215] To a solution of 2-[5-amino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)-acetamide (0.3 g) in tetrahydrofuran (7 mL) was added sodium carbonate (0.4 g) and the mixture cooled in an ice bath to 0 °C. Acetyl chloride (0.11 mL) was added and the solution warmed to room temperature and let stir for 1 h. The mixture was poured into ethyl acetate and washed (1 N HCl, a saturated solution of sodium carbonate, and H₂O). The resulting solution was dried and the solvent removed by evaporation. The resulting material was purified by chromatography, eluting with methanol:dichloromethane (gradient, 5:95, 12:88) to give the title compound (0.46 g) as a light yellow powder; TLC: R_F=0.47, methanol:dichloromethane (5:95); NMR (DMSO/D₂O): 8.77 (s,1), 7.81 (d,1), 7.36 (d,1), 7.09 (t,1), 4.83 (m,2), 4.09 (d,1), 2.26 (m,1), 2.13 (s,3), 0.93 (d,3), 0.80 (d,3).

Analysis for $C_{18}H_{19}F_3N_4O_4S \cdot 0.25 H_2O$:			
Calculated:	C, 48.2;	H, 4.38;	N, 12.5
Found:	C, 48.0;	H, 4.35;	N, 12.4

[0216] This procedure will be referred to as Acylation Method A.

EXAMPLES 36-74

[0217] The following compounds of formula I wherein R^0 is isopropyl and R and R^6 have the indicated values were prepared from the corresponding compounds of formula I wherein R is hydrogen using Acylation Method A and the required acyl chloride.

[0218] Example 36: R=acetyl, R^6 =4-fluorophenyl; chromatography solvent: methanol:dichloromethane (gradient, 0:100, 7:93); TLC: R_f =0.27, ethyl acetate:dichloromethane (35:65); NMR (DMSO/ D_2O): 9.51 (s,1), 8.8 (s,1), 7.55 (m,2), 7.26 (m,2), 4.66 (broad d,1), 4.52 (m,2), 4.01 (d,1), 2.24 (m,1), 2.14 (s,3), 0.85 (d,3), 0.75 (d,3).

Analysis for $C_{20}H_{20}F_4N_4O_4 \cdot 1.0 H_2O$:			
Calculated:	C, 50.6;	H, 4.67;	N, 11.8
Found:	C, 50.4;	H, 4.63;	N, 11.8

[0219] Example 37: R=acetyl, R^6 =3-pyridyl; tan solid; chromatography solvent: methanol:dichloromethane (gradient, 5:95, 10:90); TLC: R_f =0.4, methanol:dichloromethane (10:90); 300 MHz NMR (DMSO/ D_2O): 8.58 (m,2), 7.81 (m,1), 7.40 (m,1), 7.34 (s,1), 4.59 (m,2), 3.98 (d,1), 2.16 (m,1), 0.90 (d,3), 0.68 (d,3).

Analysis for $C_{19}H_{20}F_3N_5O_4 \cdot 0.7 H_2O$:			
Calculated:	C, 50.5;	H, 4.77;	N, 15.5
Found:	C, 50.6;	H, 4.75;	N, 15.3

[0220] Example 38: R=methoxycarbonyl, R^6 =3-pyridyl; white solid; purified by trituration from diethyl ether:hexane (50:50); TLC: R_f =0.6, methanol:dichloromethane (10:90); NMR (DMSO/ D_2O): 8.62 (m,2), 8.44 (s,1), 7.86 (m,1), 7.45 (m,1), 4.62 (m,2), 3.95 (d,1), 3.66 (s,3), 2.15 (m,1), 0.78 (d,3), 0.64 (d,3).

Analysis for $C_{19}H_{20}F_3N_5O_5 \cdot 1.0 H_2O$:			
Calculated:	C, 48.2;	H, 4.68;	N, 14.8
Found:	C, 48.5;	H, 4.87;	N, 14.2

[0221] Example 39: R=4-methoxyphenoxy carbonyl, R^6 =3-pyridyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f =0.4, methanol:dichloromethane (7:93); 300 MHz NMR (DMSO/ D_2O): 8.65 (m,2), 8.43 (s,1), 7.88 (m,1), 7.44 (m,1), 7.10 (d,2), 6.92 (d,2), 4.55 (m,2), 3.97 (d,1), 3.62 (s,3), 2.17 (m,1), 0.78 (d,3), 0.67 (d,3).

Analysis for $C_{25}H_{24}F_3N_5O_6$:			
Calculated:	C, 54.8;	H, 4.41;	N, 12.8
Found:	C, 54.2;	H, 4.43;	N, 12.5

[0222] **Example 40:** R=methoxycarbonyl, R^6 =4-methoxyphenyl; white solid; purified by recrystallization from ethyl acetate; TLC: R_f =0.46, ethyl acetate; NMR (DMSO/ D_2O): 8.85 (d,1), 8.79 (s,1), 7.42 (d,2), 6.99 (d,2), 4.66 (t,1), 4.58 (dd,1), 3.81 (s,1), 3.68 (s,3), 2.16 (m,1), 0.90 (d,3), 0.84 (d,3).

Analysis for $C_{21}H_{23}F_3N_4O_6$:			
Calculated:	C, 52.1;	H, 4.79;	N, 11.6
Found:	C, 51.9;	H, 4.82;	N, 11.5

[0223] **Example 41:** R=4-fluorophenoxycarbonyl, R^6 =phenyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (gradient, 25:75, 40:60); TLC: R_f =0.8, tetrahydrofuran:dichloromethane (20:80); NMR (DMSO/ D_2O): 8.42 (s,1), 7.48 (m,5), 7.25 (d,4), 4.65 (d,1), 4.48 (d,1), 4.05 (d,1), 2.23 (m,1), 0.85 (d,3), 0.77 (d,3).

Analysis for $C_{25}H_{22}F_4N_4O_5$:			
Calculated:	C, 56.2;	H, 4.15;	N, 10.5
Found:	C, 56.3;	H, 4.31;	N, 10.4

[0224] **Example 42:** R=methoxycarbonyl, R^6 =phenyl; white solid; chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 5:95 to 20:80); TLC: R_f =0.38, tetrahydrofuran:dichloromethane (10:90); NMR (DMSO/ D_2O): 8.79 (d,1), 8.43 (s,1), 7.49 (m,5), 4.60 (m,2), 4.02 (s,1), 3.67 (s,3), 2.23 (m,1), 0.80 (overlapping d,6).

Analysis for $C_{20}H_{21}F_3N_4O_5$:			
Calculated:	C, 52.9;	H, 4.66;	N, 12.3
Found:	C, 52.8;	H, 4.65;	N, 12.3

[0225] **Example 43:** R=methoxycarbonyl, R^6 =4-fluorophenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f =0.37, methanol:dichloromethane (5:95); NMR: 8.88 (s,1), 8.83 (s,1), 8.44 (s,1), 7.53 (dd,2), 7.31 (dd,2), 4.64 (t,1), 4.57 (dd,2), 3.66 (s,3), 2.15 (m,1), 0.87 (d,3), 0.82 (d,3).

Analysis for $C_{20}H_{20}F_4N_4O_5$:			
Calculated:	C, 50.9;	H, 4.27;	N, 11.9
Found:	C, 51.0;	H, 4.32;	N, 11.9

[0226] **Example 44:** R=methoxycarbonyl, R^6 =4-chlorophenyl; white solid; chromatography solvent: metha-

not:dichloromethane (5:95); TLC: $R_f=0.41$, methanol:dichloromethane (5:95); NMR: 8.88 (s,1), 8.83 (d,1), 8.45 (s,1), 7.54 (d,2), 7.49 (d,2), 4.64 (t,1), 4.57 (dd,2), 3.68 (s,3), 2.15 (m,1), 0.88 (d,3), 0.81 (d,3).

5

Analysis for $C_{20}H_{20}ClF_3N_4O_5$:			
Calculated:	C, 49.1;	H, 4.12;	N, 11.4
Found:	C, 48.8;	H, 4.16;	N, 11.3

10

[0227] Example 45: R=cyclopentyloxycarbonyl, $R^6=4$ -fluorophenyl; using cyclopentyl chloroformate; white solid; chromatography solvent: methanol:dichloromethane (5:95); NMR: 8.84 (d,1), 8.53 (s,1), 8.42 (s,1), 7.53 (dd,2), 7.31 (dd,2), 5.10 (m,1), 4.58 (m,3), 2.14 (m,1), 1.61 (m,8), 0.68 (d,3), 0.62 (d,3).

15

Analysis for $C_{24}H_{26}F_4N_4O_5$:			
Calculated:	C, 54.8;	H, 4.98;	N, 10.6
Found:	C, 54.6;	H, 4.97;	N, 10.5

20

[0228] Example 46: R=cyclopentyloxycarbonyl, $R^6=4$ -chlorophenyl; using cyclopentyl chloroformate; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: $R_f=0.45$, methanol:dichloromethane (5:95); NMR: 8.85 (d,1), 8.53 (s,1), 8.43 (s,1), 7.55 (d,2), 7.49 (d,2), 5.10 (m,1), 4.65 (t,1), 4.58 (dd,2), 2.16 (m,1), 1.71 (m,8), 0.88 (d,3), 0.83 (d,3).

25

30

Analysis for $C_{24}H_{26}ClF_3N_4O_5$:			
Calculated:	C, 53.1;	H, 4.83;	N, 10.3
Found:	C, 53.1;	H, 4.88;	N, 10.2

35

[0229] Example 47: R=cyclopentyloxycarbonyl, R^6 =phenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: $R_f=0.35$, methanol:dichloromethane (5:95); NMR: 8.84 (d,1), 8.52 (s,1), 8.43 (s,1), 7.48 (m,5), 5.10 (m,1), 4.67 (t,1), 4.55 (dd,2), 2.16 (m,1), 1.71 (m,8), 0.89 (d,3), 0.83 (d,3).

40

Analysis for $C_{24}H_{27}F_3N_4O_5$:			
Calculated:	C, 56.7;	H, 5.35;	N, 11.0
Found:	C, 56.7;	H, 5.49;	N, 10.9

45

[0230] Example 48: R=isopropoxycarbonyl, R^6 =phenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: $R_f=0.38$, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.39 (s,1), 7.43 (m,5), 4.85 (m,1), 4.52 (dd,2), 4.00 (d,1), 2.19 (m,1), 1.22 (d,6), 0.81 (d,3), 0.73 (d,3).

50

55

Analysis for $C_{22}H_{25}F_3N_4O_5 \cdot 0.1 H_2O$:			
Calculated:	C, 54.7;	H, 5.27;	N, 11.5

(continued)

Analysis for $C_{22}H_{25}F_3N_4O_5 \cdot 0.1 H_2O$:			
Found:	C, 54.2;	H, 5.50;	N, 11.2

[0231] Example 49: R=isopropoxycarbonyl, $R^6=4$ -fluorophenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: $R_f=0.33$, methanol:dichloromethane (5:95); 300 MHz NMR (DMSO/ D_2O): 8.38 (s,1), 7.50 (dd,2), 7.23 (t,2), 4.86 (m,1), 4.55 (dd,2), 3.98 (d,1), 2.15 (m,1), 1.22 (d,6), 0.80 (d,3), 0.70 (d,3).

Analysis for $C_{22}H_{24}F_4N_4O_5 \cdot 0.5 H_2O$:			
Calculated:	C, 51.9;	H, 4.95;	N, 11.0
Found:	C, 51.9;	H, 4.95;	N, 10.8

[0232] Example 50: R=cyclopentyloxycarbonyl, $R^6=2$ -thienyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (15:85); TLC: $R_f=0.28$, ethyl acetate:dichloromethane (15:85); NMR (DMSO/ D_2O): 8.40 (s,1), 7.79 (d,1), 7.38 (d,1), 7.13 (t,1), 5.09 (m,1), 4.88 (dd,2), 4.09 (d,1), 2.27 (m,1), 1.68 (m,8), 0.93 (d,3), 0.79 (d,3).

Analysis for $C_{22}H_{25}F_3N_4O_5S$:			
Calculated:	C, 51.4;	H, 4.90;	N, 10.9
Found:	C, 51.1;	H, 4.93;	N, 10.8

[0233] Example 51: R=isopropoxycarbonyl, $R^6=2$ -thienyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (15:85); TLC: $R_f=0.22$, ethyl acetate:dichloromethane (15:85); NMR (DMSO/ D_2O): 8.41 (s,1), 7.79 (d,1), 7.39 (d,1), 7.14 (t,1), 4.87 (m,2), 4.09 (d,1), 2.27 (m,1), 1.26 (d,6), 0.94 (d,3), 0.79 (d,3).

Analysis for $C_{20}H_{23}F_3N_4O_5S$:			
Calculated:	C, 49.2;	H, 4.75;	N, 11.4
Found:	C, 49.1;	H, 4.76;	N, 11.4

[0234] Example 52: R=methoxycarbonyl, $R^6=2$ -thienyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: $R_f=0.31$, methanol:dichloromethane (5:95); 300 MHz NMR (DMSO/ D_2O): 8.43 (s,1), 7.79 (d,1), 7.39 (d,1), 7.13 (dd,1), 4.88 (dd,2), 4.09 (d,1), 3.70 (s,3), 2.29 (m,1), 0.93 (d,3), 0.79 (d,3).

Analysis for $C_{18}H_{19}F_3N_4O_5S$:			
Calculated:	C, 47.0;	H, 4.16;	N, 12.2
Found:	C, 46.7;	H, 4.16;	N, 12.1

[0235] Example 53: R=isopropoxycarbonyl, $R^6=4$ -chlorophenyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (gradient, 10:90, 15:85); TLC: $R_f=0.43$, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.44 (s,1), 7.51 (m,4), 4.90 (m,1), 4.60 (dd,2), 4.02 (d,1), 2.22 (m,1), 1.26 (d,6), 0.85 (d,3), 0.73 (d,3).

Analysis for $C_{22}H_{24}ClF_3N_4O_5 \cdot 0.75 H_2O$:			
Calculated:	C, 49.8;	H, 4.85;	N, 10.6
Found:	C, 50.1;	H, 4.73;	N, 10.5

[0236] **Example 54:** R=cyclopentyloxycarbonyl, R^6 =4-methoxyphenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f =0.31, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.40 (s,1), 7.44 (d,2), 6.99 (d,2), 5.09 (m,1), 4.60 (dd,2), 4.04 (d,1), 3.81 (s,3), 2.23 (m,1), 1.69 (m,8), 0.77 (d,3), 0.66 (d,3).

Analysis for $C_{25}H_{29}F_3N_4O_6$:			
Calculated:	C, 55.8;	H, 5.43;	N, 10.4
Found:	C, 55.8;	H, 5.52;	N, 10.4

[0237] **Example 55:** R=isopropoxycarbonyl, R^6 =4-methoxyphenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f =0.28, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.42 (s,1), 7.46 (d,2), 7.00 (d,2), 4.91 (m,1), 4.68 (dd,2), 4.07 (d,1), 3.82 (s,3), 2.25 (m,1), 1.28 (d,6), 0.87 (d,3), 0.79 (d,3).

Analysis for $C_{23}H_{27}F_3N_4O_6$:			
Calculated:	C, 53.9;	H, 5.31;	N, 10.9
Found:	C, 54.1;	H, 5.43;	N, 10.

[0238] **Example 56:** R=4-fluorobenzyloxycarbonyl, R^6 =4-methoxyphenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95) TLC: R_f =0.22, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.37 (s,1), 7.41 (m,4), 7.18 (t,2), 6.93 (d,2), 5.64 (m,2), 5.11 (s,2), 4.00 (d,1), 3.75 (s,3), 2.18 (m,1), 0.81 (d,3), 0.72 (d,3).

Analysis for $C_{27}H_{26}F_4N_4O_6$:			
Calculated:	C, 56.1;	H, 4.53;	N, 9.68
Found:	C, 56.0;	H, 4.62;	N, 9.51

[0239] The intermediate 4-fluorobenzyloxy chloroformate was prepared as follows.

[0240] To a solution of triphosgene (0.53 g) in diethyl ether (2 mL) which had been cooled to 0 °C was added 4-fluorobenzyl alcohol (0.63 g) and quinoline (0.65 g). The solution was allowed to stir for 5 minutes and then additional ether was added (2 mL). A solid separated from solution and was removed by filtration. The solvent was evaporated and the resulting oil was then used without further purification.

[0241] **Example 57:** R=4-fluorobenzyloxycarbonyl, R^6 =phenyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (25:75); TLC: R_f =0.41, ethyl acetate:dichloromethane (25:75); 300 MHz NMR: 9.01 (s,1), 8.83 (d,1), 8.45 (s,1), 7.48 (m,7), 7.22 (t,2), 5.17 (s,2), 4.66 (t,1), 4.55 (dd,2), 2.15 (m,1), 0.89 (d,3), 0.82 (d,3).

Analysis for $C_{26}H_{24}F_4N_4O_5$:			
Calculated:	C, 57.0;	H, 4.41;	N, 10.2
Found:	C, 56.9;	H, 4.39;	N, 10.2

10 [0242] **Example 58:** R=4-fluorobenzyloxycarbonyl, R⁶=4-chlorophenyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (25:75); TLC: R_f=0.46, ethyl acetate:dichloromethane (25:75); 300 MHz NMR: 9.03 (s,1), 8.84 (d,1), 8.45 (s,1), 7.52 (m,6), 7.27 (t,2), 5.17 (s,2), 4.64 (t,1), 4.58 (dd,2), 2.15 (m,1), 0.88 (d,3), 0.82 (d,3).

Analysis for $C_{26}H_{23}ClF_4N_4O_5$:			
Calculated:	C, 53.6;	H, 3.97;	N, 9.61
Found:	C, 53.7;	H, 4.09;	N, 9.56

15
20 [0243] **Example 59:** R=4-fluorobenzyloxycarbonyl, R⁶=4-fluorophenyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (25:75); TLC: R_f=0.39, ethyl acetate:dichloromethane (25:75); 300 MHz NMR: 9.02 (s,1), 8.84 (d,1), 8.45 (s,1), 7.51 (m,4), 7.32 (t,2), 5.17 (s,2), 4.65 (t,1), 4.57 (dd,2), 2.15 (m,1), 0.88 (d,3), 0.82 (d,3).

Analysis for $C_{26}H_{23}F_5N_4O_5$:			
Calculated:	C, 55.1;	H, 4.09;	N, 9.89
Found:	C, 55.0;	H, 4.04;	N, 9.83

30
35 [0244] **Example 60:** R=4-fluorobenzyloxycarbonyl, R⁶=2-thienyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (25:75); TLC: R_f=0.51, ethyl acetate:dichloromethane (25:75); 300 MHz NMR: 9.04 (d,1), 9.03 (s,1), 8.43 (s,1), 7.85 (d,1), 7.50 (t,2), 7.33 (d,1), 7.18 (m,3), 5.19 (s,2), 4.88 (dd,2), 4.75 (t,1), 2.22 (m,1), 0.94 (d,3), 0.91 (d,3).

Analysis for $C_{24}H_{22}F_4N_4O_5S$:			
Calculated:	C, 52.0;	H, 4.00;	N, 10.1
Found:	C, 51.7;	H, 3.96;	N, 10.0

40
45 [0245] **Example 61:** R=ethoxycarbonyl, R⁶=phenyl; white solid; chromatography solvent: ethyl acetate:dichloromethane (25:75); TLC: R_f=0.32, methanol:dichloromethane (5:95); 300 MHz NMR: 8.83 (d,1), 8.67 (s,1), 8.44 (s,1), 7.48 (m,5), 4.64 (m,1), 4.55 (dd,2), 4.15 (m,2), 2.16 (m,1), 1.24 (t,3), 0.89 (d,3), 0.83 (d,3).

Analysis for $C_{21}H_{23}F_3N_4O_5$:			
Calculated:	C, 53.9;	H, 4.95;	N, 12.0
Found:	C, 53.8;	H, 4.97;	N, 11.9

[0246] **Example 62:** R=ethoxycarbonyl, R⁶=4-methoxyphenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.29, methanol:dichloromethane (5:95); 300 MHz NMR: 8.85 (d,1), 8.62 (s,1), 8.41 (s,1), 7.42 (d,2), 7.00 (d,2), 4.66 (m,1), 4.58 (m,2), 4.14 (q,2), 3.81 (s,3), 2.17 (m,1), 1.23 (t,3), 0.90 (d,3), 0.84 (d,3).

Analysis for C ₂₂ H ₂₅ F ₃ N ₄ O ₆ :			
Calculated:	C, 53.0;	H, 5.06;	N, 11.2
Found:	C, 53.0;	H, 5.08;	N, 11.2

[0247] **Example 63:** R=ethoxycarbonyl, R⁶=4-trifluoromethylphenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.36, methanol:dichloromethane (5:95); 300 MHz NMR: 8.80 (d,1), 8.75 (s,1), 8.46 (s,1), 7.85 (d,2), 7.70 (d,2), 4.60 (m,3), 4.15 (q,2), 2.13 (m,1), 1.24 (t,3), 0.84 (d,3), 0.77 (d,3).

Analysis for C ₂₂ H ₂₂ F ₆ N ₄ O ₅ :			
Calculated:	C, 49.3;	H, 4.13;	N, 10.4
Found:	C, 49.3;	H, 4.23;	N, 10.4

[0248] **Example 64:** R=ethoxycarbonyl, R⁶=3,5-difluorophenyl; yellow solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.33, methanol:dichloromethane (5:95); NMR (DMSO/D₂O): 8.39 (s,1), 7.35 (m,1), 7.14 (d,2), 4.52 (dd,2), 4.10 (q,2), 3.97 (d,1), 2.17 (m,1), 1.19 (t,3), 0.79 (d,3), 0.69 (d,3).

Analysis for C ₂₁ H ₂₁ F ₅ N ₄ O ₅ :			
Calculated:	C, 50.0;	H, 4.20;	N, 11.1
Found:	C, 49.7;	H, 4.38;	N, 10.9

[0249] **Example 65:** R=ethoxycarbonyl, R⁶=4-fluorophenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.28, methanol:dichloromethane (5:95); NMR (DMSO/D₂O): 8.44 (s,1), 7.53 (dd,2), 7.27 (dd,2), 4.58 (dd,2), 4.16 (q,2), 4.00 (d,1), 2.21 (m,1), 1.25 (t,3), 0.84 (d,3), 0.74 (d,3).

Analysis for C ₂₁ H ₂₂ F ₄ N ₄ O ₅ :			
Calculated:	C, 51.9;	H, 4.56;	N, 11.5
Found:	C, 51.8;	H, 4.55;	N, 11.5

[0250] **Example 66:** R=isobutoxycarbonyl, R⁶=2-thienyl; yellow solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.40, methanol:dichloromethane (5:95); NMR: 9.03 (d,1), 8.77 (s,1), 8.41 (s,1), 7.83 (d,1), 7.32 (d,1), 7.15 (t,1), 4.86 (dd,2), 4.74 (m,1), 3.87 (d,2), 2.20 (m,1), 1.90 (m,1), 0.91 (m,12).

Analysis for C ₂₁ H ₂₅ F ₃ N ₄ O ₅ S:			
Calculated:	C, 50.2;	H, 5.01;	N, 11.2

(continued)

Analysis for C ₂₁ H ₂₅ F ₃ N ₄ O ₅ S:			
Found:	C, 50.3;	H, 5.06;	N, 10.9

[0251] **Example 67:** R=ethoxycarbonyl, R⁶=2-thienyl; yellow solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.21, methanol:dichloromethane (5:95); NMR: 9.03 (d,1), 8.70 (s,1), 8.40 (s,1), 7.84 (d,1), 7.32 (d,1), 7.15 (dd,1), 4.86 (dd,2), 4.73 (t,1), 4.14 (q,2), 2.21 (m,1), 1.22 (t,3), 0.93 (d,3), 0.89 (d,3).

Analysis for C ₁₉ H ₂₁ F ₃ N ₄ O ₅ S:			
Calculated:	C, 48.1;	H, 4.46;	N, 11.8
Found:	C, 48.1;	H, 4.54;	N, 11.8

[0252] **Example 68:** R=isopropoxycarbonyl, R⁶=4-trifluoromethylphenyl; yellow solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.36, methanol:dichloromethane (5:95); NMR: 8.81 (d,1), 8.59 (s,1), 8.45 (s,1), 7.85 (d,2), 7.70 (d,2), 4.90 (m,1), 4.62 (m,3), 2.13 (m,1), 1.26 (d,6), 0.84 (d,3), 0.78 (d,3).

Analysis for C ₂₃ H ₂₄ F ₆ N ₄ O ₅ S:			
Calculated:	C, 50.2;	H, 4.39;	N, 10.2
Found:	C, 50.1;	H, 4.55;	N, 9.96

[0253] **Example 69:** R=2-methoxyethoxycarbonyl, R⁶=3,5-difluorophenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.33, methanol:dichloromethane (5:95); 300 MHz NMR: 8.89 (d,1), 8.42 (s,1), 7.49 (t,1), 7.23 (d,2), 4.68 (m,1), 4.52 (dd,2), 4.23 (t,2), 3.56 (t,2), 3.29 (s,3), 2.16 (m,1), 0.88 (d,3), 0.83 (d,3).

Analysis for C ₂₂ H ₂₃ F ₅ N ₄ O ₆ :			
Calculated:	C, 49.4;	H, 4.34;	N, 10.5
Found:	C, 49.1;	H, 4.29;	N, 10.4

[0254] **Example 70:** R=2-methoxyethoxycarbonyl, R⁶=phenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.26, methanol:dichloromethane (5:95); 300 MHz NMR: 8.83 (d,1), 8.77 (s,1), 8.42 (s,1), 7.50 (m,5), 4.67 (t,1), 4.55 (dd,2), 4.23 (t,2), 3.57 (t,2), 3.29 (s,3), 2.16 (m,1), 0.89 (d,3), 0.83 (d,3).

Analysis for C ₂₂ H ₂₅ F ₃ N ₄ O ₆ :			
Calculated:	C, 53.0;	H, 5.06;	N, 11.2
Found:	C, 53.0;	H, 5.02;	N, 11.2

[0255] **Example 71:** R=2-methoxyethoxycarbonyl, R⁶=4-methoxyphenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.20, methanol:dichloromethane (5:95); 300 MHz NMR: 8.87 (d,1), 8.74 (s,1), 8.40 (s,1), 7.44 (d,2), 7.02 (d,2), 4.68 (t,1), 4.60 (dd,2), 4.24 (t,2), 3.83 (s,3), 3.58 (t,2), 3.30 (s,3), 2.18 (m,1), 0.91 (d,3), 0.86 (d,3).

Analysis for C ₂₃ H ₂₇ F ₃ N ₄ O ₇ :			
Calculated:	C, 52.3;	H, 5.15;	N, 10.6
Found:	C, 52.0;	H, 5.13;	N, 10.5

[0256] Example 72: R=isopropoxycarbonyl, R⁶=3,5-difluorophenyl; white solid; chromatography solvent: methanol:dichloromethane (2.5:97.5); TLC: R_f=0.31, methanol:dichloromethane (5:95); NMR: 8.89 (d,1), 8.60 (s,1), 8.43 (s,1), 7.49 (t,1), 7.22 (d,2), 4.90 (m,1), 4.64 (m,3), 2.17 (m,1), 1.25 (d,6), 0.87 (d,3), 0.82 (d,3).

Analysis for C ₂₂ H ₂₃ F ₅ N ₄ O ₅ :			
Calculated:	C, 51.0;	H, 4.47;	N, 10.8
Found:	C, 50.7;	H, 4.44;	N, 10.7

[0257] Example 73: R=2-methoxyethoxycarbonyl, R⁶=2-thienyl; yellow solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f=0.37, methanol:dichloromethane (5:95); 300 MHz NMR: 9.04 (d,1), 8.80 (s,1), 8.39 (s,1), 7.85 (d,1), 7.33 (d,1), 7.16 (t,1), 4.88 (dd,2), 4.74 (t,1), 4.22 (t,2), 3.56 (t,2), 3.28 (s,3), 2.21 (m,1), 0.94 (d,3), 0.90 (d,3).

Analysis for C ₂₀ H ₂₃ F ₃ N ₄ O ₆ S:			
Calculated:	C, 47.6;	H, 4.60;	N, 11.1
Found:	C, 47.7;	H, 4.61;	N, 10.9

[0258] Example 74: R=2-methoxyethoxycarbonyl, R⁶=4-fluorophenyl; white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f=0.34, methanol:dichloromethane (5:95); 300 MHz NMR: 8.86 (d,1), 8.81 (s,1), 8.41 (s,1), 7.54 (m,2), 7.33 (t,2), 4.60 (m,3), 4.24 (m,2), 3.57 (m,2), 3.29 (s,3), 2.16 (m,1), 0.89 (d,3), 0.83 (d,3).

Analysis for C ₂₂ H ₂₄ F ₃ N ₄ O ₆ :			
Calculated:	C, 51.2;	H, 4.68;	N, 10.9
Found:	C, 51.1;	H, 4.67;	N, 10.9

EXAMPLE 75

2-[5-Trifluoroacetyl-amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0259] To a cooled (-5 °C) solution of the product of Example 15 (0.104 g) in dichloromethane was added trifluoroacetic anhydride (50 µL). The reaction mixture was allowed to warm to room temperature. After 1 h a further charge of trifluoroacetic anhydride (25 µL) was made. After stirring for 1 h, the mixture was evaporated and the residue was purified by chromatography, eluting with dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5) to afford the title compound; TLC: R_f=0.44, dichloromethane:methanol (98:2); NMR: 0.7-1.0 (m,6), 2.03-2.30 (m,1), 3.97-4.80 (m,3), 7.23-7.40 (m,2), 7.53-7.70 (m,2), 8.43 (s,1), 8.87 (d,1), 10.96 (s,1); MS: m/z=511(M+1).

Analysis for $C_{20}H_{17}F_7N_4O_4 \cdot 0.5 H_2O$:			
Calculated:	C, 46.25;	H, 3.49;	N, 10.78
Found:	C, 46.22;	H, 3.57;	N, 10.62

10 **EXAMPLES 76-79**

[0260] Using procedures similar to that described in Example 75, the following compounds of formula I wherein R^0 is isopropyl, R is trifluoroacetyl and R^6 is the indicated group were prepared from the corresponding compounds of formula I wherein R is hydrogen, with exceptions as noted.

15 **[0261] Example 76:** R^6 =phenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5); TLC: R_f =0.40, dichloromethane:methanol (98:2); NMR: 0.7-0.97 (m,6), 2.07-2.33 (m,1), 4.00-4.77 (m,3), 7.40-7.67 (m,5), 8.45 (s,1), 8.86 (d,1), 10.96 (d,1); MS: m/z =493(M+1).

Analysis for $C_{20}H_{18}F_6N_4O_4 \cdot 0.6 H_2O$:			
Calculated:	C, 47.74;	H, 3.84;	N, 11.13
Found:	C, 47.69;	H, 3.83;	N, 10.91

25 **[0262] Example 77:** R^6 =4-methoxyphenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5); TLC: R_f =0.52, dichloromethane:methanol (98:2); NMR: 0.70-0.97 (m,6), 2.03-2.27 (m,1), 3.81 (s,3), 4.00-4.77 (m,3), 6.97-7.07 (m,2), 7.47-7.57 (m,2), 8.40 (s,1), 8.88 (d,1), 10.92 (d,1); MS: m/z =523(M+1).

Analysis for $C_{21}H_{20}F_6N_4O_5 \cdot 0.6 H_2O$:			
Calculated:	C, 47.30;	H, 4.00;	N, 10.50
Found:	C, 47.27;	H, 4.00;	N, 10.32

35 **[0263] Example 78:** R^6 =2-thienyl: Omitting the second addition of trifluoroacetic anhydride and stirring overnight: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.2:0.8); TLC: R_f =0.44, dichloromethane:methanol (98:2); NMR: 0.83-1.0 (m,6), 2.20-2.30 (m,1), 4.73-5.03 (m,3), 7.10-8.47 (m,4), 9.06 (d,1), 10.90 (d,1); MS: m/z =499(M+1).

Analysis for $C_{18}H_{16}F_6N_4O_4S$:			
Calculated:	C, 43.37;	H, 3.23;	N, 11.24
Found:	C, 43.20;	H, 3.57;	N, 11.12

45 **[0264] Example 79:** R^6 =3,5-difluorophenyl: Omitting the second addition of trifluoroacetic anhydride and stirring overnight: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99:1); TLC: R_f =0.46, dichloromethane:methanol (98:2); NMR: 0.83-0.97 (m,6), 2.10-2.30 (m,1), 4.43-4.80 (m,3), 7.27-7.37 (m,2), 7.47-7.60 (m,1), 8.44 (s,1), 8.90 (d,1), 11.03 (d,1); MS: m/z =529(M+1).

Analysis for $C_{20}H_{16}F_8N_4O_4$:			
Calculated:	C, 45.46;	H, 3.05;	N, 10.60
Found:	C, 45.51;	H, 3.25;	N, 10.44

10 **EXAMPLE 80**

2-[5-Ethylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

15 [0265] To a solution of the product of Example 15 (0.40 g) in dimethylformamide (6 mL) was added 2,6-lutidine (0.14 mL) followed by ethyl iodide (0.12 mL). The reaction mixture was stirred at room temperature for 5 days over which time further aliquots (4) of 2,6-lutidine (0.14 mL) and ethyl iodide (0.12 mL) were added. Ethyl acetate was added and the reaction mixture was washed (H_2O , saturated aqueous copper sulfate, brine), dried ($MgSO_4$) and evaporated. The residue was purified by chromatography, with dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5) as the eluent, to give the title compound; NMR: 0.83-0.91 (dd,6), 1.19 (t,3), 2.07-2.27 (m,1), 4.43-4.73 (m,3), 5.45 (t,1), 7.12 (s,1), 7.20-7.37 (m,2), 7.43-7.63 (m,2), 8.82 (d,1); MS: $m/z=443(M+1)$.

Analysis for $C_{20}H_{22}F_4N_4O_3$:			
Calculated:	C, 54.29;	H, 5.01;	N, 12.66
Found:	C, 54.36;	H, 5.18;	N, 12.50

30 **EXAMPLES 81-83**

[0266] Using procedures similar to that described in Example 80, the following compounds of formula I wherein R^0 is isopropyl, R is ethyl and R^6 is the indicated group were prepared from the corresponding compounds of formula I wherein R is hydrogen.

35 [0267] **Example 81:** R^6 =phenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.3:0.7); TLC: R_f =0.37, dichloromethane:methanol (98:2); NMR: 0.73-1.00 (m,6), 1.17 (t,3), 2.07-2.30 (m,1), 3.03-3.23 (m,2), 4.40-4.70 (m,3), 5.40 (t,1), 7.12 (s,1), 7.33-7.57 (m,5), 8.79 (d,1); MS: $m/z=425(M+1)$.

Analysis for $C_{20}H_{23}F_3N_4O_3$:			
Calculated:	C, 56.59;	H, 5.46;	N, 13.20
Found:	C, 56.69;	H, 5.53;	N, 13.08

40 [0268] **Example 82:** R^6 =2-thienyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.3:0.7); TLC: R_f =0.37, dichloromethane:methanol (98:2); NMR: 0.83-1.07 (m,6), 1.10-2.23 (t,3), 2.13-3.13 (m,1), 3.03-3.17 (m,2), 4.67-4.87 (m,3), 5.55 (t,1), 7.03-7.20 (m,3), 7.69 (d,1), 8.97 (d,1); MS: $m/z=431(M+1)$.

Analysis for $C_{18}H_{21}F_3N_4O_3S$:			
Calculated:	C, 50.22;	H, 4.91;	N, 13.01
Found:	C, 50.13;	H, 5.00;	N, 12.93

[0269] **Example 83:** R⁶=3,5-difluorophenyl: Chromatography solvent: dichloromethane:methane (two columns, gradient, 99.5:0.5, 99.3:0.7); TLC: R_f=0.47, dichloromethane:methanol (98:2); NMR: 0.80-1.0 (m,6), 1.17 (t,3), 2.10-2.30 (m,1), 3.03-3.20 (m,2), 4.47-4.73 (m,3), 5.57 (t,1), 7.07-7.23 (m,3), 7.33-7.50 (m,1), 8.86 (d,1); MS: m/z=461(M+1).

Analysis for C ₂₀ H ₂₁ F ₅ N ₄ O ₃ :			
Calculated:	C, 52.17;	H, 4.59;	N, 12.16
Found:	C, 52.00;	H, 4.84;	N, 11.80

EXAMPLE 84

2-[2-(4-Fluorophenyl)-5-methylamino-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0270] To a solution of 2-[2-(4-fluorophenyl)-5-(N-trifluoroacetyl-N-methylamino)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (0.58 g) in tetrahydrofuran (5 mL) was added water (10 mL) followed by potassium carbonate (0.76 g). The mixture was stirred overnight, extracted into dichloromethane, washed (water, brine), dried (MgSO₄) and evaporated. The residue was purified by chromatography, with dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5) as the eluent, to give the title compound; TLC: R_f=0.34, dichloromethane:methanol (98:2); NMR: 0.70-1.0 (m,6), 2.03-2.33 (m,1), 2.71 (d,3), 4.00-4.70 (m,3), 5.63-5.73 (m,1), 7.03 (s,1), 7.17-7.30 (m,2), 7.43-7.60 (m,2), 8.81 (d,1); MS: m/z=429(M+1).

Analysis for C ₁₉ H ₂₀ F ₄ N ₄ O ₃ · 0.4 H ₂ O:			
Calculated:	C, 52.39;	H, 4.81;	N, 12.86
Found:	C, 52.38;	H, 4.76;	N, 12.76

[0271] The intermediate N-trifluoroacetyl-N-methylamino compound was prepared as follows.

[0272] To a solution of the product of Example 75 (0.57 g) in dimethylformamide was added Na₂CO₃ (0.32 g), followed by methyl iodide (0.52 mL). The mixture was stirred overnight. Dichloromethane was added and the mixture was filtered. The filtrates were washed (H₂O, brine), dried (MgSO₄), and evaporated. The residue was purified by chromatography, with dichloromethane:acetone (gradient, 95:5, 90:10) as the eluent, to give the N-trifluoroacetyl-N-methylamino compound; TLC: R_f=0.33, dichloromethane:acetone (90:10); MS: m/z=525(M+1).

EXAMPLES 85-88

[0273] Using procedures similar to that described in Example 84, the following compounds of formula I wherein R⁰ is isopropyl, R is methyl and R⁶ is the indicated group were prepared from the corresponding 2-[2-aryl-5-(N-trifluoroacetyl-N-methylamino)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamides wherein R⁶ is the indicated aryl group.

[0274] **Example 85:** R⁶=phenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5); TLC: R_f=0.36, dichloromethane:methanol (98:2); NMR: 0.73-1.0 (m,6), 2.07-2.33 (m,1), 2.73 (d,3), 4.07-4.73 (m,3), 5.63-5.73 (m,1), 7.05 (s,1), 7.37-7.53 (m,5), 8.81 (d,1); MS: m/z=411(M+1).

Analysis for C ₁₉ H ₂₁ F ₃ N ₄ O ₃ · 0.3 H ₂ O:			
Calculated:	C, 54.88;	H, 5.23;	N, 13.47
Found:	C, 54.93;	H, 5.20;	N, 13.48

[0275] Example 86: $R^6=4$ -methoxyphenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5); TLC: $R_f=0.41$, dichloromethane:methanol (98:2); NMR: 0.80-1.00 (m,6), 2.07-2.27 (m,1), 2.71 (d,3), 3.79 (s,3), 4.40-4.70 (m,3), 5.58 (d,1), 6.90-7.00 (m,2), 7.03 (s,1), 7.27-7.40 (m,2), 8.79 (d,1); MS: $m/z=441(M+1)$.

Analysis for $C_{20}H_{23}F_3N_4O_4$:			
Calculated:	C, 54.54;	H, 5.26;	N, 12.72
Found:	C, 54.16;	H, 5.42;	N, 12.49

[0276] Example 87: $R^6=2$ -thienyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.3:0.7); TLC: $R_f=0.30$, dichloromethane:methanol (98:2); NMR: 0.73-1.0 (m,6), 2.17-2.33 (m,1), 2.71 (s,3), 4.07-4.33 (m,1), 4.67-4.97 (m,2), 5.70-5.87 (m,1), 7.07-7.30 (m,3), 7.63-7.73 (m,1), 8.06 (d,1); MS: $m/z=417(M+1)$.

Analysis for $C_{17}H_{19}F_3N_4O_3S \cdot 0.6 CH_3OH \cdot 0.5 H_2O$:			
Calculated:	C, 47.54;	H, 5.07;	N, 12.60
Found:	C, 48.06;	H, 5.04;	N, 12.23

[0277] Example 88: $R^6=3,5$ -difluorophenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.3:0.7); TLC: $R_f=0.30$, dichloromethane:methanol (98:2); NMR: 0.70-0.97 (m,6), 2.07-2.27 (m,1), 2.71 (d,3), 4.37-4.73 (m,3), 5.77-5.90 (m,1), 7.03 (s,1), 7.10-7.23 (m,2), 7.33-7.47 (m,1), 8.85 (d,1); MS: $m/z=447(M+1)$.

Analysis for $C_{19}H_{19}F_5N_4O_3$:			
Calculated:	C, 51.12;	H, 4.29;	N, 12.55
Found:	C, 51.06;	H, 4.37;	N, 12.38

EXAMPLES 85.a.-88.a.

[0278] Using procedures similar to that described in Example 84.a., the following 2-[2-aryl-5-(N-trifluoroacetyl)-N-methylamino)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamides wherein R^6 is the indicated aryl group were prepared from the corresponding compounds of formula I wherein R is trifluoroacetyl, which were prepared in Examples 76-79.

[0279] Example 85.a.: $R^6=phenyl$: Chromatography solvent: dichloromethane:acetone (gradient, 95.0:5.0, 90.0:10.0); TLC: $R_f=0.30$, dichloromethane:acetone (90:10); MS: $m/z=507(M+1)$.

[0280] Example 86.a.: $R^6=4$ -methoxyphenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 98.5:1.5); TLC: $R_f=0.33$, dichloromethane:methanol (98:2); MS: $m/z=537(M+1)$.

[0281] Example 87.a.: $R^6=2$ -thienyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.0:1.0); TLC: $R_f=0.37$, dichloromethane:methanol (98:2); MS: $m/z=513(M+1)$.

[0282] Example 88.a.: $R^6=3,5$ -difluorophenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.3:0.7); TLC: $R_f=0.39$, dichloromethane:methanol (98:2); MS: $m/z=543(M+1)$.

EXAMPLE 89

2-[2-(3,5-Difluorophenyl)-5-formamido-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0283] The title compound was prepared using the method described in Chen and Benoiton, *Synthesis* (1979), 709.

[0284] To a suspension of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.193 g) in dichloromethane (8 mL) was added 90% formic acid (1 mL). This mixture was stirred at room temperature for 10 min. This mixture was then added N-methylmorpholine (0.10 mL) followed by the product from Example 27 (0.40 g). After 3 days, the reaction was diluted with dichloromethane, washed (brine), dried (MgSO₄) and evaporated. The residue was purified by chromatography, with dichloromethane:methanol (gradient, 99.5:0.5, 98:2) as the eluent, to give the title compound; TLC: R_f=0.53, dichloromethane:acetone (70:30); NMR: 0.67-1.00 (m,6), 2.07-2.30 (m,1), 4.33-4.83 (m,3), 7.17-7.30 (m,2), 7.43-7.57 (m,1), 8.38 (s,1), 8.83-8.97 (m,2), 10.12 (s,1); MS: m/z=461(M+1).

Analysis for C₁₉H₁₇F₅N₄O₄ • 0.4 H₂O:

Calculated:	C, 48.80;	H, 3.83;	N, 11.98
Found:	C, 48.81;	H, 3.94;	N, 11.77

EXAMPLE 90

2-[2-(4-Fluorophenyl)-6-oxo-5-(2,2,2-trifluoroethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0285] To a solution of the compound from Example 15 (414 mg) in tetrahydrofuran (8 mL) and dichloromethane (8 mL) at 0 °C was added pyridine (320 mg) followed by 2,2,2-trifluoroethyl chloroformate (300 µL). After 1 h the reaction mixture was diluted with diethyl ether and quenched with ice. The phases were separated and the organic phase was washed (dilute hydrochloric acid, brine), dried, and evaporated to afford a gummy solid. This solid was triturated with diethyl ether:hexanes (10 mL, 1:1) to afford a white powder, which was collected by filtration and dried under vacuum to yield the title compound (460 mg); R_f=0.48, chloroform:methanol (20:1); NMR DMSO/D₂O: 8.40 (s,1), 7.56 (m,2), 7.28 (m,2), 4.83-4.56 (m,4), 2.22 (m,1), 0.80 (dd,6); MS: m/z=541(M+1).

Analysis for C₂₁H₁₉F₇N₄O₅ • 0.5 H₂O:

Calculated:	C, 45.91;	H, 3.67;	N, 10.20
Found:	C, 46.05;	H, 3.52;	N, 10.25

[0286] The intermediate 2,2,2-trifluoroethyl chloroformate was prepared using a procedure similar to that described in U.S. Patent Number 3,852,464, except that bis(trichloromethyl) carbonate was used in place of phosgene.

EXAMPLES 91-108

[0287] Using procedures similar to that described in Example 90 and using the required acyl, sulfonyl, or aminosulfonyl chloride, the following compounds of formula I wherein R⁰ is isopropyl and R and R⁶ have the indicated values were prepared from the corresponding compounds of formula I wherein R is hydrogen.

[0288] **Example 91:** R=cyclohexylaminosulfonyl, R⁶=4-fluorophenyl: Purified by trituration with methyl tert-butyl ether:diethyl ether; TLC: R_f=0.33, tetrahydrofuran:dichloromethane (1:9); NMR: 8.92 (s,1), 8.82 (d,1), 7.96 (s,1), 7.64 (d,1), 7.51 (m,2), 7.31 (t,2), 4.61 (m,3), 1.78-1.18 (m,10), 0.88 (d,3), 0.82 (d,3); MS: m/z=576(M+1).

Analysis for C₂₄H₂₉F₄N₅O₅S:

Calculated:	C, 50.08;	H, 5.08;	N, 12.17
Found:	C, 50.11;	H, 5.20;	N, 11.90

[0289] **Example 92:** R=benzylaminosulfonyl, R⁶=4-fluorophenyl: Chromatography solvent: dichloromethane:meth-

EP 0 528 633 B1

anol (30:1); TLC: R_f =0.29, dichloromethane:methanol (20:1); NMR: 9.15 (s,1), 8.85 (d,1), 8.2 (m,1), 7.95 (s,1), 7.53 (m,2), 7.30 (m,7), 4.67 (t,1), 4.13 (d,2), 2.2 (m,1), 0.83 (dd,6); MS: m/z =584(M+1).

5

Analysis for $C_{25}H_{25}F_4N_5O_5S$:			
Calculated:	C, 51.46;	H, 4.32;	N, 12.00
Found:	C, 51.35;	H, 4.49;	N, 11.86

10

[0290] **Example 93:** R=cyclohexylaminosulfonyl, R^6 =phenyl: Purified by trituration with diethyl ether; TLC: R_f =0.31, dichloromethane:methanol (20:1); NMR (DMSO/ D_2O): 8.0 (s,1), 7.5 (mm,5), 4.60 (dd,2), 4.0 (m,1), 3.1 (m,1), 2.2 (m,1), 1.7-1.4 (mm,4), 1.3-1.1 (mm,6), 0.82 (qd,6); MS: m/z =558(M+1).

15

Analysis for $C_{24}H_{30}F_3N_5O_5S \cdot 0.5 H_2O$:			
Calculated:	C, 50.87;	H, 5.51;	N, 12.36
Found:	C, 50.92;	H, 5.46;	N, 12.18

20

[0291] **Example 94:** R=benzylaminosulfonyl, R^6 =phenyl: Chromatography solvent:dichloromethane:methanol (30:1); TLC: R_f =0.28, dichloromethane:methanol (20:1); NMR: 9.10 (s,1), 8.84 (d,1), 8.15 (m,1), 7.95 (s,1), 7.47 (m,5), 7.29 (m,5), 4.68 (dd,1), 4.53 (2d,2), 4.12 (d,2), 2.17 (m,1), 0.89 (dd,6); MS: m/z =566(M+1).

25

30

Analysis for $C_{25}H_{26}F_3N_5O_5S$:			
Calculated:	C, 52.26;	H, 4.74;	N, 12.19
Found:	C, 52.15;	H, 4.69;	N, 12.10

35

[0292] **Example 95:** R=benzylsulfonyl, R^6 =4-fluorophenyl: Chromatography solvent: dichloromethane:methanol (99:1); TLC: R_f =0.44, dichloromethane:methanol (20:1); NMR (DMSO/ D_2O): 7.72 (s,1), 7.55-7.26 (mm,9), 4.6 (s,2), 4.58 (dd,2), 2.24 (m,1), 0.86 (d,3), 0.76 (d,3); MS: m/z =569(M+1).

40

Analysis for $C_{25}H_{24}F_4N_4O_5S$:			
Calculated:	C, 52.81;	H, 4.25;	N, 9.85
Found:	C, 53.01;	H, 4.36;	N, 9.69

45

[0293] **Example 96:** R=benzylsulfonyl, R^6 =phenyl: Chromatography solvent: dichloromethane:acetonitrile (9:1), followed by trituration with diethyl ether; TLC: R_f =0.44, dichloromethane:methanol (95:5); NMR: 9.33 (s,1), 8.86 (d,1), 7.77 (s,1), 7.58-7.33 (m,10), 4.69 (m,1), 4.60 (s,2), 2.50 (m,1), 0.90 (d,3), 0.88 (d,3); MS: m/z =551(M+1).

50

Analysis for $C_{25}H_{25}F_3N_4O_5S$:			
Calculated:	C, 54.54;	H, 4.58;	N, 10.18
Found:	C, 54.80;	H, 4.53;	N, 10.09

55

[0294] **Example 97:** R=isopropylaminosulfonyl, R⁶=phenyl: Chromatography solvent: dichloromethane:methanol (98:2), followed by trituration with diethyl ether; TLC: R_f=0.28, dichloromethane:methanol (20:1); NMR: 8.96 (s,1), 8.2 (d,1), 7.96 (s,1), 7.51 (m,5), 5.55 (dd,2), 3.42 (m,1), 2.34 (m,1), 1.05 (d,6), 0.88 (d,3), 0.82 (d,3); MS: m/z=518(M+1).

Analysis for C ₂₁ H ₂₆ F ₃ N ₅ O ₅ S:			
Calculated:	C, 48.74;	H, 5.06;	N, 13.53
Found:	C, 48.53;	H, 5.11;	N, 13.45

[0295] **Example 98:** R=methoxycarbonyl, R⁶=3,5-difluorophenyl: Chromatography solvent: dichloromethane:methanol (96:4); TLC: R_f=0.47, dichloromethane:methanol (95:5); NMR: 8.46 (s,1), 7.43 (t,1), 7.22 (d,2), 4.71 (d,1), 4.45 (d,1), 4.02 (d,1), 3.70 (s,3), 2.24 (m,1), 0.85 (d,3), 0.75 (d,3); MS: m/z=491(M+1).

Analysis for C ₂₀ H ₁₉ F ₅ N ₄ O ₅ :			
Calculated:	C, 48.98;	H, 3.90;	N, 11.42
Found:	C, 49.22;	H, 3.93;	N, 11.52

[0296] **Example 99:** R=2,2,2-trifluoroethoxycarbonyl, R⁶=phenyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.42, chloroform:methanol (20:1); NMR (DMSO/D₂O): 8.41 (s,1), 7.5 (m,5), 4.79 (m,2), 4.67 (dd,2), 2.24 (m,1), 0.82 (dd,6); MS: m/z=523(M+1).

Analysis for C ₂₁ H ₂₀ F ₆ N ₄ O ₅ • 0.7 H ₂ O:			
Calculated:	C, 47.14;	H, 4.03;	N, 10.47
Found:	C, 47.03;	H, 4.07;	N, 10.32

[0297] **Example 100:** R=methylthiocarbonyl, R⁶=phenyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.38, dichloromethane:methanol (20:1); NMR: 9.98 (s,1), 8.85 (d,1), 8.60 (s,1), 7.50 (m,5), 4.68 (t,1), 4.55 (dd,2), 2.30 (s,3), 2.18 (m,1), 0.87 (dd,6); MS: m/z=471(M+1).

Analysis for C ₂₀ H ₂₁ F ₃ N ₄ O ₄ S:			
Calculated:	C, 51.06;	H, 4.50;	N, 11.91
Found:	C, 51.04;	H, 4.64;	N, 11.59

[0298] **Example 101:** R=ethylthiocarbonyl, R⁶=phenyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.41, dichloromethane:methanol (20:1); NMR: 9.88 (d,1), 8.86 (d,1), 8.59 (s,1), 7.49 (m,5), 4.69 (t,1), 4.55 (m,2), 2.85 (q, 2), 2.17 (m,1), 1.25 (t,3), 0.88 (dd,6); MS: m/z=485(M+1).

Analysis for C ₂₁ H ₂₃ F ₃ N ₄ O ₄ S • 0.35 H ₂ O:			
Calculated:	C, 51.39;	H, 4.87;	N, 11.42

(continued)

Analysis for $C_{21}H_{23}F_3N_4O_4S \cdot 0.35 H_2O$:			
Found:	C, 51.36;	H, 4.93;	N, 11.34

[0299] **Example 102:** R=methylthiocarbonyl, R⁶=4-fluorophenyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.44, dichloromethane:methanol (20:1); NMR: 10.0 (s,1), 8.85 (d,1), 8.58 (s,1), 7.55 (dd,), 7.35 (dd,2), 4.66 (t,1), 4.59 (dd,2), 2.29 (s,3), 2.18 (m,1), 0.88 (dd,6); MS: m/z=489(M+1).

Analysis for $C_{20}H_{20}F_4N_4O_4S$:			
Calculated:	C, 49.18;	H, 4.13;	N, 11.47
Found:	C, 49.23;	H, 4.19;	N, 11.47

[0300] **Example 103:** R=ethylthiocarbonyl, R⁶=4-fluorophenyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.45, dichloromethane:methanol (20:1); NMR: 9.90 (s,1), 8.85 (d,1), 8.55 (s,1), 7.55 (m,2), 7.32 (m,2), 4.67 (t,1), 4.59 (m,2), 2.85 (q,2), 2.15 (m,1), 1.25 (t,3), 0.86 (dd,6); MS: m/z=503(M+1).

Analysis for $C_{21}H_{22}F_4N_4O_4S \cdot 0.4 H_2O$:			
Calculated:	C, 49.49;	H, 4.51;	N, 10.99
Found:	C, 49.54;	H, 4.53;	N, 11.07

[0301] **Example 104:** R=methylthiocarbonyl, R⁶=2-thienyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.48, dichloromethane:methanol (20:1); NMR: 9.98 (s,1), 9.05 (d,1), 8.55 (s,1), 7.84 (d,1), 7.48 (d,1), 7.25 (t,1), 4.89 (t,1), 4.8 (dd,2), 2.29 (s,3), 2.22 (m,1), 0.95 (dd,6); MS: m/z=477(M+1).

Analysis for $C_{18}H_{19}F_3N_4O_4S_2 \cdot 0.75 H_2O$:			
Calculated:	C, 44.12;	H, 4.22;	N, 11.43
Found:	C, 44.13;	H, 4.24;	N, 11.27

[0302] **Example 105:** R=ethylthiocarbonyl, R⁶=2-thienyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.47, dichloromethane:methanol (20:1); NMR: 9.90 (s,1), 9.07 (d,1), 8.55 (s,1), 7.85 (d,1), 7.48 (d,1), 7.17 (t,1), 4.89 (t,1), 4.84 (dd,2), 1.25 (t,3), 2.87 (q, 2), 2.24 (m,1), 0.94 (dd,6); MS: m/z=491(M+1).

Analysis for $C_{18}H_{19}F_3N_4O_4S_2 \cdot 0.75 H_2O$:			
Calculated:	C, 45.28;	H, 4.50;	N, 11.12
Found:	C, 45.44;	H, 4.45;	N, 11.02

[0303] **Example 106:** R=2,2,2-trifluoroethoxycarbonyl, R⁶=2-thienyl: Purified by trituration with diethyl ether:hexanes (1:5); TLC: R_f=0.49, dichloromethane:methanol (20:1); NMR: 9.47 (s,1), 9.05 (d,1), 8.35 (s,1), 7.87 (d,1), 7.40 (d,1), 7.16 (t,1), 4.98-4.73 (mm,3), 2.26 (m,1), 0.96 (dd,6); MS: m/z=529(M+1).

Analysis for $C_{19}H_{18}F_8N_4O_5S \cdot 0.5 H_2O$:			
Calculated:	C, 42.46;	H, 3.56;	N, 10.42
Found:	C, 42.52;	H, 3.71;	N, 10.39

10 [0304] Example 107: R=tert-butylaminosulfonyl, R⁶=4-fluorophenyl: Chromatography solvent: dichloromethane:methanol (95:5); TLC: R_f=0.17, dichloromethane:methanol (95:5); NMR (DMSO/D₂O): 7.97 (s,1), 7.55 (m,2), 7.28 (t,2), 4.68 (d,2), 4.47 (d,1), 4.02 (d,1), 2.21 (m,1), 1.23 (s,9), 0.84 (d,3), 0.74 (d,3); MS: m/z=550(M+1).

Analysis for $C_{22}H_{27}F_4N_5O_5S \cdot 0.5 H_2O$:			
Calculated:	C, 47.3;	H, 5.05;	N, 12.5
Found:	C, 47.4;	H, 5.09;	N, 12.4

20 [0305] Example 108: R=tert-butylaminosulfonyl, R⁶=phenyl: Chromatography solvent: dichloromethane:methanol (95:5); TLC: R_f=0.21, dichloromethane:methanol (95:5); NMR (DMSO/D₂O): 7.94 (s,1), 7.44 (s,5), 4.50 (dd,2), 4.00 (m,1), 2.50 (m,1), 1.19 (s,9), 0.80 (d,3), 0.73 (d,3); MS: m/z=532(M+1).

Analysis for $C_{22}H_{28}F_3N_5O_5S \cdot 0.25 H_2O$:			
Calculated:	C, 49.3;	H, 5.36;	N, 13.1
Found:	C, 49.3;	H, 5.44;	N, 12.9

EXAMPLE 109

25 2-[5-Dimethylaminooxycarbonylamino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxo-propyl)acetamide.

40 [0306] A slurry of the compound from Example 15 (0.414 g) in dry dichloromethane (10 mL) was cooled in an ice/salt bath to an internal temperature of less than 1 °C. Bis(trichloromethyl) carbonate (0.1994 g) as a solution in dry dichloromethane (5 mL) was added dropwise at such a rate to maintain the internal temperature below 2 °C. After the addition was complete, the reaction was stirred an additional 10 min. Triethylamine (1.18 mL) as a solution in dry dichloromethane was added dropwise, maintaining an internal temperature of less than 4 °C; and, after the addition was complete, the reaction was stirred an additional 20 min. N,N-Dimethylhydroxylamine hydrochloride (0.1883 g) was added
 45 as a solid to the reaction mixture and it was stirred for 30 min. The reaction mixture was diluted with ethyl acetate, washed (saturated ammonium chloride, brine), dried and evaporated to yield a solid. This solid was triturated with diethyl ether to give the title compound (0.3338 g) as a white solid; TLC: R_f=0.4, dichloromethane:methanol (20:1); NMR: 8.55 (s,1), 7.54 (m,2), 7.29 (m,2), 4.59 (dd,2), 4.02 (dd,1), 2.82 (s,6), 2.22 (m,1), 0.84 (d,3), 0.74 (d,3); MS: m/z=502(M+1).

Analysis for $C_{21}H_{23}F_4N_5O_5$:			
Calculated:	C, 50.30;	H, 4.62;	N, 13.97
Found:	C, 50.32;	H, 4.68;	N, 13.79

EXAMPLES 110-112

[0307] The following compounds of formula I wherein R^0 is isopropyl and R and R^6 have the indicated values were prepared using procedures similar to that described in Example 109 by replacing 2-[5-amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxo-propyl)acetamide with the compound of formula I wherein R is hydrogen and R^6 is the indicated group and by replacing N,N-dimethylhydroxylamine hydrochloride with the required nucleophile.

[0308] Example 110: R=methylaminocarbonyl, R^6 =4-fluorophenyl: Using methylamine hydrochloride; chromatography solvent: dichloromethane:methanol (20:1), followed by washing an ethyl acetate solution of the appropriate fractions with 0.1 N HCl, drying and evaporation to give a solid; TLC: R_f =0.19, dichloromethane:methanol (20:1); NMR: 8.65 (s,1), 7.53 (m,2), 7.26 (t,2), 4.64 (d,1), 4.45 (d,1), 4.04 (d,1), 2.65 (s,3), 2.22 (m,1), 0.86 (d,3), 0.76 (d,3); MS: m/z =472(M+1).

Analysis for $C_{20}H_{21}F_4N_5O_4 \cdot 0.33 H_2O$:			
Calculated:	C, 50.32;	H, 4.57;	N, 14.67
Found:	C, 50.42;	H, 4.54;	N, 14.77

[0309] Example 111: R=isopropylaminocarbonyl, R^6 =phenyl: Using isopropylamine; chromatography solvent: dichloromethane:methanol (97:3); TLC: R_f =0.28, dichloromethane:methanol (95:5); NMR (DMSO- D_6): 8.64 (s,1), 7.46 (m,5), 4.61 (d,1), 4.45 (d,1), 4.08 (d,1), 3.72 (m,1), 2.24 (m,1), 1.09 (d,6), 0.87 (d,3), 0.79 (d,3); MS: m/z =482(M+1).

Analysis for $C_{22}H_{26}F_3N_5O_4$:			
Calculated:	C, 54.88;	H, 5.44;	N, 14.54
Found:	C, 54.60;	H, 5.53;	N, 14.47

[0310] Example 112: R=methylaminocarbonyl, R^6 =phenyl: Using methylamine hydrochloride; chromatography solvent: dichloromethane:methanol (96:4), followed by crystallization from dichloromethane; TLC: R_f =0.21, dichloromethane:methanol (95:5); NMR (DMSO- D_6): 8.66 (s,1), 7.47 (m,5), 4.61 (d,1), 4.45 (d,1), 4.08 (d,1), 2.66 (s,3), 2.25 (m,1), 0.86 (d,3), 0.79 (d,3); MS: m/z =454(M+1).

Analysis for $C_{20}H_{22}F_3N_5O_4 \cdot 0.5 H_2O$:			
Calculated:	C, 51.95;	H, 5.01;	N, 15.14
Found:	C, 51.86;	H, 4.98;	N, 15.09

EXAMPLE 113

2-[2-(4-Fluorophenyl)-5-isopropylaminocarbonyl-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxo-propyl)acetamide.

[0311] To a slurry of 2-[5-amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)-acetamide (0.414 g) in dry tetrahydrofuran (3 mL) and dichloromethane (5 mL) was added isopropyl isocyanate (110 μ L). After overnight stirring, dimethylformamide (1.5 mL), triethylamine (100 μ L) and isopropyl isocyanate (50 μ L) were added. The reaction mixture was heated to reflux for 24 h. The mixture was cooled to room temperature and cuprous chloride (120 mg) followed by three portions of isopropyl isocyanate (50 μ L, 50 μ L, and 25 μ L) were added. The reaction mixture was diluted with diethyl ether and rinsed with dilute hydrochloric acid and brine. The organic phase

was dried and evaporated to afford an oil which was purified by chromatography, with chloroform:methanol (gradient 100:0, 97:3) as the eluent. This afforded 160 mg of a yellow-green solid. Analysis by NMR showed line broadening, indicative of contamination by a heavy metal. The remaining 135 mg sample was dissolved in methyl *tert*-butyl ether and rinsed with 5% disodium EDTA solution followed by dilute aqueous ammonia. The organic phase was dried and evaporated to afford the title compound (120 mg) as an off-white powder; TLC: R_f =0.46, chloroform:methanol (9:1); NMR: 8.62 (s,1), 7.52 (dd,2), 7.26 (dd,2), 4.6 (dd,2), 4.04 (d,1), 2.25 (m,1), 1.09 (d,6), 0.85 (d,3), 0.76 (d,3); MS: m/z =500(M+1).

Analysis for $C_{22}H_{25}F_4N_5O_4 \cdot 0.25 H_2O$:			
Calculated:	C, 52.43;	H, 5.10;	N, 13.90
Found:	C, 52.78;	H, 5.24;	N, 13.58

EXAMPLE 114

2-(6-Oxo-2-phenyl-5-ureido-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0312] To a solution of the compound from Example 6 (0.40 g) in tetrahydrofuran (10 mL) cooled to 0 °C was added dropwise chlorosulfonyl isocyanate (0.16 g) and the mixture was stirred for 45 min. The reaction mixture was neutralized with saturated aqueous sodium carbonate solution (3 mL), diluted with ethyl acetate (10 mL), and the separated organic phase was washed (water, brine). TLC examination revealed that the major reaction product was present in the aqueous phase. Following saturation of the aqueous phase with sodium chloride, repeated ethyl acetate extractions (4 x 25 mL) were performed. The combined organic layers were dried (magnesium sulfate) and evaporated. Chromatography with dichloromethane:tetrahydrofuran (10:1) as the eluent, followed by overnight drying (50 °C, 27 Pa) gave the title compound (0.061 g) as an off-white solid; TLC: R_f =0.36, dichloromethane:methanol (10:1); NMR: 8.85 (d,1), 8.69 (s,1), 8.35 (s,1), 7.48 (m,5), 6.47 (broad s, 2), 4.70 (dd,1), 4.55 (dd,2), 2.2 (m,1), 0.89 (dd,6); MS: m/z =440(M+1).

Analysis for $C_{19}H_{20}F_3N_5O_4 \cdot 0.60 H_2O$:			
Calculated:	C, 50.69;	H, 4.75;	N, 15.56
Found:	C, 50.82;	H, 4.83;	N, 15.24

EXAMPLE 115

2-[2-(4-Fluorophenyl)-5-methylsulfonylamino-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0313] To a solution of the compound from Example 15 (0.45 g) in tetrahydrofuran (11 mL) was added pyridine (0.89 mL) and the mixture cooled in an ice bath to 0 °C. Methylsulfonyl chloride (0.17 mL) was added and the solution warmed to room temperature and allowed to stir for 24 h. The reaction mixture was poured into ethyl acetate and washed with a saturated solution of potassium dihydrogen phosphate, and then H₂O. The resulting solution was dried and the solvent removed by evaporation. The resulting material was purified by chromatography, eluting with tetrahydrofuran:dichloromethane (gradient, 10:90, 20:80) to give the title compound (0.48 g) as a white powder; TLC: R_f =0.67, tetrahydrofuran:dichloromethane (20:80); NMR (DMSO/D₂O): 7.97 (s,1), 7.55 (m,2), 7.27 (t,2), 4.68 (d,1), 4.45 (d,1), 4.00 (d,1), 3.07 (s,3), 2.16 (s,1), 0.83 (d,3), 0.72 (d,3).

Analysis for $C_{19}H_{20}F_4N_4O_5S \cdot 0.3 H_2O$:			
Calculated:	C, 45.8;	H, 4.17;	N, 11.3

(continued)

Analysis for $C_{19}H_{20}F_4N_4O_5S \cdot 0.3 H_2O$:			
Found:	C, 45.8;	H, 4.32;	N, 11.0

EXAMPLES 116-125

[0314] Using procedures similar to that described in Example 115 and using the required sulfonyl chloride, the following compounds of formula I wherein R^0 is isopropyl and R and R^6 have the indicated values were prepared from the corresponding compounds of formula I wherein R is hydrogen.

[0315] Example 116: R=methylsulfonyl, R^6 =3-pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 10:90); TLC: R_f =0.40, methanol/dichloromethane (7:93); NMR: 8.70 (d,1), 8.81 (t,2), 8.03 (s,1), 7.90 (m,2), 7.48 (m,1), 4.72 (d,1), 4.51 (d,1), 4.00 (d,1), 3.11 (s,3), 2.19 (m,1), 0.82 (d,3), 0.70 (d,3).

Analysis for $C_{18}H_{20}F_3N_4O_5S \cdot 0.4 H_2O$:			
Calculated:	C, 44.8 ;	H, 4.34;	N, 14.5
Found:	C, 44.7 ;	H, 4.35;	N, 14.3

[0316] Example 117: R=methylsulfonyl, R^6 =2-thienyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 8:92); TLC: R_f =0.34, tetrahydrofuran:dichloromethane (10:90); NMR: 7.95 (s,1), 7.83 (d,1), 7.42 (d,1), 7.12 (t,1), 4.85 (d,1), 4.51 (m,2), 4.08 (d,1), 3.06 (s,3), 2.25 (m,1), 0.90 (d,3), 0.75 (d,3).

Analysis for $C_{17}H_{19}F_3N_4O_5S_2$:			
Calculated:	C, 42.5 ;	H, 3.99;	N, 11.7
Found:	C, 42.1 ;	H, 4.12;	N, 11.4

[0317] Example 118: R=methylsulfonyl, R^6 =4-trifluoromethylphenyl: Chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f =0.12, methanol:dichloromethane (5:95); NMR (DMSO- D_2O): 7.99 (s,1), 7.79 (d,2), 7.67 (d,2), 4.59 (m,2), 3.99 (d,1), 3.09 (s,3), 2.15 (m,1), 0.75 (d,3), 0.63 (d,3).

Analysis for $C_{20}H_{20}F_6N_4O_5S \cdot 0.5 H_2O$:			
Calculated:	C, 43.5 ;	H, 3.84;	N, 10.1
Found:	C, 43.6 ;	H, 3.91;	N, 10.5

[0318] Example 119: R=4-methoxyphenylsulfonyl, R^6 =4-trifluoromethylphenyl: Chromatography solvent: ethyl acetate:hexane (gradient, 50:50, 100:0); TLC: R_f =0.57, ethyl acetate; NMR (DMSO- D_2O): 7.89 (s,1), 7.82 (d,2), 7.74 (d,2), 7.60 (d,2), 7.06 (d,2), 4.49 (m,2), 3.88 (d,1), 3.78 (s,3), 2.12 (m,1), 0.71 (d,3), 0.59 (d,3).

Analysis for $C_{26}H_{24}F_6N_4O_6S$:			
Calculated:	C, 49.2 ;	H, 3.81;	N, 8.83
Found:	C, 49.0 ;	H, 3.77;	N, 8.77

[0319] **Example 120:** R=methylsulfonyl, R⁶=3,5-difluorophenyl: Chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 15:85, 30:70); TLC: R_f=0.41, tetrahydrofuran:dichloromethane (10:90); NMR (DMSO-D₂O): 7.96 (s,1), 7.40 (m,1), 7.19 (d,2), 4.68 (d,1), 4.40 (d,1), 3.99 (s,1), 3.08 (s,3), 2.20 (m,1), 0.81 (d,3), 0.70 (d,3).

Analysis for C ₁₉ H ₁₉ F ₅ N ₄ O ₅ S • 0.6 H ₂ O:			
Calculated:	C, 43.8 ;	H, 3.90;	N, 10.8
Found:	C, 43.8 ;	H, 3.80;	N, 10.6

[0320] **Example 121:** R=phenylsulfonyl, R⁶=3,5-difluorophenyl: Chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 5:95 to 15:85); TLC: R_f=0.50, tetrahydrofuran:dichloromethane (17:83); NMR (DMSO/D₂O): 7.90 (m,3), 7.60 (m,3), 7.40 (t,1), 7.17 (d,2), 4.60 (d,1), 4.35 (d,1), 3.96 (d,1), 2.20 (m,1), 0.81 (d,3), 0.67 (d,3).

Analysis for C ₂₄ H ₂₁ F ₅ N ₄ O ₅ S:			
Calculated:	C, 50.4 ;	H, 3.70;	N, 9.79
Found:	C, 50.0 ;	H, 3.72;	N, 9.69

[0321] **Example 122:** R=phenylsulfonyl, R⁶=4-nitrophenyl: Chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f=0.40, methanol:dichloromethane (5:95); 300 MHz NMR (DMSO/D₂O): 8.21 (d,2), 7.86 (m,3), 7.63 (m,4), 4.52 (m,2), 3.89 (d,1), 2.12 (m,1), 0.72 (d,3), 0.60 (d,3).

Analysis for C ₂₄ H ₂₂ F ₃ N ₅ O ₇ S:			
Calculated:	C, 49.6;	H, 3.81;	N, 12.0
Found:	C, 49.4;	H, 4.00;	N, 12.0

[0322] **Example 123:** R=4-chlorophenylsulfonyl, R⁶=4-trifluoromethylphenyl: Chromatography solvent: ethyl acetate:dichloromethane (50:50); TLC: R_f=0.55, ethyl acetate; 300 MHz NMR (DMSO/D₂O): 7.96 (s,1), 7.85 (d,2), 7.76 (d,2), 7.60 (m,4), 4.60 (d,1), 4.38 (d,2), 3.89 (d,1), 2.11 (m,1), 0.71 (d,3), 0.59 (d,3).

Analysis for C ₂₅ H ₂₁ F ₆ N ₄ O ₅ S:			
Calculated:	C, 47.0;	H, 3.31;	N, 8.77
Found:	C, 46.7;	H, 3.36;	N, 8.74

[0323] **Example 124:** R=methylsulfonyl, R⁶=4-methoxyphenyl: Chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f=0.28, methanol:dichloromethane (5:95); 300 MHz NMR: 9.30 (s,1), 8.86 (d,1), 7.98 (s,1), 7.46 (d,2), 7.02 (d,2), 4.67 (t,1), 4.58 (m,2), 3.82 (s,3), 3.07 (s,3), 2.17 (t,1), 0.90 (d,3), 0.84 (d,3).

Analysis for C ₂₀ H ₂₃ F ₃ N ₄ O ₆ S:			
Calculated:	C, 47.6;	H, 4.60;	N, 11.1

(continued)

Analysis for $C_{20}H_{23}F_3N_4O_6S$:			
Found:	C, 47.4;	H, 4.63;	N, 11.0

[0324] Example 125: R=5-dimethylamino-1-naphthylsulfonyl, R⁶=phenyl: Chromatography solvent: ethyl acetate:dichloromethane (25:75); TLC: R_F=0.16, ethyl acetate:dichloromethane (25:75); NMR: (DMSO/D₂O) 8.47 (d,1), 8.33 (d,1), 8.26 (d,2), 7.81 (s,1), 7.61 (m,2), 7.36 (m,5), 7.25 (d,1), 4.39 (dd,2), 2.79 (s,6), 2.17 (m,1), 0.76 (d,3), 0.69 (d,3).

Analysis for $C_{30}H_{30}F_3N_5O_5S \cdot 0.5 H_2O$:			
Calculated:	C, 56.4;	H, 4.89;	N, 11.0
Found:	C, 56.2;	H, 4.94;	N, 11.7

EXAMPLE 126

2-[2-(4-Dimethylaminophenyl)-5-methylsulfonylamino-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0325] To a solution of the product from Example 136 (300 mg) in ethanol (50 mL) and tetrahydrofuran (5 mL) was added formaldehyde (2 mL of a 37% solution of formaldehyde in H₂O) and 10% (w/w) palladium on carbon (100 mg). The resulting mixture was placed under a hydrogen atmosphere (3.5 bar) and shaken overnight. The solution was filtered, dried, and evaporated. The resulting oil was chromatographed, eluting with methanol:dichloromethane (gradient, 5:95, 10:90), to give a solid which was washed with diethyl ether:hexane (1:1) to provide the title compound (0.23 g); TLC: R_F=0.2, methanol:methylene chloride (5:95); NMR (DMSO-D₂O): 7.91 (d,1), 7.33 (d,2), 6.87 (d,2), 4.58 (m,2), 3.96 (d,1), 3.01 (s,3), 2.95 (s,6), 2.20 (m,1), 0.82 (d,3), 0.74 (d,3).

Analysis for $C_{21}H_{26}F_3N_5O_5S \cdot 0.3 H_2O$:			
Calculated:	C, 48.2;	H, 5.12;	N, 13.4
Found:	C, 48.5;	H, 5.53;	N, 13.1

EXAMPLE 127

2-[2-(4-Aminophenyl)-5-methylsulfonylamino-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0326] The product from Example 136 was subjected to a hydrogenation procedure similar to the hydrogenolysis described in Example 2.b. to give the title compound; chromatography solvent: methanol:dichloromethane (gradient, 5:95, 15:85); TLC: R_F=0.25, methanol:dichloromethane (5:95); NMR (DMSO/D₂O): 7.90 (d,1), 7.20 (d,2), 6.58 (d,2), 4.70 (m,2), 4.00 (d,1), 3.01 (s,3), 2.21 (m,1), 0.86 (d,3), 0.78 (d,3).

Analysis for $C_{19}H_{22}F_3N_5O_5S \cdot 0.5 H_2O$:			
Calculated:	C, 45.9;	H, 4.45;	N, 14.1
Found:	C, 46.0;	H, 4.65;	N, 13.9

EXAMPLES 128-129

[0327] Using procedures similar to that described in Example 1, the following compounds of formula I, wherein R⁰ is isopropyl and R and R⁶ have the indicated values were prepared by oxidation of the corresponding alcohols of formula II.

[0328] Example 128: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=phenyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 10:90); TLC: R_F=0.50, methanol:dichloromethane (5:95); NMR (DMSO/D₂O): 8.41 (s,1), 7.50 (m,5), 7.06 (s,2), 5.11 (s,2), 4.50 (m,2), 4.01 (m,1), 2.20 (m,1), 0.80 (d,3), 0.73 (d,3).

Analysis for C ₂₇ H ₂₈ F ₃ N ₅ O ₅ • 0.1 H ₂ O:			
Calculated:	C, 57.8;	H, 5.06;	N, 12.5
Found:	C, 57.6;	H, 5.00;	N, 12.4

[0329] Example 129: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=4-fluorophenyl: Chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 10:90 to 40:60); TLC: R_F=0.33, tetrahydrofuran:dichloromethane (30:70); NMR (DMSO/D₂O): 8.45 (s,1), 7.54 (m,2), 7.26 (t,2), 7.09 (s,2), 5.14 (s,2), 4.68 (d,1), 4.42 (d,1), 4.03 (d,1), 2.41 (s,6), 2.21 (m,1), 0.84 (d,3), 0.74 (d,3).

Analysis for C ₂₇ H ₂₇ F ₄ N ₅ O ₅ • 0.5 H ₂ O:			
Calculated:	C, 55.3;	H, 4.81;	N, 11.9
Found:	C, 55.2;	H, 4.85;	N, 12.0

[0330] The intermediate alcohols of formula II used in Examples 128-129 were prepared as follows.

EXAMPLES 128.a.-129.a.

[0331] The following compounds of formula XIV wherein R⁰ is isopropyl, R_p is *tert*-butyldimethylsilyl and R and R⁶ have the indicated values were prepared from the corresponding compounds of formula XIII by a procedure similar to that described in Example 3.a. except using the required carbinol.

[0332] Example 128.a.: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=phenyl: Chromatography solvent: ethyl acetate:dichloromethane (gradient, 50:50, 75:25); TLC: R_F=0.5, methanol:dichloromethane (3:97); MS: m/z=676(M+1).

[0333] Example 129.a.: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=4-fluorophenyl: Chromatography solvent: ethyl acetate:dichloromethane (gradient, 50:50 to 75:25); TLC: R_F=0.38, methanol:dichloromethane (5:95); MS: m/z=694 (M+1).

EXAMPLES 128.b.-129.b.

[0334] The following compounds of formula II wherein R⁰ is isopropyl and R and R⁶ have the indicated values were prepared from the corresponding compounds of formula XIV, described in Examples 128.a.-129.a., by procedures similar to that described in Example 2.d.

[0335] Example 128.b.: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=phenyl: Purified by crystallization from diethyl ether:hexane (1:1); TLC: R_F=0.20, ethyl acetate; MS: m/z=562(M+1).

[0336] Example 129.b.: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=4-fluorophenyl: Purified by crystallization from diethyl ether:ethyl acetate (1:1); TLC: R_F=0.30, ethyl acetate:dichloromethane (75:25); MS: m/z=580 (M+1).

EXAMPLES 130-132

[0337] Using procedures similar to that described in Example 1, the following compounds of formula I wherein R⁰ is isopropyl and R and R⁶ have the indicated values were prepared by oxidation of the corresponding alcohols of formula II.

[0338] Example 130: R=4-pyridylmethoxycarbonyl, R⁶=3-pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 10:90); TLC: R_F=0.30, methanol:dichloromethane (10:90); 300 MHz NMR (DMSO/D₂O): 8.67 (m,2), 8.55 (m,2), 8.47 (s,1), 7.90 (m,2), 7.44 (m,3), 5.23 (s,2), 4.67 (m,2), 3.99 (m,1), 2.19 (m,1), 0.82 (d,3), 0.69 (d,3).

Analysis for C ₂₄ H ₂₃ F ₃ N ₆ O ₅ • 0.6 H ₂ O:			
Calculated:	C, 53.1;	H, 4.49;	N, 15.5
Found:	C, 53.0;	H, 4.53;	N, 15.5

[0339] Example 131: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=3,5-difluorophenyl: Chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 15:85, 40:60); TLC: R_F=0.37, tetrahydrofuran:dichloromethane (25:75); NMR (DMSO/D₂O): 8.45 (s,1), 7.42 (t,1), 7.20 (d,2), 7.08 (s,2), 5.13 (s,2), 4.70 (d,1), 4.45 (d,1), 4.01 (d,1), 2.40 (s,6), 2.23 (m,1), 0.84 (d,3), 0.73 (d,3).

Analysis for C ₂₇ H ₂₆ F ₅ N ₅ O ₅ • 0.5 H ₂ O:			
Calculated:	C, 53.6;	H, 4.50;	N, 11.6
Found:	C, 53.4;	H, 4.47;	N, 11.6

[0340] Example 132: R=2,6-dimethylpyrid-4-ylmethoxycarbonyl, R⁶=4-methoxyphenyl: Chromatography solvent: methanol:dichloromethane (5:95); TLC: R_F=0.21, methanol:dichloromethane (5:95); NMR (DMSO/D₂O): 8.44 (s,1), 7.45 (d,2), 7.11 (s,2), 6.99 (d,2), 5.16 (s,2), 4.60 (dd,2), 4.05 (d,1), 3.81 (s,3), 2.44 (s,6), 2.25 (m,1), 0.87 (d,3), 0.78 (d,3).

Analysis for C ₂₈ H ₃₀ F ₃ N ₅ O ₆ • 0.7 H ₂ O:			
Calculated:	C, 55.9;	H, 5.25;	N, 11.6
Found:	C, 55.7;	H, 5.27;	N, 11.7

[0341] The intermediate alcohols of formula II, used in Examples 130-132, were prepared as follows.

EXAMPLES 130.a-132.a.

[0342] Using procedures similar to that outlined in Example 2.a., the following compounds of formula XII wherein R⁰ is isopropyl, R_p is *tert*-butyldimethylsilyl and R⁶ has the indicated value were prepared from the corresponding compounds of formula II.

[0343] Example 130.a: R⁶=3-pyridyl: Chromatography solvent: diethyl ether:hexane (gradient, 80:20 to 100:0); TLC: R_F=0.35, methanol:dichloromethane (5:95); MS: m/z=648(M+1).

[0344] Example 131.a: R⁶=3,5-difluorophenyl: Purified by recrystallization from diethyl ether:hexane (1:1); TLC: R_F=0.6, tetrahydrofuran:dichloromethane (5:95); MS: m/z=683(M+1).

[0345] Example 132.a: R⁶=4-methoxyphenyl: Chromatography solvent: ethyl acetate:dichloromethane (10:90); TLC: R_F=0.33, ethyl acetate:dichloromethane (10:90); MS: m/z=677(M+1).

EXAMPLES 130.b-132.b

[0346] The following compounds of formula XIII wherein R⁰ is isopropyl, R_p is *tert*-butyldimethylsilyl and R⁶ has the indicated value were prepared from the corresponding compounds of formula XII, described in Examples 130.a.-132.a., by procedures similar to that described in Example 2.b.

[0347] Example 130.b.: $R^6=3$ -pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 10:90); TLC: $R_f=0.20$, methanol:dichloromethane (5:95); MS: $m/z=514(M+1)$.

[0348] Example 131.b.: $R^6=3,5$ -difluorophenyl: Purified by recrystallization from diethyl ether:hexane (1:1); TLC: $R_f=0.44$, tetrahydrofuran:dichloromethane (15:85); MS: $m/z=549(M+1)$.

5 [0349] Example 132.b.: $R^6=4$ -methoxyphenyl: Purified by recrystallization from diethyl ether:hexane (1:1); TLC: $R_f=0.55$, methanol:dichloromethane (10:90); MS: $m/z=543(M+1)$.

EXAMPLES 130.c-132.c.

10 [0350] The following compounds of formula XIV wherein R^0 is isopropyl, R_p is *tert*-butyldimethylsilyl and R and R^6 have the indicated values were prepared from the corresponding compounds of formula XIII, described in Examples 130.b.-132.b., by procedures similar to that described in Example 3.a., except using the required carbinol.

[0351] Example 130.c.: $R=4$ -pyridylmethoxycarbonyl, $R^6=3$ -pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95, 10:90); TLC: $R_f=0.3$, methanol:dichloromethane (7:93); MS: $m/z=649(M+1)$.

15 [0352] Example 131.c.: $R=2,6$ -dimethylpyrid-4-ylmethoxycarbonyl, $R^6=3,5$ -difluorophenyl: Chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 5:95, 15:85); TLC: $R_f=0.43$, tetrahydrofuran:dichloromethane (10:90); MS: $m/z=712(M+1)$.

[0353] Example 132.c.: $R=2,6$ -dimethylpyrid-4-ylmethoxycarbonyl, $R^6=4$ -methoxyphenyl: Chromatography solvent: methanol:dichloromethane (5:95); TLC: $R_f=0.25$, methanol:dichloromethane (5:95); MS: $m/z=706(M+1)$.

20

EXAMPLES 130.d.-132.d.

[0354] The following compounds of formula II wherein R^0 is isopropyl and R and R^6 have the indicated values were prepared from the corresponding compounds of formula XIV, described in Examples 130.c.-132.c., by procedures similar to that described in Example 2.d.

25

[0355] Example 130.d.: $R=4$ -pyridylmethoxycarbonyl, $R^6=3$ -pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 5:95 to 10:90); TLC: $R_f=0.45$, methanol:dichloromethane (5:95); MS: $m/z=535(M+1)$.

[0356] Example 131.d.: $R=2,6$ -dimethylpyrid-4-ylmethoxycarbonyl, $R^6=3,5$ -difluorophenyl: Purified by crystallization from diethyl ether:tetrahydrofuran (4:1); TLC: $R_f=0.17$, tetrahydrofuran:dichloromethane (15:85); MS: $m/z=598(M+1)$.

30

[0357] Example 132.d.: $R=2,6$ -dimethylpyrid-4-ylmethoxycarbonyl, $R^6=4$ -methoxyphenyl: Chromatography solvent, methanol:dichloromethane (5:95); TLC: $R_f=0.50$, methanol:dichloromethane (10:90); MS: $m/z=592(M+1)$.

EXAMPLES 133-134

35

[0358] Using procedures similar to that described in Example 1 the following compounds of formula I wherein R^0 is isopropyl and R and R^6 have the indicated values were prepared by oxidation of the corresponding alcohols of formula II.

40

[0359] Example 133: R =ethoxycarbonyl, $R^6=3$ -pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 4:96 to 8:92); TLC: $R_f=0.44$, tetrahydrofuran:dichloromethane (40:60); NMR (DMSO- D_6): 8.88 (m,2), 8.45 (s,1), 7.90 (m,2), 7.45 (m,1), 4.73 (d,1), 4.50 (d,1), 4.15 (m,2), 3.99 (m,1), 2.20 (m,1), 1.23 (t,3), 0.82 (d,3), 0.70 (d,3).

45

Analysis for $C_{20}H_{22}F_3N_5O_5 \cdot 0.6 H_2O$:			
Calculated:	C, 50.0;	H, 4.87;	N, 14.5
Found:	C, 49.7;	H, 4.88;	N, 14.4

50

[0360] Example 134: R =isopropoxycarbonyl, $R^6=3$ -pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 4:96, 8:92); TLC: $R_f=0.50$, tetrahydrofuran:dichloromethane (35:65); NMR (DMSO- D_6): 8.69 (s,2), 8.45 (s,1), 7.91 (d,1), 7.46 (m,1), 4.87 (m,1), 4.6 (m,2), 4.00 (m,1), 2.20 (m,1), 1.26 (d,6), 0.82 (d,3), 0.70 (d,3).

55

Analysis for $C_{21}H_{24}F_3N_5O_5 \cdot 1.0 H_2O$:			
Calculated:	C, 50.3;	H, 5.22;	N, 14.0
Found:	C, 50.3;	H, 5.16;	N, 13.6

[0361] The intermediate alcohols of formula II, used in Examples 133-134, were prepared as follows

EXAMPLES 133.a.-134.a.

[0362] The following compounds of formula XIV wherein R^0 is isopropyl, R_p is *tert*-butyldimethylsilyl and R and R^6 have the indicated values were prepared from the corresponding compounds of formula XIII and the indicated chloroformate using procedures similar to that described in Example 35.

[0363] Example 133.a.: R=ethoxycarbonyl, R^6 =3-pyridyl: Using ethyl chloroformate; chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 10:90 to 30:70); TLC: R_f =0.48, methanol:dichloromethane (10:90); MS: m/z =586(M+1).

[0364] Example 134.a.: R=isopropoxycarbonyl, R^6 =3-pyridyl: Using isopropyl chloroformate; chromatography solvent: tetrahydrofuran:dichloromethane (gradient, 10:90, 30:70); TLC: R_f =0.43, methanol:dichloromethane (10:90); MS: m/z =600(M+1).

EXAMPLES 133.b-134.b.

[0365] The following compounds of formula II wherein R^0 is isopropyl, and R and R^6 have the indicated values were prepared from the corresponding compounds of formula XIV, described in Examples 133.a.-134.a., by procedures similar to that described in Example 2.d.

[0366] Example 133.b.: R=ethoxycarbonyl, R^6 =3-pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 4:96, 10:90); TLC: R_f =0.25, tetrahydrofuran:dichloromethane (35:65); MS: m/z =472(M+1).

[0367] Example 134.b.: R=isopropoxycarbonyl, R^6 =3-pyridyl: Chromatography solvent: methanol:dichloromethane (gradient, 4:96, 8:92); TLC: R_f =0.24, methanol:dichloromethane (8:92); MS: m/z =486(M+1).

EXAMPLE 135

[0368] 2-[5-Amino-2-(4-hydroxyphenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0368] To a suspension of the product from Example 28 (0.68 g) in dichloromethane (20 mL) was added boron tribromide (32 mL, 1 M solution in dichloromethane) and the resulting solution was allowed to stir for 3 days. The excess boron tribromide was quenched by addition of methanol (5 mL) and the solution was brought to neutral pH by addition of sodium bicarbonate. The product was extracted into ethyl acetate, the solution dried, and the solvent was evaporated. The resulting material was purified by chromatography, eluting with methanol:dichloromethane (5:95), to give the title compound (0.4 g) as a white solid; TLC: R_f =0.3, methanol:dichloromethane (5:95); NMR (DMSO- d_6): 7.32 (s,1), 7.22 (dd,2), 6.74 (dd,2), 4.48 (dd,2), 4.1 (m,1), 2.20 (m,1), 0.83 (d,3), 0.76 (d,3).

Analysis for $C_{18}H_{19}F_3N_4O_4$:			
Calculated:	C, 52.4;	H, 4.64;	N, 13.6
Found:	C, 52.4;	H, 4.80;	N, 13.3

EXAMPLE 136

2-[5-Methylsulfonylamino-2-(4-nitrophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0369] The product from Example 33 was subjected to a procedure similar to that described in Example 115 to yield crude material, which was purified by trituration with ether, to give the title compound as a white solid; TLC: R_f =0.4, methanol:dichloromethane (5:95); NMR (DMSO/ D_2O): 8.29 (d,2), 7.97 (s,1), 7.70 (d,2), 4.55 (m,2), 3.92 (d,1), 3.08 (s,3), 2.13 (m,1), 0.73 (d,3), 0.64 (d,3).

Analysis for $C_{19}H_{20}F_3O_7N_5S$:			
Calculated:	C, 43.9;	H, 3.88;	N, 13.5
Found:	C, 43.8;	H, 3.95;	N, 13.3

EXAMPLE 137

2-[5-Phenylsulfonylamino-2-(4-methoxyphenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0370] The product from Example 28 was subjected to a procedure similar to that described in Example 115 substituting phenyl sulfonylchloride for methyl sulfonyl chloride to give the title compound as a white solid; chromatography solvent: methanol:dichloromethane (5:95); TLC: R_f =0.38, methanol:dichloromethane (5:95); NMR: 10.06 (s,1), 8.81 (d,1), 7.91 (m,3), 7.61 (m,3), 7.40 (d,2), 6.99 (d,2), 4.64 (t,1), 4.50 (dd,2), 3.80 (s,3), 2.15 (m,1), 0.88 (d,3), 0.82 (d,3).

Analysis for $C_{25}H_{25}F_3N_4O_6S$:			
Calculated:	C, 53.00;	H, 4.45;	N, 9.89
Found:	C, 52.95;	H, 4.61;	N, 9.86

EXAMPLE 138

2-[2-(4-Fluorophenyl)-6-oxo-5-ureido-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0371] A flask was charged with 0.4 g of the product from Example 15 dissolved in a mixture of water (6 mL), tetrahydrofuran (1 mL) and acetic acid (4 mL) at room temperature. To this was added 0.31 g sodium isocyanate and the reaction mixture stirred for 1.5 h at room temperature and then 0.5 h at 55 °C. Additional sodium isocyanate, 2 x 70 mg and 1 x 35 mg, was sequentially added over an additional 1 h. The mixture was cooled, diluted with ethyl acetate, and partitioned with water. The organic layer was dried over sodium sulfate, filtered and evaporated to afford a residual solid which was triturated with diethyl ether. The residual solid was collected and dried under vacuum to afford the title compound (200 mg) as an off-white solid; TLC: R_f =0.20, dichloromethane:methanol (15:1); NMR (DMSO/ D_2O): 8.65 (s,1), 7.54 (m,2), 7.28 (m,2), 4.56 (dd,2), 4.05 (d,1), 2.24 (m,1), 0.82 (dd,6); MS: m/z =458(M+1).

Analysis for $C_{19}H_{19}F_4N_5O_4 \cdot 0.75 H_2O$:			
Calculated:	C, 48.46;	H, 4.39;	N, 14.87
Found:	C, 48.10;	H, 4.21;	N, 15.16

EXAMPLES 139-153

[0372] Using procedures similar to that described in Example 90 and using the required acyl or sulfonyl chloride, the following compounds of formula I wherein R^0 is isopropyl and R and R^6 have the indicated values were prepared from the corresponding compounds of formula I wherein R is hydrogen.

[0373] Example 139: R=methoxyaminocarbonyl, R^6 =phenyl: Chromatography solvent: dichloromethane:methanol (3:1) followed by trituration with dichloromethane:diethyl ether; TLC: R_f =0.29, dichloromethane:methanol (20:1); NMR (DMSO/ D_2O): 8.63 (s,1), 7.47 (m,5), 4.63 (d,1), 4.06 (d,1), 2.34 (m,1), 0.85 (d,3), 0.77 (d,3); MS: m/z =470(M+1).

Analysis for $C_{20}H_{22}F_3N_5O_5 \cdot 0.5 H_2O$:			
Calculated:	C, 50.21;	H, 4.84;	N, 14.64
Found:	C, 50.33;	H, 4.54;	N, 14.26

[0374] Example 140: R=acetylaminosulfonyl, R^6 =4-fluorophenyl: Purified by crystallization from diethyl ether; TLC: R_f =0.14, chloroform:methanol (6:1); NMR (DMSO/ D_2O): 7.99 (s,1), 7.53 (dd,2), 7.28 (t,2), 4.67 (d,1), 4.50 (d,1), 2.2 (m,1), 1.95 (s,3), 0.84 (d,3), 0.74 (d,3); MS: FAB m/z =536(M+1), m/z =534(M-1).

Analysis for $C_{20}H_{21}F_4N_5O_6S$:			
Calculated:	C, 44.86;	H, 3.95;	N, 13.08
Found:	C, 45.19;	H, 4.00;	N, 13.21

[0375] Example 141: R=acetylaminosulfonyl, R^6 =phenyl: Purified by crystallization from diethyl ether; TLC: R_f =0.23, chloroform:methanol (6:1); NMR (DMSO/ D_2O): 8.00 (s,1), 7.57-7.47 (m,5), 4.61 (d,1), 4.50 (d,1), 2.24 (m,1), 1.96 (s,3), 0.85 (d,3), 0.77 (d,3); MS: FAB m/z =518(M+1), 516(M-1); High resolution exact mass analysis for $C_{20}H_{23}F_3N_5O_6S$ (M+1): Calculated: m/z =518.1321, Found: m/z =518.1319.

Analysis for $C_{20}H_{22}F_3N_5O_6S$:			
Calculated:	C, 46.42;	H, 4.28;	N, 13.53
Found:	C, 46.85;	H, 4.42;	N, 13.60

[0376] Example 142: R=benzoylaminosulfonyl, R^6 =phenyl: Purified by crystallization from ethyl acetate:ether; TLC: R_f =0.18, chloroform:methanol (6:1); NMR (DMSO/ D_2O): 8.07 (s,1), 7.84 (m,2), 7.64 (m,1), 7.54-7.45 (m,7), 4.60 (d,1), 4.46 (d,1), 2.20 (m,1), 0.82 (d,3), 0.72 (d,3); MS: FAB m/z =580(M+1), m/z =578(M-1).

Analysis for $C_{25}H_{24}F_3N_5O_6S \cdot 0.7 H_2O$:			
Calculated:	C, 50.71;	H, 4.32;	N, 11.83
Found:	C, 50.62;	H, 4.33;	N, 11.82

[0377] Example 143: R=methoxycarbonylaminosulfonyl, R^6 =2-thienyl: Purified by trituration with methyl *tert*-butyl ether; TLC: R_f =0.14, chloroform:methanol (6:1); NMR (DMSO/ D_2O): 7.97 (s,1), 7.82 (d,1), 7.43 (d,1), 7.15 (dd,1), 4.89 (dd,2), 3.64 (s,3), 2.26 (m,1), 0.93 (d,3), 0.78 (d,3); MS: FAB m/z =540(M+1), 538(M-1).

Analysis for $C_{18}H_{20}F_3N_5O_7S_2$:			
Calculated:	C, 40.07;	H, 3.74;	N, 12.98
Found:	C, 39.91;	H, 3.74;	N, 12.87

10 **[0378] Example 144:** R=4-acetylaminoxyphenylsulfonyl, R⁶=4-fluorophenyl: Purified by titration with diethyl ether; TLC: R_F=0.22, dichloromethane:methanol (20:1); NMR (DMSO/D₂O): 7.93 (s,1), 7.85 (d,2, J=8.0), 7.74 (d,2, J=8.0), 7.51 (m,2), 7.25 (dd,2, J=8.8, 6.8), 4.60 (d,1, J=17.0), 4.39 (d,1, J=16.8), 3.99 (m,1), 2.20 (m,1), 2.09 (s,3), 0.82 (d,3, J=6.4), 0.71 (d,3, J=6.4); MS: m/z=612(M+1).

Analysis for $C_{26}H_{25}F_4N_5O_6S \cdot 0.5 H_2O$:			
Calculated:	C, 50.32;	H, 4.22;	N, 11.28
Found:	C, 50.19;	H, 4.15;	N, 11.24

20 **[0379] Example 145:** R=4-acetylaminoxyphenylsulfonyl, R⁶=phenyl: Purified by titration with diethyl ether; TLC: R_F=0.25, dichloromethane:methanol (20:1); NMR (DMSO/D₂O): 7.94 (s,1), 7.86 (d,2, J=8.8), 7.74 (d,2, J=8.8), 7.53 (m,1), 7.45 (m,4), 4.56 (d,1, J=16.8), 4.37 (d,1, J=16.8), 4.02 (d,1, J=2.8), 2.21 (m,1), 2.09 (s,3), 0.80 (d,3, J=6.8); 0.74 (d,3, J=6.8); MS: m/z=594(M+1).

Analysis for $C_{26}H_{26}F_3N_5O_6S$:			
Calculated:	C, 52.21;	H, 4.46;	N, 11.71
Found:	C, 52.24;	H, 4.52;	N, 11.66

30 **[0380] Example 146:** R=4-acetylaminoxyphenylsulfonyl, R⁶=2-thienyl: Purified by recrystallization from ethyl acetate; TLC: R_F=0.47, dichloromethane:methanol (10:1); NMR (DMSO/D₂O): 7.91 (s,1), 7.83 (d,2, J=8.8), 7.81 (d,1, J=4.7), 7.74 (d,2, J=8.8), 7.38 (d,1, J=4.7), 7.12 (dd,1, J=4.2, 4.6), 4.86 (d,1, J=16.6), 4.74 (d,1, J=16.6), 4.06 (d,1, J=2.7), 2.24 (m,1), 2.09 (s,3), 0.90 (d,3, J=6.8), 0.74 (d,3, J=6.8); MS: m/z=600(M+1).

Analysis for $C_{24}H_{24}F_3N_5O_6S_2 \cdot 0.5 H_2O$:			
Calculated:	C, 47.36;	H, 4.14;	N, 11.51
Found:	C, 47.58;	H, 4.06;	N, 11.56

45 **[0381] Example 147:** R=benzoylaminoxyphenylsulfonyl, R⁶=2-thienyl: Purified by titration with ethyl acetate=diethyl ether; TLC: R_F=0.14, dichloromethane:methanol (10:1); NMR (DMSO/D₂O): 8.04 (s,1), 7.84 (m,3), 7.65 (dd,1, J=7.2, 7.5), 7.52 (dd,2, J=7.5, 7.8), 7.42 (d,1, J=3.6), 7.13 (dd,1, J=4.0, 4.9), 4.91 (d,1, J=16.8), 4.80 (d,1, J=16.8), 4.08 (d,1, J=2.8), 2.24 (m, 1), 0.91 (d,3, J=6.8), 0.74 (d,3, J=6.8); MS: m/z=586(M+1).

Analysis for $C_{23}H_{22}F_3N_5O_6S_2 \cdot 0.5 H_2O$:			
Calculated:	C, 46.46;	H, 3.90;	N, 11.78

(continued)

Analysis for $C_{23}H_{22}F_3N_5O_6S_2 \cdot 0.5 H_2O$:			
Found:	C, 46.63;	H, 3.85;	N, 11.79

[0382] Example 148: R=benzylaminosulfonyl, R⁶=2-thienyl: Purified by trituration with hexane:diethyl ether (20:1); TLC: R_f=0.42, dichloromethane:methanol (20:1); NMR (DMSO/D₂O): 7.90 (s,1), 7.82 (d,1), 7.40 (d,1), 7.38 (m,5), 7.15 (t,1), 4.83 (q,2), 4.11 (s,2), 2.28 (m,1), 0.95 (d,3), 0.80 (d,3); MS: m/z=572(M+1) also exact mass for C₂₃H₂₄F₃N₅O₅S₂, Calculated: 571.1171, Found: 571.1170.

Analysis for $C_{23}H_{24}F_3N_5O_5S_2$:			
Calculated:	C, 48.33;	H, 4.23;	N, 12.25
Found:	C, 48.52;	H, 4.19;	N, 11.68

[0383] Example 149: R=2,2,2-trifluoroethylsulfonyl, R⁶=2-thienyl: Purified by trituration with diethyl ether; TLC: R_f=0.79, dichloromethane:tetrahydrofuran:acetic acid (90:10:1); NMR (DMSO/D₂O): 8.01 (s,1), 7.87 (dd,1, J=5.1, 0.9), 7.48 (dd,1, J=3.8, 0.9), 7.16 (dd,1, J=5.1, 3.8), 4.96 (d,1, J=16.8), 4.84 (d,1, J=16.8), 4.57 (q,2, J=9.8), 4.10 (d,1, J=2.9), 2.26 (m,1), 0.93 (d,3, J=6.8), 0.78 (d,3, J=6.8); MS: m/z=549(M+1).

Analysis for $C_{18}H_{18}F_6N_4O_5S_2$:			
Calculated:	C, 39.42;	H, 3.36;	N, 10.22
Found:	C, 39.30;	H, 3.36;	N, 10.20

[0384] Example 150: R=2,2,2-trifluoroethylaminosulfonyl, R⁶=2-thienyl: Purified by trituration with diethyl ether; TLC: R_f=0.56, dichloromethane:tetrahydrofuran:acetic acid (90:10:1); NMR (DMSO/D₂O): 7.95 (s,1), 7.84 (dd,1, J=1.0, 5.1), 7.42 (dd,1, J=1.0, 3.8), 7.15 (dd,1, J=3.8, 5.1), 4.94 (d,1, J=16.8), 4.82 (d,1, J=16.8), 4.10 (d,1, J=2.8), 3.70 (q,2, J=9.6), 2.28 (m,1), 0.94 (d,3, J=6.8), 0.79 (d,3, J=6.8); MS: m/z=564(M+1).

Analysis for $C_{18}H_{19}F_6N_5O_5S_2 \cdot 0.5 H_2O$:			
Calculated:	C, 37.76;	H, 3.52;	N, 12.23
Found:	C, 37.85;	H, 3.48;	N, 12.26

[0385] Example 151: R=cyanomethylsulfonyl, R⁶=2-thienyl: Purified by trituration with diethyl ether containing a trace of ethyl acetate; TLC: R_f=0.36, dichloromethane:methanol (90:10); NMR (DMSO/D₂O): 8.02 (s,1), 7.88 (dd,1, J=0.9, 5.1), 7.48 (dd,1, J=0.9, 3.8), 7.17 (dd,1, J=3.8, 5.1); 4.97 (d,1, J=16.9), 4.86 (d,1, J=16.9), 4.10 (d,1, J=2.8), 2.27 (m,1), 0.94 (d,3, J=6.8), 0.78 (d,3, J=6.8); MS: m/z=506(M+1).

Analysis for $C_{18}H_{18}F_3N_5O_6S$:			
Calculated:	C, 42.77;	H, 3.59;	N, 13.85
Found:	C, 42.63;	H, 3.59;	N, 13.90

[0386] Example 152: R=cyanomethylsulfonyl, R⁶=4-fluorophenyl: Purified by trituration with diethyl ether containing

a trace of ethyl acetate; TLC: R_f =0.33, dichloromethane:methanol (90:10); NMR(DMSO/ D_2O): 8.04 (s,1), 7.58 (m,2), 7.31 (m,2), 4.70 (d,1, J =16.8), 4.50 (d,1, J =16.8), 4.03 (broad s,1), 2.23 (m,1), 0.85 (d,3, J =6.4), 0.74 (d,3, J =6.4); MS: m/z =518($M+1$).

Analysis for $C_{20}H_{19}F_4N_5O_5S \cdot 1.0 H_2O$:

Calculated:	C, 44.86;	H, 3.95;	N, 13.08
Found:	C, 44.97;	H, 3.67;	N, 13.02

[0387] **Example 153:** R=2,2,2-trifluoroethylsulfonyl, R^6 =4-fluorophenyl: Purified by trituration with diethyl ether containing a trace of ethyl acetate; TLC: R_f =0.59, dichloromethane:methanol (90:10); NMR (DMSO/ D_2O): 7.99 (s,1), 7.56 (dd,2, J =5.4,5.4), 7.29 (t,2, J =8.8), 4.74 (d,1, J =18), 4.47 (d,1, J =18), 4.04 (d,1, J =2.6), 3.71 (q,2, J =9.4), 2.23 (m,1), 0.85 (d,3, J =6.8), 0.75 (d, J =6.8); MS: m/z =578($M+1$).

Analysis for $C_{20}H_{20}F_7N_5O_5S \cdot 0.5 H_2O$:

Calculated:	C, 41.10;	H, 3.62;	N, 11.98
Found:	C, 40.98;	H, 3.56;	N, 11.92

EXAMPLE 154

[0388]

2-[5-(2,6-Dimethylpyrid-4-ylmethoxycarbonylamino)-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0389] Using a similar procedure to that described in Example 1, 2-[5-(2,6-dimethylpyrid-4-ylmethoxycarbonylamino)-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was oxidized to afford the title compound, obtained as a yellow solid; TLC: R_f =0.4, methanol:dichloromethane (5:95); 300 MHz NMR (DMSO/ D_2O): 8.38 (s,1), 7.74 (d,1), 7.34 (d,1), 7.03 (m,3), 5.10 (s,2), 4.82 (m,2), 4.02 (d,1), 3.95 (m,2), 2.38 (s,6), 2.22 (m,1), 0.87 (d,3), 0.74 (d,3).

Analysis for $C_{25}H_{26}F_3N_5O_5S \cdot 0.25 H_2O$

Calculated:	C, 52.7;	H, 4.63;	N, 12.4
Found:	C, 52.7;	H, 4.66;	N, 12.2

[0390] The intermediate 2-[5-(2,6-dimethylpyrid-4-ylmethoxycarbonylamino)-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide was prepared as follows:

a. 2-[5-Benzoyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-pivaloyloxypropyl)acetamide.

[0391] To a solution of 2-[5-Benzoyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide (1.95 g), triethylamine (1.4 mL), and 4-dimethylaminopyridine (10 mg) in dichloromethane (35 mL) was added pivaloyl chloride (0.53 mL); and the resulting solution allowed to stir for 2 h. The solution was diluted with ethyl acetate, washed (saturated aqueous sodium bicarbonate (twice), saturated aqueous ammonium chloride, and brine), dried, and evaporated. Purification by chromatography, eluting with ethyl acetate:dichloromethane (10:90), afforded the ester (1.80 g) as a white foam; TLC: R_f =0.34, ethyl acetate:dichloromethane

(10:90); MS: $m/z=623(M+1)$.

b. 2-[5-Amino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-pivaloyloxypropyl)acetamide.

[0392] Using a procedure similar to that described in Example 12, 2-[5-benzyloxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-pivaloyloxypropyl)acetamide was deprotected to afford the amine, obtained as a yellow foam; TLC: $R_f=0.5$, ethyl acetate; MS: $m/z=489(M+1)$.

c. 2-[5-(2,6-Dimethylpyrid-4-ylmethoxycarbonylamino)-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyridinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-pivaloyloxypropyl)acetamide.

[0393] Using a procedure similar to that described in Example 3.a., but using 2,6-dimethylpyrid-4-ylcarbinol, 2-[5-amino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-pivaloyloxypropyl)acetamide was acylated to provide the urethane, obtained as yellow foam; TLC: $R_f=0.36$, ethyl acetate; MS: $m/z=652(M+1)$.

d. 2-[5-(2,6-Dimethylpyrid-4-ylmethoxycarbonylamino)-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-2-hydroxy-1-isopropylpropyl)acetamide.

To a solution of 2-[5-(2,6-dimethylpyrid-4-ylmethoxycarbonylamino)-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyridinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-pivaloyloxypropyl)acetamide (1.44 g) in tetrahydrofuran (10 mL) and methanol (10 mL) was added a solution of lithium hydroxide monohydrate (0.92 g) in water (20 mL). The resulting solution was stirred for 2 h, diluted with saturated ammonium chloride (30 mL), and extracted with ethyl acetate. The ethyl acetate solution was dried and evaporated. The resulting material was chromatographed, eluting with ethyl acetate, to provide the alcohol (0.39 g) as a yellow foam; TLC: $R_f=0.24$, ethyl acetate; MS: $m/z=568(M+1)$.

EXAMPLE 155

2-[5-Ethylamino-2-(4-methoxyphenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0395] Using a procedure similar to that described in Example 80, the title compound was prepared from the product of Example 28. Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 99.3:0.7), then triturated with hexane:diethyl ether (80:20); TLC: $R_f=0.41$, dichloromethane:acetone (70:30); NMR: 0.73-1.03 (m,6), 1.17 (t,3), 2.07-2.27 (m,1) 3.03-3.20 (m,2), 3.79 (s,3) 4.03-4.73 (m,3), 5.30-5.43 (m,1), 6.96 (d,2), 7.10 (s,1), 7.30-7.47 (m,2), 8.80 (d,1); MS: $m/z=455(M+1)$.

Analysis for $C_{21}H_{25}F_3N_4O_4 \cdot 0.2 H_2O$:			
Calculated:	C, 55.06;	H, 5.58;	N, 12.23
Found:	C, 55.43;	H, 5.64;	N, 11.80

EXAMPLES 156-159

[0396] Using a procedure similar to that described in Example 89, the following compounds of formula I wherein R^0 is isopropyl, R is formyl and R^6 is the indicated group were prepared from the corresponding compounds of formula I wherein R is hydrogen.

[0397] Example 156: $R^6=4$ -methoxyphenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 97.5:2.5); TLC: $R_f=0.46$, dichloromethane:acetone (70:30); NMR: 0.73-1.00 (m,6), 2.03-2.30 (m,1), 3.81 (s,3), 4.02-4.77 (m,3), 6.87-7.07 (m,2), 7.43 (d,2), 8.36 (s,1), 8.80-8.93 (m,2), 10.00 (s,1); MS: $m/z=455(M+1)$.

Analysis for $C_{20}H_{21}F_3N_4O_5 \cdot 0.3 H_2O$:

Calculated:	C, 52.24;	H, 4.73;	N, 12.18
Found:	C, 52.24;	H, 4.73;	N, 12.07

[0398] Example 157: $R^6=2$ -thienyl: Chromatography solvent: dichloromethane:methanol (gradient 99.5:0.5, 97.5:2.5); TLC: $R_f=0.55$, dichloromethane:acetone (70:30); NMR: 0.73-1.17 (m,6), 2.13-2.33 (m,1), 4.07-5.07 (m,3), 7.10-7.23 (m,1), 7.30-7.47 (m,1), 7.77-7.90 (m,1), 8.36 (s,1), 8.88 (d,1), 9.05 (d,1), 10.07 (s,1); MS: $m/z=431(M+1)$.

Analysis for $C_{17}H_{17}F_3N_4O_4S \cdot 0.3 H_2O$:

Calculated:	C, 46.85;	H, 4.07;	N, 12.85
Found:	C, 46.83;	H, 4.05;	N, 12.62

[0399] Example 158: $R^6=4$ -fluorophenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 97.5:2.5), then re-chromatographed with dichloromethane:acetone (80:20); TLC: $R_f=0.47$, dichloromethane:acetone (70:30); NMR: 0.60-1.03 (m,6), 2.03-2.27 (m,1), 4.00-4.77 (m,3), 7.27-7.43 (m,2), 7.50-7.70 (m,2), 8.37 (s,1), 9.70-9.90 (m,2), 10.07 (s,1); MS: $m/z=443(M+1)$.

Analysis for $C_{19}H_{18}F_4N_4O_4 \cdot 0.5 H_2O$:

Calculated:	C, 50.55;	H, 4.24;	N, 12.41
Found:	C, 50.52;	H, 4.31;	N, 12.20

[0400] Example 159: $R^6=$ phenyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 97.5:2.5); TLC: $R_f=0.51$, dichloromethane:acetone (70:30); NMR: 0.73-1.00 (m,6), 2.07-2.30 (m,1), 4.07-4.77 (m,3), 7.37-7.60 (m,5), 8.37 (s,1), 8.85 (d,2), 10.07 (s,1); MS: $m/z=425(M+1)$.

Analysis for $C_{19}H_{18}F_3N_4O_4$:

Calculated:	C, 53.77;	H, 4.51;	N, 13.20
Found:	C, 53.43;	H, 4.63;	N, 12.94

EXAMPLES 160-161

[0401] Using a procedure similar to that described in Example 90 but using 4-nitrobenzylaminosulfonyl chloride, the following compounds of formula I wherein R^0 is isopropyl, R is 4-nitrobenzylaminosulfonyl and R^6 is the indicated group were prepared from the corresponding compounds of formula I wherein R is hydrogen.

[0402] Example 160: $R^6=2$ -thienyl: Chromatography solvent: dichloromethane:methanol (gradient, 99.5:0.5, 98:2); TLC: $R_f=0.36$, dichloromethane:methanol (98:2); NMR: 0.77-1.13 (m,6), 2.17-2.30 (m,1), 4.26 (d,2), 4.70-5.00 (m,3), 7.13-7.20 (m,1), 7.32 (d,1), 7.54 (d,2), 7.86 (d,1), 7.92 (s,1), 8.14 (d,2), 8.27-8.37 (m,1), 9.03 (d,1), 9.24 (s,1); MS: $m/z=617(M+1)$.

Analysis for $C_{23}H_{23}F_3N_6O_7S_2$:			
Calculated:	C, 44.87;	H, 3.60;	N, 13.65
Found:	C, 44.63;	H, 3.82;	N, 13.70

[0403] **Example 161:** R^6 =phenyl: Chromatography solvent: dichloromethane:methanol (gradient 99.5:0.5, 98:2); TLC: R_f =0.35, dichloromethane:methanol (98:2); NMR: 0.70-0.93 (m,6), 2.10-2.30 (m,1), 4.27 (d,2), 4.03-4.70 (m,3), 7.43-7.63 (m,7), 7.95 (s,1), 8.17 (d,2), 8.33-8.43 (m,1), 8.84 (d,1), 9.24 (s,1); MS: m/z =611(M+1).

Analysis for $C_{25}H_{25}F_3N_6O_7S \cdot 0.3 H_2O$:			
Calculated:	C, 48.74;	H, 4.18;	N, 13.64
Found:	C, 48.69;	H, 4.11;	N, 13.60

[0404] The intermediate 4-nitrobenzylaminosulfonyl chloride was prepared from 4-nitrobenzylamine hydrochloride and sulfonyl chloride at reflux in acetonitrile. The solvent was evaporated and the crude material used without further purification.

EXAMPLE 162

2-[5-(4-Acetylamino benzylaminosulfonylamino)-6-oxo-2-thienyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0405] To a solution of 2-[5-(4-nitrobenzylaminosulfonylamino)-6-oxo-2-thienyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (0.431 g) in acetic acid (12 mL) was added acetic anhydride (0.265 mL), followed by iron powder (0.783 g). The reaction was stirred at room temperature overnight. Methanol was added and the reaction mixture filtered through diatomaceous earth to remove excess iron. Ethyl acetate was added and the resultant solution washed (water (twice), brine), dried ($MgSO_4$) and evaporated. The crude material was purified by chromatography, with dichloromethane:methanol (gradient, 99.5:0.5, 98:2) as the eluent, to provide the title compound (0.247 g); TLC: R_f =0.27, dichloromethane:methanol (98:2); NMR: 0.87-0.97 (m,6), 2.17-2.30 (m,1), 3.32 (s,3), 4.02 (d,2), 4.73-4.77 (m,1), 4.77-4.93 (m,2), 7.13-7.20 (m,3), 7.33 (d,1), 7.47 (d,2), 7.80-7.87 (m,1), 7.91 (s,1), 7.97-8.03 (m,1), 9.04 (d,1), 9.09 (s,1), 9.87 (s,1); MS: m/z =629(M+1).

Analysis for $C_{25}H_{27}F_3N_6O_6S_2 \cdot 0.4 H_2O$			
Calculated:	C, 47.22;	H, 4.40;	N, 13.21
Found:	C, 47.17;	H, 4.26;	N, 13.11

EXAMPLE 163

2-[2-(4-Acetylamino phenyl)-5-benzoyloxycarbonylamino-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0406] 2-[5-Benzoyloxycarbonylamino-2-(4-nitrophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)-acetamide (2.01 g) was reductively acetylated using a similar procedure to that described in Example 162. The crude material was purified by chromatography, with dichloromethane:methanol (gradient 99.5:0.5, 98:2) as the eluent, to provide the title compound (1.51 g) as a white solid; TLC: R_f =0.47, dichloromethane:acetone (70:30); NMR: 0.77-1.00 (m,6), 2.00-2.33 (m,1), 2.07 (s,3), 4.03-4.73 (m,3), 5.17 (s,2), 7.30-7.53 (m,7), 7.60-7.77 (m,2), 8.43

(s,1), 9.83-9.03 (m,2), 10.19 (d,1); MS: $m/z=588(M+1)$.

Analysis for $C_{28}H_{28}F_3N_5O_6 \cdot 0.7 H_2O$			
Calculated:	C, 56.03;	H, 4.93;	N, 11.66
Found:	C, 56.00;	H, 5.03;	N, 11.45

EXAMPLE 164

2-[5-Benzyloxycarbonylamino-2-(4-formylaminophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

[0407] To a solution of 2-[5-benzyloxycarbonylamino-2-(4-nitrophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide (2.02 g) in 90% aqueous formic acid (56 mL) was added iron powder (3.9 g). The mixture was stirred at room temperature overnight. Methanol was added and the reaction mixture filtered through diatomaceous earth to remove the excess iron. Ethyl acetate was added and the resultant solution washed with water (twice), brine, then dried ($MgSO_4$) and evaporated. The crude material was purified by chromatography, with dichloromethane:methanol (gradient, 99.5:0.5, 98.2) as the eluent, followed by trituration with hexane:diethyl ether (80:20), to provide the title compound (1.12 g) as an off-white solid; TLC: $R_f=0.39$, dichloromethane:acetone (70:30); NMR: 0.67-1.03 (m,6), 2.03-2.33 (m,1), 4.00-4.80 (m,3), 5.17 (s,2), 7.20-7.60 (m,7), 7.57-7.77 (m,2), 8.22 (s,1), 8.43 (s,1), 8.83-9.07 (m,2), 10.30-10.57 (m,1); MS: $m/z=574(M+1)$.

Analysis for $C_{27}H_{26}F_3N_5O_6 \cdot 0.6 H_2O$			
Calculated:	C, 55.49;	H, 4.69;	N, 11.98
Found:	C, 55.64;	H, 4.86;	N, 11.59

EXAMPLES 165-167

[0408] Using a procedure similar to that described in Example 12, the following compounds of formula I wherein R^0 is isopropyl, R is hydrogen, and R^6 is the indicated group were prepared from the corresponding compounds of formula I wherein R is benzyloxycarbonyl.

[0409] Example 165: $R^6=4$ -acetylaminophenyl: After the addition of sodium bicarbonate the product precipitated and was collected by filtration. The crude product was purified by trituration with hexane:diethyl ether (70:30) followed by chromatography, eluting with dichloromethane:methanol (gradient 95:5, 90:10); TLC: $R_f=0.06$, dichloromethane:methanol (98:2); NMR ($DMSO/D_2O$): 0.81 (d,3), 0.88 (d,3), 2.17-2.30 (m,1), 4.08 (s,1), 4.40-4.67 (m,2), 7.37 (d,3), 7.60 (d,2); MS: $m/z=454(M+1)$.

Analysis for $C_{20}H_{22}F_3N_5O_4 \cdot 0.7 H_2O \cdot 0.4 CH_3OH$			
Calculated:	C, 51.16;	H, 5.26;	N, 14.62
Found:	C, 51.45;	H, 5.31;	N, 14.19

[0410] Example 166: $R^6=4$ -formylamino: After the addition of sodium bicarbonate the product precipitated and was collected by filtration. The crude product was purified by trituration with hexane:diethyl ether (70:30) followed by chromatography, eluting with dichloromethane:methanol (gradient 95:5, 90:10); TLC: $R_f=0.05$, dichloromethane:methanol (95:5); NMR ($DMSO/D_2O$): 0.77 (d,3), 0.87 (d,3), 2.17-2.30 (m,1), 4.03-4.10 (m,1), 4.40-4.67 (m,2), 7.37-7.47 (m,3), 7.61 (d,2), 8.30 (s,1); MS: $m/z=440(M+1)$.

Analysis for $C_{19}H_{20}F_3N_5O_4 \cdot 0.6 H_2O \cdot 0.4 CH_3OH$			
Calculated:	C, 50.32;	H, 4.96;	N, 15.12
Found:	C, 50.31;	H, 4.93;	N, 15.15

[0411] Example 167: $R^6=4$ -trifluoroacetylaminophenyl: After addition of sodium bicarbonate the product precipitated and was collected by filtration. The crude product was purified by trituration with hexane:diethyl ether (70:30) followed by chromatography, eluting with dichloromethane:methanol (gradient, 95:5, 90:10); TLC: $R_f=0.10$, dichloromethane:methanol (95:5); NMR (DMSO/ D_2O): 0.70-1.03 (m,6), 2.17-2.33 (m,1), 4.00-4.13 (m,1), 4.37-4.70 (m,2), 7.37 (s,1), 7.48 (d,2), 7.72 (d,2); MS: $m/z=508(M+1)$.

Analysis for $C_{20}H_{19}F_8N_5O_4 \cdot 0.5 H_2O$			
Calculated:	C, 46.51;	H, 3.90;	N, 13.56
Found:	C, 46.62;	H, 3.93;	N, 13.29

[0412] The intermediate 2-[5-benzoyloxycarbonylamino-6-oxo-2-(4-trifluoroacetylaminophenyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide, which is also an Example of the invention, was prepared from 2-[5-benzoyloxycarbonylamino-2-(4-nitrophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide using a procedure similar to that described in Example 165 but employing trifluoroacetic acid in place of acetic acid and trifluoroacetic anhydride in place of acetic anhydride.

FORMULAE

5

10

15

20

25

30




35

40


45

50

55

**I****II****Vb**

Scheme I



Scheme II

5

10

15

20

25

30

35

40


45


50

55

Claims

1. A compound of formula I





I

wherein:

R^0 is (1-5C)alkyl;

R is hydrogen; or

R is an acyl group of formula A.X.CO- in which A.X-, taken together, is hydrogen, trifluoromethyl, 2,2,2-trifluoroethoxy, amino, methoxyamino, 2,2,2-trifluoroethylamino, RbRcN.O-, RaOCONH-, R^1SO_2NH -, RaOCO-, RbRcNCO- or RaCO-; or

R is an acyl group of formula A.X.CJ- in which

J is oxygen or sulfur;

X is a direct bond, imino, oxy or thio; and

A is as defined below or

A is tetrahydropyran-4-yl, 1-methylpiperid-4-yl, or 5-methyl-1,3-dioxacyclohex-5-ylmethyl; or

R is a sulfonyl group of formula D.W.SO₂- in which D.W-, taken together, is hydroxy, amino, di(1-4C)alkylamino, 2,2,2-trifluoroethylamino, 2,2,2-trifluoroethyl, 3,3,3-trifluoropropyl or trifluoromethyl; or

W is a direct bond, imino, carbonylimino, oxycarbonylimino or iminocarbonylimino; and

D is as defined below; or

R is an alkyl, aryl or heteroaryl group G as defined below;

The group A, D or G is (1-6C)alkyl, (3-6C)cycloalkyl, (3-6C)cycloalkyl-(1-3C)alkyl, aryl, aryl(1-3C)alkyl, heteroaryl or heteroaryl(1-3C)alkyl wherein an aryl moiety is selected from phenyl, indenyl and naphthyl, and a heteroaryl moiety is selected from furyl, imidazolyl, tetrazolyl, pyridyl, pyridyl N-oxide, thienyl, pyrimidinyl, pyrimidinyl N-oxide, indolyl, quinolyl and quinolyl N-oxide, and wherein an aryl or heteroaryl moiety may bear one or more halogeno, nitro, methyl or trifluoromethyl groups and further wherein the group A, D or G may bear one or more substituents selected from a group consisting of hydroxy, (1-4C)alkoxy, acyloxy containing 1 to 5 carbon atoms, COORa, CONRbRc, COO(CH₂)₂NR¹R², cyano, SO₂R¹, CONR³SO₂R¹, NR⁴R⁵, NR⁶CHO, NR⁷COR⁸, NR⁹COOR¹⁰, NR¹¹CQR¹²R¹³, NR¹⁴SO₂R¹⁵, SO₂NR¹⁶R¹⁷, SO₂NR¹⁸COR¹⁹ and P(O)(ORa)₂ in which

Q is oxygen or sulfur;

Ra-Rn are independently hydrogen, benzyl or (1-4C)alkyl; or, independently, a group NRbRc, NR¹R², NR³R⁴ or NR⁵R⁶ is a cyclic radical selected from a group consisting of 1-pyrrolidinyl, piperidino, morpholino or 1-piperazinyl which may bear a (1-4C)alkyl substituent at the 4-position; or, independently, a group NR⁷R⁸ is a cyclic radical selected from a group consisting of 2-pyrrolidinon-1-yl, succinimido, oxazolidin-2-on-3-yl, 2-benzoxazolinon-3-yl, phthalimido and cis-hexahydrophthalimido; and

R¹-R⁴ are independently trifluoromethyl, (1-6C)alkyl, (3-6C)cycloalkyl, aryl or heteroaryl, wherein an aryl moiety is selected from phenyl, indenyl and naphthyl, and a heteroaryl moiety is selected from furyl, imidazolyl, tetrazolyl, pyridyl, pyridyl N-oxide, thienyl, pyrimidinyl, pyrimidinyl N-oxide, indolyl, quinolyl and quinolyl N-oxide, and wherein an aryl or heteroaryl moiety may bear one or more substituents selected from a group consisting of (1-4C)alkyl, hydroxy, (1-4C)alkoxy, halogeno and trifluoromethyl;

R⁶ is (1-5C)alkyl which has no tertiary carbon, (3-7C)cycloalkyl, aryl or heteroaryl wherein an aryl moiety is selected from phenyl, indenyl and naphthyl, and a heteroaryl moiety is selected from furyl, imidazolyl, tetrazolyl, pyridyl, pyridyl N-oxide, thienyl, pyrimidinyl, pyrimidinyl N-oxide, indolyl, quinolyl and quinolyl N-oxide, and wherein an aryl or heteroaryl moiety independently may bear one or more of the substituents defined for the group A or an aryl or heteroaryl moiety thereof; and

provided that no aliphatic carbon is bonded to more than one nitrogen or oxygen, except as part of a cyclic ketal or where the nitrogen bears a carbonyl group; or,

for a compound of formula I which is acidic or basic, a pharmaceutically acceptable salt thereof.

2. A compound as claimed in Claim 1 wherein:

R⁰ is methyl, ethyl, propyl, isopropyl or isobutyl; R is hydrogen, A.X.CO-, A.X.CJ-, D.W.SO₂- or G in which W is a direct bond or imino;

G is (1-3C)alkyl, aryl(1-3C)alkyl or heteroaryl(1-2C)alkyl which may bear one or more substituents as defined in Claim 1 for G or a part thereof;

(1-6C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, pentyl, 3-methylbutyl, 1-ethylpropyl, hexyl or 4-methylpentyl; (3-6C)cycloalkyl is cyclopropyl, cyclopentyl or cyclohexyl; the (1-3C)alkyl portion of (3-6C)cycloalkyl(1-3C)alkyl, aryl(1-3C)alkyl or heteroaryl(1-3C)alkyl is methylene, ethylene or trimethylene; C(1-4)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl or t-butyl; acyloxy containing 1 to 5 carbon atoms is acetoxo; (1-4C)alkoxy is methoxy, ethoxy, propoxy, isopropoxy or t-butoxy; halogeno is bromo, chloro or fluoro; A.X-, taken together, is 2,2,2-trifluoroethoxy; COORa is carboxy or methoxycarbonyl; CONRbRc is carbamoyl or N,N-dimethylcarbamoyl; NRgCOR² is trifluoroacetyl amino; CONRdSO₂R¹ is N-phenylsulfonylcarbamoyl or N-(4-chlorophenylsulfonyl)carbamoyl; A.X- is tris(hydroxymethyl)methylamino, tris(acetoxymethyl)methylamino or 2,2-bis(hydroxymethyl)propoxy; D.W-, taken together, is 2,2,2-trifluoroethylamino or 3,3,3-trifluoropropyl; and

R⁶ is isopropyl, cyclopentyl, cyclohexyl, phenyl, furyl, thienyl or pyridyl in which a phenyl or heteroaryl may bear one or two substituents as defined in Claim 1.

3. A compound as claimed in Claim 1 or 2 wherein R⁰ is isopropyl; R is hydrogen, A.X.CO-, A.X.CJ-, D.W.SO₂- or G in which A.X-, taken together, is 2,2,2-trifluoroethoxy; J is oxygen; X is a direct bond, imino or oxy; A is methyl, ethyl, phenyl, benzyl, phenethyl, pyridyl, thienyl, 5-tetrazolyl, thiazolyl, pyridylmethyl, thenyl, 5-tetrazolylmethyl, 2-(pyridyl)ethyl, 2-(thienyl)ethyl or 2-(thiazolyl)ethyl wherein the phenyl or heteroaryl group may bear one or two halogeno or methyl groups and further wherein the group A may bear a substituent selected from hydroxy, methoxy, t-butoxy, acetoxo, pivaloyloxy, carboxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl, dimethylcarbamoyl, 2-(dimethylamino)ethoxycarbonyl, cyano, methylsulfonyl, phenylsulfonyl, N-methylsulfonylcarbamoyl, N-phenylsulfonylcarbamoyl, amino, dimethylamino, oxazolidin-2-on-3-yl, acetyl amino, trifluoroacetyl amino, ureido, methylsulfonyl, sulfamoyl, dimethylphosphoryl or diethylphosphoryl; D.W-, taken together, is 2,2,2-trifluoroethylamino or 3,3,3-trifluoropropyl; D is methyl, ethyl, isopropyl, tert-butyl, cyclohexyl, phenyl, benzyl, phenethyl, pyridyl, thienyl, 5-tetrazolyl, thiazolyl, quinolinyl, pyridylmethyl, thenyl, 5-tetrazolylmethyl, 2-(pyridyl)ethyl, 2-(thienyl)ethyl or 2-(thiazolyl)ethyl wherein the phenyl or heteroaryl group may bear one or two halogeno or methyl groups and further wherein the group D may bear a substituent selected from hydroxy, methoxy, t-butoxy, acetoxo, pivaloyloxy, carboxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl, dimethylcarbamoyl, 2-(dimethylamino)ethoxycarbonyl, cyano, methylsulfonyl, phenylsulfonyl, N-methylsulfonylcarbamoyl, N-phenylsulfonylcarbamoyl, N-(4-chlorophenylsulfonyl)carbamoyl, methylsulfonylamino, amino, dimethylamino, oxazolidin-2-on-3-yl, acetyl amino, trifluoroacetyl amino, ureido, methylsulfonyl, sulfamoyl, dimethylphosphoryl or diethylphosphoryl; G is methyl, ethyl, benzyl, phenethyl, pyridyl, pyridylmethyl, thenyl, 5-tetrazolylmethyl, or 2-(pyridyl)ethyl, wherein an alkyl carbon may bear an oxo group and wherein the phenyl or heteroaryl group may bear one or two halogeno or methyl groups and further wherein the group G may bear a substituent selected from hydroxy, methoxy, acetoxo, carboxy, methoxycarbonyl, ethoxycarbonyl, carbamoyl, dimethylcarbamoyl, phenylcarbamoyl, pyridylcarbamoyl, methylsulfonylamino, amino, dimethylamino, acetyl amino, nicotinoylamino, or trifluoroacetyl amino.

4. A compound as claimed in Claim 1, 2 or 3 wherein R is hydrogen, formyl, trifluoroacetyl, 2,2,2-trifluoroethoxycarbonyl, hydroxyoxalyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, 2-methoxyethoxycarbonyl, 4-fluorophenoxycarbonyl, 4-bromophenoxycarbonyl, 4-methoxyphenoxycarbonyl, benzoyloxycarbonyl, 4-fluorobenzoyloxycarbonyl, 4-pyridylmethoxycarbonyl, 3-methylpyrid-4-ylmethoxycarbonyl, 2, 6-dimethylpyrid-4-ylmethoxycarbonyl, 2-pyridylmethoxycarbonyl, 6-methylpyrid-2-ylmethoxycarbonyl, 2-dimethylaminoethoxycarbonyl, acetyl, carbamoyl/methylaminocarbonyl, 4-(N-phenylsulfonylcarbamoyl)phenylacetyl, methylthiocarbonyl, sulfo, aminosulfonyl, dimethylaminosulfonyl, 2,2,2-trifluoroethylaminosulfonyl, 3,3,3-trifluoroethylsulfonyl, trifluoromethylsulfonyl, methylsulfonyl (which may bear a methoxycarbonyl, carboxy or ethylsulfonyl substituent), methylaminosulfonyl, isopropylaminosulfonyl, butylsulfonyl, butylaminosulfonyl, tert-butylaminosulfonyl, cyclohexylaminosulfonyl, phenylsulfonyl (in which the phenyl may bear a chloro, nitro, amino, formylamino, acetyl amino, trifluoroacetyl amino, methoxy, carboxy, N-(4-chlorophenylsulfonyl)carbamoyl, or methylsulfonylamino substituent at the 3- or 4-position), anilino, pyridylsulfonyl, quinolinylsulfonyl, benzylsulfonyl (in which the phenyl ring may bear a nitro or amino substituent at the 3- or 4-position), pyridyl/methylsulfonyl, 2-(pyridyl)ethylsulfonyl, benzylaminosulfonyl, methyl, ethyl, benzyl, phenethyl or pyridylmethyl.

5. A compound as claimed in Claim 4 wherein R is hydrogen, formyl, trifluoroacetyl, 2,2,2-trifluoroethoxycarbonyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, 2-methoxyethoxycarbonyl, 4-bromophenoxycarbonyl, benzoyloxycarbonyl, 2,6-dimethylpyrid-4-ylmethoxycarbonyl, methylthiocarbonyl, tert-butylaminosulfonyl, 4-acetylami-

nophenylsulfonyl, 4-[N-(4-chlorophenylsulfonyl)carbamoyl]phenylsulfonyl, benzylsulfonyl, benzylaminosulfonyl or ethyl.

6. A compound as claimed in any one of Claims 1-5 wherein R^6 is 2-furyl, 2-thienyl, 3-pyridyl or phenyl in which the phenyl may bear one or two halogeno, trifluoromethyl, methyl, hydroxy, methoxy, tert-butoxy, methoxycarbonyl or carboxy substituents.

7. A compound as claimed in Claim 1 in which R^0 is isopropyl, R is hydrogen, formyl, 2,2,2-trifluoroethoxycarbonyl, isopropoxycarbonyl, methylthiocarbonyl or ethyl, and R^6 is 2-furyl, 2-thienyl, 3-pyridyl or phenyl in which the phenyl may bear one or two halogeno, trifluoromethyl, methyl, hydroxy, methoxy, tert-butoxy, methoxycarbonyl or carboxy substituents.

8. A compound as claimed in Claim 1 selected from:

(a) 2-[5-amino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(b) 2-[5-amino-2-(4-fluorophenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(c) 2-[5-isopropoxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(d) 2-[5-ethylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(e) 2-[6-oxo-2-phenyl-5-(2,2,2-trifluoroethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(f) 2-[5-methylthiocarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(g) 2-[2-(4-fluorophenyl)-5-methylthiocarbonylamino-6-oxo-1,2-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(h) 2-[6-oxo-2-(2-thienyl)-5-(2,2,2-trifluoroethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide,

(i) 2-[5-formylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide, and


(j) 2-[5-formylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acetamide.

9. A salt as claimed in Claim 1 selected from

(a) for an acidic compound of formula I, an alkali metal salt, an alkaline earth metal salt, an aluminum salt, an ammonium salt, or a salt made from an organic base which affords a pharmaceutically acceptable cation; and
(b) for a basic compound of formula I, an acid-addition salt made with an acid which provides a pharmaceutically acceptable anion.

10. A method of making a compound of formula I, or a pharmaceutically acceptable salt thereof, as claimed in any one of Claims 1-9 which is characterized by:

(A) Oxidizing a corresponding alcohol of formula II;



II

(B) For a compound of formula I which contains an N-H residue, removal by using a conventional method of the nitrogen protecting group of a corresponding compound bearing a conventional nitrogen protecting group;

(C) For a compound of formula I wherein R is an acyl group, acylation of a corresponding amine of formula I wherein R is hydrogen;

(D) For a compound of formula I wherein R is a sulfonyl group, sulfonylation of a corresponding amine of formula I wherein R is hydrogen with a corresponding sulfonic acid of formula D.W.SO₂.OH, or an activated derivative thereof;

(E) For a compound of formula I in which R is a group G, substitution of the group L of a corresponding compound of formula G-L, wherein L is a conventional leaving group, with a corresponding amine of formula I wherein R is hydrogen, optionally using a conventional catalyst;

(F) For a compound of formula I which bears a hydroxy substituent on an aryl or heteroaryl group, cleaving the alkyl ether or acyloxy ester of a corresponding compound of formula I which bears a (1-4C)-alkoxy or acyloxy containing 1 to 5 carbon atoms substituent on an aryl or heteroaryl group;

(G) For a compound of formula I which bears a group of formula COORa in which Ra is hydrogen, decomposing the ester group of a corresponding ester made with a conveniently removed acid protecting group;

(H) For a compound of formula I bearing a moiety of formula COORa, CONRbRc, COO(CH₂)₂NReRf or CONRdSO₂R¹, acylation of a corresponding compound of formula HORa, HNRbRc, HO(CH₂)₂NReRf or HNRdSO₂R¹ with a corresponding acid of formula I bearing a moiety of formula COORa in which Ra is hydrogen, or an activated derivative thereof;


(I) For a compound of formula I bearing a containing 1 to 5 carbon atoms acyloxy group or a group of formula NRgCHO, NRgCOR², NRgCOOR², NRhCQNRiRj or NRkSO₂R³, acylation or sulfonylation of a corresponding compound of formula I bearing a hydroxy group or an amino group of formula NHRg, NHRh or NHRk with an activated derivative of a corresponding acid of formula HOCHO, HOCOR², HOCOR², HOCQNRiRj (including an isocyanate or isothiocyanate) or HOSO₂R³, respectively, using a conventional method;

(J) For a compound of formula I which bears a heteroaryl N-oxide group, oxidation of a corresponding compound of formula I which bears a heteroaryl group using a conventional oxidant; or

(K) For a compound of formula I which bears a primary amino group, reduction of a corresponding compound bearing a nitro group using a conventional reducing method; and

whereafter, for any of the above procedures, when a pharmaceutically acceptable salt of an acidic or basic compound of formula I is required, reacting the acidic or basic form of such a compound of formula I with a base or acid affording a physiologically acceptable counterion or by any other conventional procedure; and wherein R, R⁰, R⁶, D, W, G, Ra-Rk, R¹-R³ and Q, except where more particularly described, have the meaning defined in any one of Claims 1-9.


11. A compound of formula II,



II

wherein R, R⁰ and R⁶ are defined as in Claim 1, or a salt thereof.

12. A compound of formula Vb,




Vb

wherein R has a value defined for G in Claim 1, R⁰ and R⁶ are defined as in Claim 1, and Rx is a group which protects and activates a primary amino group for substitution, or a salt thereof.

13. A pharmaceutical composition comprising a compound as defined in Claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable diluent or carrier.

Patentansprüche

1. Verbindung der Formel I



I

wobei:

R⁰ für einen (1-5C)-Alkylrest steht;

R ein Wasserstoffatom ist; oder

R ein Acylrest der Formel A.X.CO- ist, worin A.X- zusammengekommen ein Wasserstoffatom, eine Trifluormethyl-, 2,2,2-Trifluorethoxy-, Amino-, Methoxyamino- oder 2,2,2-Trifluorethylaminogruppe oder ein Rest RbRcN.O-, RaOCONH-, R¹SO₂NH-, RaOCO-, RbRcNCO- oder RaCO- ist; oder

R ein Acylrest der Formel A.X.CJ- ist, worin

J ein Sauerstoffatom oder Schwefelatom ist;

X eine direkte Bindung, eine Imino-, Oxy- oder Thiogruppe ist; und

A die nachstehend definierte Bedeutung hat; oder

A eine Tetrahydropyran-4-yl-, 1-Methylpiperid-4-yl- oder 5-Methyl-1,3-dioxacyclohex-5-ylmethylgruppe ist; oder

R eine Sulfonylgruppe der Formel D.W.SO₂- ist, worin D.W- zusammengekommen eine Hydroxy-, Amino-, Di(1-4C)-Alkylamino-, 2,2,2-Trifluorethylamino-, 2,2,2-Trifluorethyl-, 3,3,3-Trifluorpropyl- oder Trifluormethylgruppe ist; oder

W eine direkte Bindung, eine Imino-, Carbonylimino-, Oxycarbonylimino- oder Iminocarbonyliminogruppe ist; und

D die nachstehend definierte Bedeutung hat; oder

R ein Alkyl-, Aryl- oder Heteroarylrest G wie nachstehend definiert ist; der Rest A, D oder G ein (1-6C)Alkyl-, (3-6C)Cycloalkyl-, (3-6C)Cycloalkyl-(1-3C)alkyl-, Aryl-, Aryl(1-3C)alkyl-, Heteroaryl- oder Heteroaryl(1-3C)alkylrest ist, wobei eine Aryleinheit ausgewählt ist aus einer Phenyl-, Indenyl- und Naphthylgruppe, und eine Heteroaryleinheit ausgewählt ist aus einer Furyl-, Imidazolyl-, Tetrazolyl-, Pyridyl-, Pyridyl-N-oxid-, Thienyl-, Pyrimidinyl-, Pyrimidinyl-N-oxid-, Indolyl-, Chinolyl- und Chinolyl-N-oxidgruppe und wobei eine Aryl- oder Heteroaryleinheit ein oder mehrere Halogenatome, Nitro-, Methyl- oder Trifluormethylgruppen tragen kann, und wobei außerdem der Rest A, D oder G einen oder mehrere Substituenten tragen kann, die aus einer aus

Hydroxy, (1-4C)Alkoxy, Acyloxy mit 1 bis 5 Kohlenstoffatomen COORa, CONRbRc, COO(CH₂)₂NReRf Cyano, SO₂R¹, CONRdSO₂R¹, NReRf, NRgCHO, NRgCOR², NRgCOOR², NRhCQRiRj, NRkSO₂R³, SO₂NRlRm, SO₂NRnCOR⁴ und P(O)(ORa)₂ bestehenden Gruppe ausgewählt sind, wobei

Q ein Sauerstoffatom oder Schwefelatom ist;

5 Ra-Rn unabhängig voneinander ein Wasserstoffatom, ein Benzyl- oder (1-4C)Alkylrest sind; oder ein Rest NRbRc, NReRf, NRlRj oder NRlRm unabhängig ein cyclischer Rest ist, der aus einer aus 1-Pyrrolidinyl, Piperidino, Morpholino oder 1-Piperazinyl, das einen (1-4C)Alkylsubstituenten an der Stellung 4 tragen kann, bestehenden Gruppe ausgewählt ist; oder ein Rest NReRf unabhängig ein cyclischer Rest ist, der aus einer aus 2-Pyrrolidinon-1-yl Succinimido, Oxazolidin-2-on-3-yl, 2-Benzoxazolinon-3-yl, Phthalimido und cis-Hexahydrophthalimido bestehenden Gruppe ausgewählt ist; und

10 R¹-R⁴ unabhängig eine Trifluormethylgruppe, ein (1-6C)Alkyl-, (3-6C)Cycloalkyl-, Aryl- oder Heteroarylrest sind, wobei eine Aryleinheit ausgewählt ist aus einer Phenyl-, Indenyl- und Naphthylgruppe, und eine Heteroaryleinheit ausgewählt ist aus einer Furyl-, Imidazolyl-, Tetrazolyl-, Pyridyl-, Pyridyl-N-oxid-, Thienyl-, Pyrimidinyl-, Pyrimidinyl-N-oxid-, Indolyl-, Chinolyl- und Chinolyl-N-oxidgruppe, und wobei eine Aryl- oder Heteroaryleinheit einen oder mehr Substituenten tragen kann, der aus einer aus (1-4C)Alkyl, Hydroxy, (1-4C)Alkoxy, Halogen und Trifluormethyl bestehenden Gruppe ausgewählt ist;

15 R⁵ ein (1-5C)Alkylrest ohne tertiäres Kohlenstoffatom, (3-7C)Cycloalkyl-, Aryl- oder Heteroarylrest ist, wohn eine Aryleinheit ausgewählt ist aus einer Phenyl-, Indenyl- und Naphthylgruppe, und eine Heteroaryleinheit ausgewählt ist aus einer Furyl-, Imidazolyl-, Tetrazolyl-, Pyridyl-, Pyridyl-N-oxid-, Thienyl-, Pyrimidinyl-, Pyrimidinyl-N-oxid-, Indolyl-, Chinolyl- und Chinolyl-N-oxidgruppe und wobei eine Aryl- oder Heteroaryleinheit unabhängig einen oder mehrere der für den Rest A oder eine Aryl- oder Heteroaryleinheit davon definierten Substituenten tragen kann; und

20 mit der Maßgabe, dass kein aliphatisches Kohlenstoffatom an mehr als ein Stickstoff- oder Sauerstoffatom gebunden ist, außer als Teil eines cyclischen Ketals, oder wenn das Stickstoffatom eine Carbonylgruppe trägt; oder

25 für eine saure oder basische Verbindung der Formel I, ein pharmazeutisch verträgliches Salz davon.

2. Verbindung nach Anspruch 1, wobei:

30 R⁰ eine Methyl-, Ethyl-, Propyl-, Isopropyl- oder Isobutylgruppe ist;

R ein Wasserstoffatom, A.X.CO-, A.X.CJ-, D.W.SO₂- oder G ist, worin

W eine direkte Bindung oder eine Iminogruppe ist;

G ein (1-3C)Alkyl-, Aryl(1-3C)alkyl- oder Heteroaryl(1-2C)alkylrest ist, der einen oder mehrere in Anspruch 1 für G oder einen Teil davon definierte Substituenten tragen kann;

35 (1-6C)Alkyl eine Methyl-, Ethyl-, Propyl-, Isopropyl-, Butyl-, Isobutyl-, tert.-Butyl-, Pentyl-, 3-Methylbutyl-, 1-Ethylpropyl-, Hexyl-, oder 4-Methylpentylgruppe ist; (3-6C)Cycloalkyl eine Cyclopropyl-, Cyclopentyl- oder Cyclohexylgruppe ist; der (1-3C)Alkylanteil des (3-6C)Cycloalkyl(1-3 C)alkyl-, Aryl(1-3C)alkyl- oder Heteroaryl(1-3C)alkylrestes eine Methylen-, Ethylen- oder Trimethylengruppe ist; C(1-4)Alkyl eine Methyl-, Ethyl-, Propyl-, Isopropyl-, Butyl-, Isobutyl- oder tert.-Butylgruppe ist; Acyloxy mit 1 bis 5 Kohlenstoffatomen eine Acetoxygruppe ist; (1-4C)Alkoxy eine Methoxy-, Ethoxy-, Propoxy-, Isopropoxy- oder tert.-Butoxygruppe ist; ein

40 Halogenatom Brom, Chlor oder Fluor ist; A.X- zusammengekommen eine 2,2,2-Trifluorethoxygruppe ist; COORa eine Carboxy- oder Methoxycarbonylgruppe ist; CONRbRc eine Carbamoyl- oder N,N-Dimethylcarbamoylgruppe ist; NRgCOR² eine Trifluoracetylaminogruppe ist; CONRdSO₂R¹ eine N-Phenylsulfonylcarbamoyl- oder N-(4-Chlorphenylsulfonyl)carbamoylgruppe ist; A.X- eine Tris(hydroxymethyl)methylamino-, Tris(acetoxymethyl)methylamino- oder 2,2-Bis(hydroxymethyl)propoxygruppe ist; D.W- zusammengekommen eine 2,2,2-Trifluorethylamino- oder 3,3,3-Trifluorpropylgruppe ist; und

45 R⁶ eine Isopropyl-, Cyclopentyl-, Cyclohexyl-, Phenyl-, Furyl-, Thienyl- oder Pyridylgruppe ist, wobei eine Phenylgruppe oder ein Heteroarylrest einen oder zwei Substituenten nach Anspruch 1 tragen kann.

50 3. Verbindung nach Anspruch 1 oder 2, wobei R⁰ Isopropyl ist; R ein Wasserstoffatom, A.X.CO-, A.X.CJ-, D.W.SO₂- oder G ist, wobei A.X- zusammengekommen eine 2,2,2-Trifluorethoxygruppe ist; J ein Sauerstoffatom ist; X eine direkte Bindung, eine Imino- oder Oxygruppe ist; A eine Methyl-, Ethyl-, Phenyl-, Benzyl-, Phenethyl-, Pyridyl-, Thienyl-, 5-Tetrazolyl-, Thiazolyl-, Pyridylmethyl-, Thenyl-, 5-Tetrazolylmethyl-, 2-(Pyridyl)ethyl-, 2-(Thienyl)ethyl- oder 2-(Thiazolyl)ethylgruppe ist, wobei die Phenyl- oder Heteroarylgruppe ein oder zwei Halogenatome oder

55 Methylgruppen tragen kann, oder ferner wobei der Rest A einen Substituenten tragen kann, ausgewählt aus einer Hydroxy-, Methoxy-, tert.-Butoxy-, Acetoxy-, Pivaloyloxy-, Carboxy-, Methoxycarbonyl-, Ethoxycarbonyl-, Carbamoyl-, Dimethylcarbamoyl-, 2-(Dimethylamino)ethoxycarbonyl-, Cyano-, Methylsulfonyl-, Phenylsulfonyl-, N-Methylsulfonylcarbamoyl-, N-Phenylsulfonylcarbamoyl-, Amino-, Dimethylamino-, Oxazolidin-2-on-3-yl-, Acetyl-

- amino-, Trifluoracetyl-amino-, Ureldo-, Methylsulfonyl-, Sulfamoyl-, Dimethylphosphoryl- oder Diethylphosphorylgruppe; D,W-zusammengenommen eine 2,2,2-Trifluorethylamino- oder 3,3,3-Trifluorpropylgruppe ist; D eine Methyl-, Ethyl-, Isopropyl-, tert.-Butyl-, Cyclohexyl-, Phenyl-, Benzyl-, Phenethyl-, Pyridyl-, Thienyl-, 5-Tetrazolyl-, Thiazolyl-, Chinolinyl-, Pyridylmethyl-, Thienyl-, 5-Tetrazolylmethyl-, 2-(Pyridyl)ethyl-, 2-(Thienyl)ethyl- oder 2-(Thiazolyl)ethylgruppe ist, wobei die Phenyl- oder Heteroarylgruppe ein oder zwei Halogenatome oder Methylgruppen tragen kann und wobei der Rest D einen Substituenten tragen kann, ausgewählt aus einer Hydroxy-, Methoxy-, tert.-Butoxy-, Acetoxy-, Pivaloyloxy-, Carboxy-, Methoxycarbonyl-, Ethoxycarbonyl-, Carbamoyl-, Dimethylcarbamoyl-, 2-(Dimethylamino)ethoxycarbonyl-, Cyano-, Methylsulfonyl-, Phenylsulfonyl-, N-Methylsulfonylcarbamoyl-, N-Phenylsulfonylcarbamoyl-, N-(4-Chlorphenylsulfonyl)carbamoyl-, Methylsulfonylamino-, Amino-, Dimethylamino-, Oxazolidin-2-on-3-yl-, Acetyl-amino-, Trifluoracetyl-amino-, Ureldo-, Methylsulfonyl-, Sulfamoyl-, Dimethylphosphoryl- oder Diethylphosphorylgruppe; G eine Methyl-, Ethyl-, Benzyl-, Phenethyl-, Pyridyl-, Pyridylmethyl-, Thienyl-, 5-Tetrazolylmethyl-, oder 2-(Pyridyl)ethylgruppe ist, wobei ein Alkylkohlenstoffatom eine Oxogruppe tragen kann und wobei die Phenyl- oder Heteroarylgruppe ein oder zwei Halogenatome oder Methylgruppen tragen kann und wobei ferner der Rest G einen Substituenten tragen kann, ausgewählt aus einer Hydroxy-, Methoxy-, Acetoxy-, Carboxy-, Methoxycarbonyl-, Ethoxycarbonyl-, Carbamoyl-, Dimethylcarbamoyl-, Phenylcarbamoyl-, Pyridylcarbamoyl-, Methylsulfonylamino-, Amino-, Dimethylamino-, Acetyl-amino-, Nicotinoylamino- oder Trifluoracetyl-amino-
gruppe.
4. Verbindung nach Anspruch 1, 2 oder 3, wobei R ein Wasserstoffatom, eine Formyl-, Trifluoracetyl-, 2,2,2-Trifluorethoxycarbonyl-, Hydroxyoxalyl-, Methoxycarbonyl-, Ethoxycarbonyl-, Isopropoxycarbonyl-, 2-Methoxyethoxycarbonyl-, 4-Fluorphenoxycarbonyl-, 4-Bromphenoxycarbonyl-, 4-Methoxyphenoxycarbonyl-, Benzoyloxycarbonyl-, 4-Fluorbenzoyloxycarbonyl-, 4-Pyridylmethoxycarbonyl-, 3-Methylpyrid-4-ylmethoxycarbonyl-, 2,6-Dimethylpyrid-4-ylmethoxycarbonyl-, 2-Pyridylmethoxycarbonyl-, 6-Methylpyrid-2-ylmethoxycarbonyl-, 2-Dimethylaminoethoxycarbonyl-, Acetyl-, Carbamoylmethylaminocarbonyl-, 4-(N-Phenylsulfonylcarbamoyl)phenylacetyl-, Methylthiocarbonyl-, Sulfo-, Aminosulfonyl-, Dimethylaminosulfonyl-, 2,2,2-Trifluorethylaminosulfonyl-, 3,3,3-Trifluorethylsulfonyl-, Trifluormethylsulfonyl-, Methylsulfonyl- (die einen Methoxycarbonyl-, Carboxy- oder Ethylsulfonyl-Substituenten tragen kann), Methylaminosulfonyl-, Isopropylaminosulfonyl-, Butylsulfonyl-, Butylaminosulfonyl-, tert.-Butylaminosulfonyl-, Cyclohexylaminosulfonyl-, Phenylsulfonyl-, (worin die Phenylgruppe ein Chloratom, einen Nitro-, Amino-, Formylamino-, Acetyl-amino-, Trifluoracetyl-amino-, Methoxy-, Carboxy-, N-(4-Chlorphenylsulfonyl)carbamoyl- oder Methylsulfonylamino-Substituenten an der Stellung 3 oder 4 tragen kann), Anilino-, Pyridylsulfonyl-, Chinolinylsulfonyl-, Benzylsulfonyl- (worin der Phenylring einen Nitro- oder Aminosubstituenten an der Stellung 3 oder 4 tragen kann), Pyridylmethylsulfonyl-, 2-(Pyridyl)ethylsulfonyl-, Benzylaminosulfonyl-, Methyl-, Ethyl-, Benzyl-, Phenethyl- oder Pyridylmethylgruppe ist.
5. Verbindung nach Anspruch 4, worin der Rest R ein Wasserstoffatom, eine Formyl-, Trifluoracetyl-, 2,2,2-Trifluorethoxycarbonyl-, Methoxycarbonyl-, Ethoxycarbonyl-, Isopropoxycarbonyl-, 2-Methoxyethoxycarbonyl-, 4-Bromphenoxycarbonyl-, Benzoyloxycarbonyl-, 2,6-Dimethylpyrid-4-ylmethoxycarbonyl-, Methylthiocarbonyl-, tert.-Butylaminosulfonyl-, 6-Acetylaminophenylsulfonyl-, 4-(N-(4-Chlorphenylsulfonyl)carbamoyl)phenylsulfonyl-, Benzylsulfonyl-, Benzylaminosulfonyl- oder Ethylgruppe ist.
6. Verbindung nach einem der Ansprüche 1 bis 5, wobei R⁶ eine 2-Furyl-, 2-Thienyl-, 3-Pyridyl- oder Phenylgruppe ist, worin die Phenylgruppe einen oder zwei Halogen-, Trifluormethyl-, Methyl-, Hydroxy-, Methoxy-, tert.-Butoxy-, Methoxycarbonyl- oder Carboxy-substituenten tragen kann.
7. Verbindung nach Anspruch 1, worin R⁰ eine Isopropylgruppe ist, R ein Wasserstoffatom, eine Formyl-, 2,2,2-Trifluorethoxycarbonyl-, Isopropoxycarbonyl-, Methylthiocarbonyl- oder Ethylgruppe ist und R⁶ eine 2-Furyl-, 2-Thienyl-, 3-Pyridyl- oder Phenylgruppe ist, worin die Phenylgruppe einen oder zwei Halogen-, Trifluormethyl-, Methyl-, Hydroxy-, Methoxy-, tert.-Butoxy-, Methoxycarbonyl- oder Carboxy-Substituenten tragen kann.
8. Verbindung nach Anspruch 1, ausgewählt aus:

- (a) 2-[5-Amino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid,
 (b) 2-[5-Amino-2-(4-fluorphenyl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid,
 (c) 2-[5-Isopropoxycarbonylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid,
 (d) 2-[5-Ethylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid.

pyl)acetamid,

(e) 2-[6-Oxo-2-phenyl-5-(2,2,2-trifluoroethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid,

(f) 2-(5-Methylthiocarbonylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid,

(g) 2-[2-(4-Fluorophenyl)-5-methylthiocarbonylamino-6-oxo-1,2-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid,

(h) 2-[6-Oxo-2-(2-thienyl)-5-(2,2,2-trifluoroethoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid,

(i) 2-[5-Formylamino-6-oxo-2-(2-thienyl)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid, und

(j) 2-(5-Formylamino-6-oxo-2-phenyl-1,6-dihydro-1-pyrimidinyl)-N-(3,3,3-trifluor-1-isopropyl-2-oxopropyl)acetamid.


9. Salz nach Anspruch 1, ausgewählt aus

(a) für eine saure Verbindung der Formel I, einem Alkalimetallsalz, einem Erdalkalimetallsalz, einem Aluminiumsalz, einem Ammoniumsalz oder einem Salz aus einer organischen Base, die ein pharmazeutisch verträgliches Kation liefert; und

(b) für eine basische Verbindung der Formel I, einem Säureadditionssalz, hergestellt mit einer Säure, die ein pharmazeutisch verträgliches Anion bereitstellt.

10. Verfahren zur Herstellung einer Verbindung der Formel I oder eines pharmazeutisch verträglichen Salzes davon nach einem der Ansprüche 1 bis 9, gekennzeichnet durch:

(A) Oxidieren eines entsprechenden Alkohols der Formel II;



II

(B) für eine Verbindung der Formel I, die einen Rest N-H enthält, Entfernen der Stickstoff-Schutzgruppe einer entsprechenden Verbindung, die eine herkömmliche Stickstoff-Schutzgruppe trägt, durch ein herkömmliches Verfahren;

(C) für eine Verbindung der Formel I, worin R ein Acylrest ist, Acylieren eines entsprechenden Amins der Formel I, wobei R ein Wasserstoffatom ist;

(D) für eine Verbindung der Formel I, worin R eine Sulfonylgruppe ist, Sulfonieren eines entsprechenden Amins der Formel I, wobei R ein Wasserstoffatom ist, mit einer entsprechenden Sulfonsäure der Formel D.W.SO₂.OH oder einem aktivierten Derivat davon;

(E) für eine Verbindung der Formel I, worin R ein Rest G ist, Substitution des Restes L einer entsprechenden Verbindung der Formel G-L, wobei L eine herkömmliche Abgangsgruppe ist, mit einem entsprechenden Amin der Formel I, wobei R ein Wasserstoffatom ist, gegebenenfalls mit einem herkömmlichen Katalysator;

(F) für eine Verbindung der Formel I, die einen Hydroxysubstituenten an einem Aryl- oder Heteroarylrest trägt, Spalten des Alkylethers oder Acyloxyesters einer entsprechenden Verbindung der Formel I, die einen (1-4C)Alkoxy- oder Acyloxy substituenten mit 1 bis 5 Kohlenstoffatomen an einem Aryl- oder Heteroarylrest trägt;

(G) für eine Verbindung der Formel I, die einen Rest der Formel COORa trägt, in dem Ra ein Wasserstoffatom ist, Zersetzen der Estergruppe eines entsprechenden Esters, hergestellt mit einer geeignet entfernten Säureschutzgruppe;

(H) für eine Verbindung der Formel I, die eine Einheit der Formel COORa, CONRbRc, COO(CH₂)₂NReRf oder CONRdSO₂R¹ trägt, Acylieren einer entsprechenden Verbindung der Formel HORa, HNRbRc, HO(CH₂)₂NReRf oder HNRdSO₂R¹ mit einer entsprechenden Säure der Formel I, die eine Einheit der Formel COORa trägt, worin Ra ein Wasserstoffatom darstellt, oder einem aktivierten Derivat davon;


(I) für eine Verbindung der Formel I, die einen Acyloxyrest mit 1 bis 5 Kohlenstoffatomen oder einen Rest der Formel NRgCHO , NRgCOR^2 , NRgCOOR^2 , NRhCQNRiRj oder NRkSO_2R^3 trägt, Acylieren oder Sulfonieren einer entsprechenden Verbindung der Formel I, die eine Hydroxygruppe oder eine Aminogruppe der Formel NHRg , NHRh oder NHRk trägt, mit einem aktivierten Derivat einer entsprechenden Säure der Formel HOCHO , HOCOR^2 , HOCOOR^2 , HOCQNRiRj (einschließlich eines Isocyanates oder Isothiocyanates) bzw. HOSO_2R^3 , unter Verwendung eines herkömmlichen Verfahrens;

(J) für eine Verbindung der Formel I, die einen Heteroaryl-N-oxidrest trägt, Oxidieren einer entsprechenden Verbindung der Formel I, die einen Heteroarylrest trägt, mit einem herkömmlichen Oxidationsmittel; oder

(K) für eine Verbindung der Formel I, die eine primäre Aminogruppe trägt, Reduktion einer entsprechenden Verbindung, die eine Nitrogruppe trägt, unter Verwendung eines herkömmlichen Reduktionsverfahrens; und danach für jedes der vorstehenden Verfahren, wenn ein pharmazeutisch verträgliches Salz einer sauren oder basischen Verbindung der Formel I benötigt wird, Umsetzen der sauren oder basischen Form einer solchen Verbindung der Formel I mit einer Base oder Säure, die ein physiologisch verträgliches Gegenion bereitstellt, oder durch jedes andere herkömmliche Verfahren; und


wobei R , R^0 , R^6 , D , W , G , Ra-Rk , $\text{R}^1\text{-R}^3$ und Q , wenn nicht eingehender beschrieben, die in einem der Ansprüche 1 bis 9 definierte Bedeutung haben.

11. Verbindung der Formel II,



wobei R , R^0 und R^6 die in Anspruch 1 definierte Bedeutung haben, oder ein Salz davon.

12. Verbindung der Formel Vb,




worin R die für G in Anspruch 1 definierte Bedeutung hat, R^0 und R^6 die in Anspruch 1 definierte Bedeutung haben, und Rx ein Rest ist, der eine primäre Aminogruppe schützt und zur Substitution aktiviert, oder ein Salz davon.

13. Arzneimittel, umfassend eine Verbindung nach Anspruch 1 oder ein pharmazeutisch verträgliches Salz davon und ein pharmazeutisch verträgliches Verdünnungsmittel oder einen pharmazeutisch verträglichen Träger.

Revendications

1. Composé de formule I



I

où

R⁰ est alkyle en C₁₋₅;

R est l'hydrogène ; ou bien

R est un groupe acyle de formule A.X.CO- où A.X-, considéré globalement, est hydrogène, trifluorométhyle, 2,2,2-trifluoroéthoxy, amino, méthoxyamino, 2,2,2-trifluoroéthylamino, RbRcN.O-, RaOCONH-, R¹SO₂NH-, RaOCO-, RbRcNCO- ou RaCO- ; ou bien

R est un groupe acyle de formule A.X.CJ- où

J est l'oxygène ou le soufre ;

X est une liaison directe, imino, oxy ou thio ; et

A est tel que défini ci-dessous ou bien

A est tétrahydropyran-4-yle, 1-méthylpipérid-4-yle, ou 5-méthyl-1,3-dioxacyclohex-5-ylméthyle ; ou bien

R est un groupe sulfonyl de formule D.W.SO₂- où D.W-, considéré globalement, est hydroxyle, amino, dialkylamino en C₁₋₄, 2,2,2-trifluoroéthylamino, 2,2,2-trifluoroéthyle, 3,3,3-trifluoropropyle ou trifluorométhyle ; ou bien

W est une liaison directe, imino, carbonylimino, oxycarboxylimino ou iminocarbonylimino ; et

D est tel que défini ci-dessous ; ou bien

R est un groupe alkyle, aryle ou hétéroaryle G tel que défini ci-dessous ;

le groupe A, D ou G est alkyle en C₁₋₆, cycloalkyle en C₃₋₆, cycloalkyle en C₃₋₆-alkyle en C₁₋₃, aryle, aryl-alkyle en C₁₋₃, hétéroaryle ou hétéroaryl-alkyle en C₁₋₃ où une entité aryle est choisie parmi phényle, indényle et naphthyle, et une entité hétéroaryle est choisie parmi furyle, imidazolyle, tétrazolyle, pyridyle, pyridyl-N-oxyde, thiényl, pyrimidinyle, pyrimidinyl-N-oxyde, indolyle, quinolyle et quinolyl-N-oxyde, et où une entité aryle ou hétéroaryle peut porter un ou plusieurs groupes halogéno, nitro, méthyle ou trifluorométhyle et où, de plus, le groupe A, D ou G peut porter un ou plusieurs substituants choisis dans un groupe consistant en hydroxyle, alcoxy en C₁₋₄, acyloxy contenant 1 à 5 atomes de carbone, COORa, CONRbRc, COO(CH₂)₂NReRf, cyano, SO₂R¹, CONRdSO₂R¹, NReRf, NRgCHO, NRgCOR², NRgCOOR², NRhCQNRiRj, NRkSO₂R³, SO₂NRlRm, SO₂NRnCOR⁴ et P(O)(ORa)₂ où

Q est l'oxygène ou le soufre ;

Ra-Rn sont indépendamment l'hydrogène, benzyle ou alkyle en C₁₋₄ ; ou bien, indépendamment, un groupe NRbRc, NReRf, NRiRj ou NRlRm est un radical cyclique choisi dans un groupe consistant en 1-pyrrolidinyle, pipéridino, morpholino ou 1-pipérazinyle qui peut porter un substituant alkyle en C₁₋₄ en position 4 ; ou bien, indépendamment, un groupe NReRf est un radical cyclique choisi dans un groupe consistant en 2-pyrrolidinon-1-yle, succinimido, oxazolidin-2-on-3-yle, 2-benzoxazolinon-3-yle, phthalimido et cis-hexahydrophthalimido ; et

R¹-R⁴ sont indépendamment trifluorométhyle, alkyle en C₁₋₆, cycloalkyle en C₃₋₆, aryle ou hétéroaryle où une entité aryle est choisie parmi phényle, indényle et naphthyle, et une entité hétéroaryle est choisie parmi furyle, imidazolyle, tétrazolyle, pyridyle, pyridyl-N-oxyde, thiényl, pyrimidinyle, pyrimidinyl-N-oxyde, indolyle, quinolyle et quinolyl-N-oxyde, et où une entité aryle ou hétéroaryle peut porter un ou plusieurs substituants choisis dans un groupe consistant en alkyle en C₁₋₄, hydroxyle, alcoxy en C₁₋₄, halogéno et trifluorométhyle ;

R⁶ est alkyle en C₁₋₅ qui n'a pas de carbone tertiaire, cycloalkyle en C₃₋₇, aryle ou hétéroaryle, où une entité aryle est choisie parmi phényle, indényle et naphthyle, et une entité hétéroaryle est choisie parmi furyle, imidazolyle, tétrazolyle, pyridyle, pyridyl-N-oxyde, thiényl, pyrimidinyle, pyrimidinyl-N-oxyde, indolyle, quinolyle et quinolyl-N-oxyde, et où une entité aryle ou hétéroaryle peut porter indépendamment un ou plusieurs des substituants définis pour le groupe A ou une entité aryle ou hétéroaryle de celui-ci ; et

à condition qu'aucun carbone aliphatique ne soit lié à plus d'un azote ou d'un oxygène, sauf s'il fait partie d'un cétal cyclique ou si l'azote porte un groupe carbonyle ; ou bien pour un composé de formule I qui est acide ou basique, sel pharmaceutiquement acceptable de celui-ci.

2. Composé selon la revendication 1, où :

R^0 est méthyle, éthyle, propyle, isopropyle ou isobutyle ;

R est l'hydrogène, A.X.CO-, A.X.CJ-, D.W.SO₂- ou G où

W est une liaison directe ou imino ;

G est alkyle en C₁₋₃, aryle-alkyle en C₁₋₃ ou hétéroaryl-alkyle en C₁₋₂ qui peut porter un ou plusieurs substituants tels que définis dans la revendication 1 pour G ou une partie de celui-ci ;

alkyle en C₁₋₆ est méthyle, éthyle, propyle, isopropyle, butyle, isobutyle, t-butyle, pentyle, 3-méthylbutyle, 1-éthylpropyle, hexyle ou 4-méthylpentyle ; cycloalkyle en C₃₋₆ est cyclopropyle, cyclopentyle ou cyclohexyle ; la partie alkyle en C₁₋₃ de cycloalkyle en C₃₋₆-alkyle en C₁₋₃, arylalkyle en C₁₋₃ ou hétéroaryl-alkyle en C₁₋₃ est méthylène, éthylène ou triméthylène ; alkyle en C₁₋₄ est méthyle, éthyle, propyle, isopropyle, butyle, isobutyle ou t-butyle ; acyloxy contenant 1 à 5 atomes de carbone est acétoxy ; alcoxy en C₁₋₄ est méthoxy, éthoxy, propoxy, isopropoxy ou t-butoxy ; halogéno est bromo, chloro ou fluoro-, A.X.-, considéré globalement, est 2,2,2-trifluoroéthoxy ; COORa est carboxyle ou méthoxycarbonate ; CONRbRc est carbamoyle ou N,N-diméthylcarbamoyle ; NRgCOR² est trifluoroacétylamino ; CONRdSO₂R¹ est N-phénylsulfonylcarbamoyle ou N-(4-chlorophénylsulfonyl)carbamoyle ; A.X.- est tris(hydroxyméthyl)méthylamino, tris(acétoxyméthyl)méthylamino ou 2,2-bis(hydroxyméthyl)propoxy ; D.W.-, considéré globalement, est 2,2,2-trifluoroéthylamino ou 3,3,3-trifluoropropyle ; et


R⁶ est isopropyle, cyclopentyle, cyclohexyle, phényle, furyle, thiényle ou pyridyle où un phényle ou hétéroaryle peut porter un ou deux substituants tels que définis dans la revendication 1.

3. Composé selon la revendication 1 ou 2 où R⁰ est isopropyle, R est l'hydrogène, A.X.CO-, A.X.CJ-, D.W.SO₂- ou G où A.X.-, considéré globalement, est 2,2,2-trifluoroéthoxy, J est l'oxygène, X est une liaison directe, imino ou oxy, A est méthyle, éthyle, phényle, benzyle, phénéthyle, pyridyle, thiényle, 5-tétrazolyloyle, thiazolyloyle, pyridylméthyle, thényle, 5-tétrazolylméthyle, 2-(pyridyl)éthyle, 2-(thiényl)éthyle ou 2-(thiazolyl)éthyle où le groupe phényle ou hétéroaryle peut porter un ou deux groupes halogéno ou méthyle et où, en outre, le groupe A pour porter un substituant choisi parmi hydroxyle, méthoxy, t-butoxy, acétoxy, pivaloyloxy, carboxyle, méthoxycarbonate, éthoxycarbonate, carbamoyle, diméthylcarbamoyle, 2-(diméthylamino)éthoxycarbonate, cyano, méthylsulfonyl, phénylsulfonyl, N-méthylsulfonylcarbamoyle, N-phénylsulfonylcarbamoyle, amino, diméthylamino, oxazolidin-2-on-3-yle, acétylamino, trifluoroacétylamino, uréido, méthylsulfonyl, sulfamoyle, diméthylphosphoryle ou diéthylphosphoryle ; D.W.-, considéré globalement, est 2,2,2-trifluoroéthylamino ou 3,3,3-trifluoropropyle ; D est méthyle, éthyle, isopropyle, tert-butyle, cyclohexyle, phényle, benzyle, phénéthyle, pyridyle, thiényle, 5-tétrazolyloyle, thiazolyloyle, pyridylméthyle, thényle, 5-tétrazolylméthyle, 2-(pyridyl)éthyle, 2-(thiényl)éthyle ou 2-(thiazolyl)éthyle où le groupe phényle ou hétéroaryle peut porter un ou deux groupes halogéno ou méthyle et où, en outre, le groupe D peut porter un substituant choisi parmi hydroxyle, méthoxy, t-butoxy, acétoxy, pivaloyloxy, carboxyle, méthoxycarbonate, éthoxycarbonate, carbamoyle, diméthylcarbamoyle, 2-(diméthylamino)éthoxycarbonate, cyano, méthylsulfonyl, phénylsulfonyl, N-méthylsulfonylcarbamoyle, N-phénylsulfonylcarbamoyle, N-(4-chlorophénylsulfonyl)carbamoyle, méthylsulfonylamino, amino, diméthylamino, oxazolidin-2-on-3-yle, acétylamino, trifluoroacétylamino, uréido, méthylsulfonyl, sulfamoyle, diméthylphosphoryle ou diéthylphosphoryle ; G est méthyle, éthyle, benzyle, phénéthyle, pyridyle, pyridylméthyle, thényle, 5-tétrazolylméthyle, ou 2-(pyridyl)éthyle, où un carbone d'alkyle peut porter un groupe oxo et où le groupe phényle ou hétéroaryle peut porter un ou deux groupes halogéno ou méthyle et où, en outre, le groupe G peut porter un substituant choisi parmi hydroxyle, méthoxy, acétoxy, carboxyle, méthoxycarbonate, éthoxycarbonate, carbamoyle, diméthylcarbamoyle, phénylcarbamoyle, pyridylcarbamoyle, méthylsulfonylamino, amino, diméthylamino, acétylamino, nicotinoylamino, ou trifluoroacétylamino.

4. Composé selon la revendication 1, 2 ou 3, où R est hydrogène, formyle, trifluoroacétyle, 2,2,2-trifluoroéthoxycarbonate, hydroxyoxalyle, méthoxycarbonate, éthoxycarbonate, isopropoxycarbonate, 2-méthoxyéthoxycarbonate, 4-fluorophénoxycarbonate, 4-bromophénoxycarbonate, 4-méthoxyphénoxycarbonate, benzyloxycarbonate, 4-fluorobenzyloxycarbonate, 4-pyridylméthoxycarbonate, 3-méthylpyrid-4-ylméthoxycarbonate, 2,6-diméthylpyrid-4-ylméthoxycarbonate, 2-pyridylméthoxycarbonate, 6-méthylpyrid-2-ylméthoxycarbonate, 2-diméthylaminoéthoxycarbonate, acétyle, carbamoylméthylaminocarbonate, 4-(N-phénylsulfonylcarbamoyl)phénylacétyle, méthylthiocarbonate, sulfo, aminosulfonyl, diméthylaminosulfonyl, 2,2,2-trifluoroéthylaminosulfonyl, 3,3,3-trifluoroéthylsulfonyl, trifluorométhylsulfonyl, méthylsulfonyl (qui peut porter un substituant méthoxycarbonate, carboxyle ou éthylsulfonyl), méthylaminosulfonyl, isopropylaminosulfonyl, butylsulfonyl, butylaminosulfonyl, tert-butylaminosulfonyl, cyclohexylaminosulfonyl, phénylsulfonyl (où le phényle peut porter un substituant chloro, nitro, amino, formylamino, acétylamino, trifluoroacétylamino, méthoxy, carboxyle, N-(4-chlorophénylsulfonyl)carbamoyle ou méthylsulfonylamino en position 3 ou 4), anilino, pyridylsulfonyl, quinolinylsulfonyl, benzylsulfonyl (où le cycle phényle peut porter un substituant nitro ou amino en position 3 ou 4), pyridylméthylsulfonyl, 2-

(pyridyl)éthylsulfonyle, benzylaminosulfonyle, méthyle, éthyle, benzyle, phénéthyle ou pyridylméthyle.

5. Composé selon la revendication 4, où R est l'hydrogène, formyle, trifluoroacétyle, 2,2,2-trifluoroéthoxycarbonyle, méthoxycarbonyle, éthoxycarbonyle, isopropoxycarbonyle, 2-méthoxyéthoxycarbonyle, 4-bromophénoxy-carbonyle, benzyloxycarbonyle, 2,6-diméthylpyrid-4-ylméthoxycarbonyle, méthylthiocarbonyle tert-butylaminosulfonyle, 4-acétylamino-phénylsulfonyle, 4-(N-(4-chlorophénylsulfonyl)carbamoyl)phénylsulfonyle, benzylsulfonyle, benzylaminosulfonyle ou éthyle.
6. Composé selon l'une quelconque des revendications 1 à 5, où R⁶ est 2-furyle, 2-thiényle, 3-pyridyle ou phényle où le phényle peut porter un ou deux substituants halogéno, trifluorométhyle, méthyle, hydroxyle, méthoxy, tert-butoxy, méthoxycarbonyle ou carboxyle.
7. Composé selon la revendication 1, où R⁰ est isopropyle, R est l'hydrogène, formyle, 2,2,2-trifluoroéthoxycarbonyle, isopropoxycarbonyle, méthylthiocarbonyle ou éthyle, et R⁶ est 2-furyle, 2-thiényle, 3-pyridyle ou phényle où le phényle peut porter un ou deux substituants halogéno, trifluorométhyle, méthyle, hydroxyle, méthoxy, tert-butoxy, méthoxycarbonyle ou carboxyle.
8. Composé selon la revendication 1, choisi parmi :
 - (a) le 2-[5-amino-6-oxo-2-(2-thiényle)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (b) le 2-[5-amino-2-(4-fluorophényl)-6-oxo-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (c) le 2-[5-isopropoxycarbonylamino-6-oxo-2-(2-thiényle)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (d) le 2-[5-éthylamino-6-oxo-2-(2-thiényle)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (e) le 2-[6-oxo-2-phényl-5-(2,2,2-trifluoroéthoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (f) le 2-[5-méthylthiocarbonylamino-6-oxo-2-phényl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (g) le 2-[2-(4-fluorophényl)-5-méthylthiocarbonylamino-6-oxo-1,2-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (h) le 2-[6-oxo-2-(2-thiényle)-5-(2,2,2-trifluoroéthoxycarbonylamino)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide,
 - (i) le 2-[5-formylamino-6-oxo-2-(2-thiényle)-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide et
 - (j) le 2-[5-formylamino-6-oxo-2-phényl-1,6-dihydro-1-pyrimidinyl]-N-(3,3,3-trifluoro-1-isopropyl-2-oxopropyl)acétamide.
9. Sel selon la revendication 1, choisi parmi
 - (a) pour un composé acide de formule I, un sel de métal alcalin, un sel de métal alcalino-terreux, un sel d'aluminium, un sel d'ammonium ou un sel formé à partir d'une base organique qui donne un cation pharmaceutiquement acceptable ; et
 - (b) pour un composé basique de formule I, un sel d'addition d'acide formé avec un acide qui donne un anion pharmaceutiquement acceptable.
10. Procédé de production d'un composé de formule I, ou d'un sel pharmaceutiquement acceptable de celui-ci, selon l'une quelconque des revendications 1 à 9, qui est caractérisé par :
 - (A) l'oxydation d'un alcool de formule II correspondant :



II

(B) pour un composé de formule I qui contient un résidu N-H, le retrait au moyen d'un procédé conventionnel du groupe protecteur d'azote d'un composé correspondant portant un groupe protecteur d'azote conventionnel ;

(C) pour un composé de formule I où R est un groupe acyle, l'acylation d'une amine correspondante de formule I où R est l'hydrogène ;

(D) pour un composé de formule I où R est un groupe sulfonyle, la sulfonylation d'une amine correspondante de formule I où R est l'hydrogène avec un acide sulfonique correspondant de formule D.W.SO₂.OH, ou un dérivé activé de celui-ci ;

(E) pour un composé de formule I où R est un groupe G, la substitution du groupe L d'un composé correspondant de formule G-L, où L est un groupe partant conventionnel, avec une amine correspondante de formule I où R est l'hydrogène, éventuellement au moyen d'un catalyseur conventionnel ;

(F) pour un composé de formule I qui porte un substituant hydroxyle sur un groupe aryle ou hétéroaryle, le clivage de l'alkyléther ou de l'acyloxyester d'un composé correspondant de formule I qui porte un substituant alcoxy en C₁₋₄ ou acyloxy contenant 1 à 5 atomes de carbone sur un groupe aryle ou hétéroaryle ;

(G) pour un composé de formule I qui porte un groupe de formule COORa où Ra est l'hydrogène, la décomposition du groupe ester d'un ester correspondant formé avec un groupe protecteur d'acide retiré de manière conventionnelle ;

(H) pour un composé de formule I portant une entité de formule COORa, CONRbRc, COO(CH₂)₂NReRf ou CONRdSO₂R¹, l'acylation d'un composé correspondant de formule HORa, HNRbRc, HO(CH₂)₂NReRf ou HNRdSO₂R¹ avec un acide correspondant de formule I portant une entité de formule COORa où Ra est l'hydrogène, ou un dérivé activé de celui-ci ;

(I) pour un composé de formule I portant un groupe acyloxy contenant 1 à 5 atomes de carbone ou un groupe de formule NRgCHO, NRgCOR², NRgCOOR², NRhCQNRiRj ou NRkSO₂R³, l'acylation ou la sulfonylation d'un composé correspondant de formule I portant un groupe hydroxyle ou un groupe amino de formule NHRg, NHRh ou NHRk avec un dérivé activé d'un acide correspondant de formule HOCHO, HOCOR², HOCOR², HOCQNRiRj (y compris un isocyanate ou un isothiocyanate) ou HOSO₂R³, respectivement, au moyen d'un procédé conventionnel ;


(J) pour un composé de formule I qui porte un groupe hétéroaryl-N-oxyde, l'oxydation d'un composé correspondant de formule I qui porte un groupe hétéroaryle au moyen d'un oxydant conventionnel ; ou

(K) pour un composé de formule I qui porte un groupe amino primaire, la réduction d'un composé correspondant portant un groupe nitro au moyen d'un procédé réducteur conventionnel ; puis

pour l'un quelconque des protocoles ci-dessus, quand un sel pharmaceutiquement acceptable d'un composé acide ou basique de formule I est nécessaire, la réaction de la forme acide ou basique d'un tel composé de formule I avec une base ou un acide donnant un contre-ion physiologiquement acceptable ou par tout autre protocole conventionnel ; et


où R, R⁰, R⁶, D, W, G, Ra-Rk, R¹-R³ et Q ont la signification définie dans l'une quelconque des revendications 1 à 9, sauf quand ils sont décrits de manière plus particulière.

11. Composé de formule II



10 où R, R⁰ et R⁶ sont définis comme dans la revendication 1, ou sel de celui-ci.

12. Composé de formule Vb,



25 où R a une valeur définie pour G dans la revendication 1, R⁰ et R⁶ sont définis comme dans la revendication 1 et Rx est un groupe qui protège et active un groupe amino primaire en vue d'une substitution, ou sel de celui-ci.

30 13. Composition pharmaceutique comprenant un composé selon la revendication 1, ou un sel pharmaceutiquement acceptable de celui-ci, et un diluant ou support pharmaceutiquement acceptable.